

DL


430.7

T3H24

1846


Wall.

BOOK BINDER,
411 BROADWAY,
ALBANY.



1902

U.S.N.M.





Ex libris

William Healey Dall.

W. H. DALL

12/1






I Hanley

Division of Mollusks
Sectional Library







716
430.7
T3/H24
1846
Moll.

381

A

MONOGRAPH OF THE GENUS

Division of Mollusks
Sectional Library

TELLINA.

BY

SYLVANUS HANLEY, B. A. F. L. S. &c.

EXTRACTED FROM THE FIRST VOLUME OF THE
"THESAURUS CONCHYLIIORUM."



LONDON:
JANUARY 3, 1846.

I

MONOGRAPH OF THE GENUS

TRELLINA

STANLEY HANLEY, B.A. F.R.S. &c.

EXTRACTED FROM THE FIRST VOLUME OF THE
"TRELLINA" COMPANION




LONDON:

JANUARY 8, 1908







Just Published, in Imperial 8vo., price Coloured, 11. 5s. Plain 16s.

PART VI.

OF THE

THESAURUS CONCHYLIIORUM;

EDITED AND ILLUSTRATED BY

G. B. SOWERBY, JUN. F.L.S.

EACH PART contains one or more MONOGRAPHS of Genera of Recent Shells, including a description and figure of each species and variety known up to the time of publication, with its locality and essential synonyms. The work, when completed, will form a CONCHOLOGICAL LIBRARY, sufficiently comprehensive to render all other works on the subject unnecessary to collectors.

The Proprietor is happy to state that the first six Parts now before the public, have given universal satisfaction; and he will endeavour to deserve a continuance of the approbation which has been freely expressed.

The price of each Part is calculated at the rate of about one shilling for twelve figures.

Parts 1, 2, 3, 4, 5, and 6, contain Monographs of the Genera HELICINA, PUPINA, ROSTELLARIA, APORRHAI, STRUTHIOLARIA, STROMBUS, PECTEN, PTEROCERAS, HINNITES, LIMA, CYCLOSTOMA, SCALARIA, COLUMBELLA, TEREBRA, VOLUTA, TELLINA (including TELLINIDES), and LINGULA, illustrated by nearly 1,800 figures.

Monographs of the following Genera are now in preparation.

TEREBRATULA and the other genera of BRACHIOPODA
by G. B. Sowerby F.L.S.

SPONDYLUS and PLICATULA by G. B. Sowerby, Jun. F.L.S.

CYMBA and MELO, by G. B. Sowerby, F.L.S.

CYCLAS by G. B. Sowerby, F.L.S.

MARGINELLA, by G. B. Sowerby, Jun. F.L.S.

In returning thanks for support already received, the Proprietor begs to mention that such support would have been much more extensive but for a disinclination on the part of some to take the numbers of a periodical publication during its progress. It should, however, be remembered, that *each Part* of the 'Thesaurus Conchyliorum' is

COMPLETE IN ITSELF,

independently of the general design which it assists in carrying out. Each Monograph, containing all the information that can be collected respecting the Species of a Genus, might, with propriety, be published separately, and under a distinct title.

The Proprietor may further observe; that it will be his aim to arrange for the completion of Families as speedily as circumstances will permit. Thus, one Family is completed by the already published Monographs of the genera Helicina, Pupina, and Cyclostoma, which include all *operculated land shells*. Another Family is completed by the genera Rostellaria, Aporrhais, Struthiolaria, Strombus, and Pteroceras. To the already given Monographs

of Pecten and Lima, will shortly be added those of Spondylus and Plicatula; the Family of Volutidæ will be completed in an early Part, by the Genera Cymba and Melo. Part seven will contain, with other genera, all the remaining Brachiopoda.

GEOLOGISTS, who have long needed a Work like the present to enable them to compare their fossil specimens with the recent species of Shells, will find *uncoloured copies* well adapted to that purpose.

The following Learned Societies and individuals are known to have already placed the *Thesaurus Conchyliorum* in their Libraries:—

HIS MAJESTY THE KING OF DENMARK. (Private library.)

HIS GRACE THE DUKE OF NORTHUMBERLAND,

THE MOST NOBLE THE MARQUIS OF NORTHAMPTON.

The Linnean Society of London.	J. Dennison, Esq. Liverpool.
The United Service Institution, London.	Benjamin Butler, Esq. Manilla.
The Edinburgh College Library.	John Bates, Esq. R.N. Sec. Royal Yacht Soc.
Imperial Academy of Sciences, Moscow.	Cowes, Isle of Wight.
Museum of Bremen.	Thos. Norris, Esq.
Imperial Museum of Zoology, Vienna.	Lyttleton T. Powis, Esq. London.
Royal Asiatic Society, Calcutta.	W. Metcalfe, Esq. London.
Royal Museum of the Jardin des Plantes, Paris.	E. Logan, Esq. New York.
Royal Museum of Berlin.	G. Hubbard, Esq.
Philosophical Institution of York.	Wm. Durrant, Esq.
Royal Museum of Stockholm.	Wm. K. Loftus, Esq. Cambridge.
Trinity College, Dublin (coloured copy).	J. Meynell, Esq. York.
Sir David Barclay, Bart.	Lieut. Hankey R.N.
The Hon. Col. Macadam Cathcart.	J. G. Gaskoin, Esq.
M. le Baron Delessert, Paris.	C. I. F. Bunbury, Esq. F.G.S.&c.
Count Keysserling St. Petersburg.	Sylvanus Hanley, Esq. F.L.S.
Lady Buxton.	J. G. Gabriel, Esq.
Miss Brookes.	Wm. Bean, Esq.
Miss Saul.	Edward Broughton, Esq. Edinburgh.
Miss Thompson, Stamford.	J. Miers, Esq. F.R.S. &c.
Dr. Schmidt, Bremen.	W. J. Broderip, Esq. F.R.S. &c.
Dr. J. C. Jay, New York.	R. B. Hinds, Esq. R.N.
Dr. Jonas Hamburg.	M. Largilliert, Rouen.
The Rev. Dr. J. Pye Smith, F.R.S. &c.	Thos. Bell, Esq. F.R.S. &c.
Rev. L. Vernon Harcourt.	M. Petit de la Saussaye, Paris.
Rev. Edward Budge.	M. D'Alcide D'Orbigny, Paris.
Rev. J. F. Stainforth.	Samuel Stutchbury, Esq. Bristol.
Rev. W. Crotch.	Dr. Charles Theodore Menke, Pyrmont, Waldeck, Germany.
Rev. S. C. E. Neville Rolfe.	T. Bridges, Esq. F.R.S. Valparaiso,
Rev. Thomas Image.	James Tennant, Esq. F.G.S.
Rev. E. Rust.	M. Giovanni Michelotti, Turin.
W. W. Benson, Esq. Rockville, Landour (Civil Service).	W. Borrer, Esq. F.R.S. &c.
Hugh Cuming, Esq. F.L.S.	M. Krantz, Berlin.
E.G.L. Gruner, Esq. Bremen	M. Deshayes, Paris.
Isaac Lea, Esq. Philadelphia.	W. Parker Hammond, Esq.
Thomas Lombe Taylor, Esq.	T. Berrington, Esq.
J. D. C. Sowerby, Esq. F.L.S.	R. Campbell, Esq.
R. Shuttleworth, Esq. Berne, Switzerland.	M. Janelle, Paris.

The Proprietor will feel obliged if those Purchasers of the Work whose names are omitted from the above List, would enable him to render it more complete by kindly forwarding their names to the Publisher for insertion; and any who may wish to purchase, or to inspect with a view to purchase Copies, may have them forwarded to any part of Great Britain, FREE OF EXPENSE,

SOWERBY, BOOKSELLER AND DEALER IN MINERALS AND SHELLS,
50, GREAT RUSSELL STREET, BLOOMSBURY.

A MONOGRAPH OF THE GENUS

TELLINA.

By SYLVANUS HANLEY, B.A., F.L.S., ETC.

1847
1

Char. Gen. Testa bivalvis, cujus latus posticum aut plicaturâ irregulari radiante tumescit, aut cardo ejus dentem dentesve duos laterales ostendit; dentes primarii duo, vel in utrâque, vel in alterâ valvulâ; margo ventralis nunquam crenatus; pallii cicatrix maximè sinuata; nymphæ haud prominentes. Ligamentum externum.

There are few genera whose definition is more difficult than that of *Tellina*. The typical species with a strong fold or umbonal ridge and two distinct lateral teeth, cannot indeed be confounded with the shells of any other genus; but the whole of the aberrant division, in which the fold is oftentimes nearly obsolete and the hinge always devoid of lateral teeth, require the strictest examination, so closely do they approximate to *Sanguinolaria* and *Psammobia*. Lamarck himself has included several species of this division in *Psammobia* and *Psammotæa* (the latter genus abolished by the unanimous agreement of all conchologists); but the authority of Deshayes gives weight to my own opinion, of the necessity of their annexation to *Tellina*. The externally projecting nymphal callosities which support the prominent ligament, characters which are universal throughout *Psammobia*, are never present in *Tellina* of this division. I may remark moreover, as a further assistance to the discrimination of those closely allied genera, that the inclination of the line which unites the posterior muscular impression to the parallel scar is *ascending* in those aberrant species which most nearly approach *Psammobia*, whilst it is *descending* in *that genus*. I speak more particularly of *Psammobia*, as there exists in some of its species a fold almost equally profound (but symmetrical) with that of the majority of the *Tellinæ*. This does not exist in *Sanguinolaria*, of which I regard *Rosea* and not *Rugosa* as the type.

Certain *Donaces*, devoid of crenulated margins, might probably

be mistaken for *Tellinæ*, but these being always perfectly equivalve, strong, wedge-shaped, or triangular shells, with a remarkably short prominent ligament, and the anterior side the more compressed, present a combination of characters which are never united in any of those *Tellens* in which the characteristic fold is imperfectly developed. The decided situation of the pallear scar suffices to separate any species from *Lucina*, *Corbis*, or *Astarte*.

~~~~~

### DESCRIPTION.

\* *Two lateral teeth in one of the valves.*

1. *TELLINA ROSTRATA* (pl. lxi. f. 157.) *Lin.*, Syst. 1118; *Lin. Mus. Ulr.* 481; *T. Spengleri*, *Chem.* vol. vi. f. 88, 89; *Wood*, *Gen. Conch.* t. 37. f. 2; *Lam.* no. 8; *Hanl. Des. Cat.* p. 62. t. 4. f. 35; *Enc. Mèth.* t. 287. f. 5.

*T.* testâ elongato-subtrigonâ, angustâ, subinæquivalvi, subæquilaterali, solidâ, convexiusculâ, subflexuosâ, extus intusque candidâ, prope nates roseo pallidè pictâ, concentricè sulcatâ; sulcis confertis, obliquis; margine ventrali subrecto aut convexiusculo; dorsali utrinque subrecto et subæqualiter declivi; areis dorsalibus sublevigatis, squamis tuberculosis, utrinque dentatis; latere antico subacuminato; postico subrostrato, biangulato; costâ umbonali angulatâ; ligamento subinfosso; dentibus lateralibus magnis, remotis.

Elongated-subtriangular, narrow, rather flexuous, almost equilateral, rather inequivalve, solid, moderately convex, white both externally and internally, with a slight tinge of rose-colour near the beaks, closely and obliquely grooved concentrically; ventral edge but little convex; dorsal margins nearly straight, and almost equally sloping; dorsal areas nearly smooth, and environed by scaly tubercles; anterior side subacuminated; posterior biangulated and subrostrated; umbonal ridge angulated; ligament sunken; lateral teeth strong and distant.

Island of Ticao, &c,

This is not the *Rostrata* of Gmelin, Lamarek, nor the majority of writers, but nevertheless, as was surmised by Schumacher from the expression "ano rimâque, subserratis," is the species indicated by Linnæus, as his specimen in the Linnæan Society's cabinet clearly evidences.

2. *TELLINA DESHAYESII* (pl. lvii. f. 66.) *Hanley*, Zool. Proc. 1844.

T. testâ *T. Rostratæ* simillimâ, sed albido-roseâ, et lamellis sub-remotis concentricè ornatâ; margine etiam ventrali convexiore.

Extremely like *T. rostrata*, but of an uniform pale pinkish hue, and the surface ornamented with concentric and rather distant lamellæ; the ventral margin more convex likewise, than in the preceding species.

Red Sea?

3. *TELLINA CUMINGII* (pl. lviii. f. 72.) *Hanley*, Zool. Proc. 1844.

T. testâ elongato-oblongâ, subæquilaterali, solidâ, compressiusculâ, vix nitidâ, albidâ aut flavescente, maculis strigis lineisve brunneis aut fusco-purpurascensibus radiatâ; concentricè sulcatâ, sulcis anticè confertis et subimbricatis, posticè sublamellosis, remotiusculis; margine ventrali convexo, utrinque sursum acclinato; dorsali utrinque subdeclivi, anticè convexiusculo, posticè subrecto, subincurvato, subdentato; latere postico attenuato, rostrato, paululum brevior, ad extremitatem biangulato; costâ umbonali valdè prominente; ligamento inconspicuo; superficie internâ albidâ aut flavidâ; dentibus lateralibus validis, subæquidistantibus.

Elongated-oblong, narrow, subequilateral, solid, rather compressed, scarcely glossy, whitish, radiated with narrow brown or chocolate-brown streaks or spots; concentrically sulcated, sulci fine, becoming elevated posteriorly, strong and coarse on the very prominent umbonal ridge, attenuated in the hollow beyond, and projecting again so as almost to dentate the straight or slightly concave ligamental margin; slopes very gentle; front dorsal area barred with coloured streaks; anterior extremity rounded, but somewhat attenuated; hinder side narrowed, subrostrated, and somewhat biangulated at the extremity; inside white or pale yellowish; lateral teeth strong, and nearly equidistant.

Guacomayo and America.

4. *TELLINA CRUCIGERA* (pl. lviii. f. 78, 79; and pl. lxii. f. 178.) *Lam.*, no. 7; *Hanl. Des. Cat.* p. 61.

T. testâ oblongâ, convexâ, inæquilaterali, nitidiusculâ, albidâ. lineis maculisque fulvo-aurantiis aut roseo-purpureis plerumque radiatâ; (rarius immaculatâ, et radio brevi roseo ad umbones utrinque ornatâ), sulcis exilibus, distinctis, paululum obliquis, concentricè exaratâ; margine ventrali convexo, posticè sursum acclinato; dorsali anticè convexiusculo et vix paululum declivi, posticè

vix convexiusculo satisque declivi; latere antico manifestè longiore; postico attenuato, subrostrato; extremitate anticâ rotundatâ; posticâ angulatâ; costâ umbonali conspicuâ; ligamento haud prominente; superficie internâ albidâ aut flavidâ; dentibus lateralibus distinctis, æquidistantibus.

Oblong, solid, convex, but slightly glossy, inequilateral, whitish, with (in the typical variety) two deep crimson short rays (thus forming a St. Andrew's cross upon the umbones), but usually only adorned with more or less interrupted rays of tawny-orange or faint lineated markings of purplish crimson, rather obliquely distinctly but not coarsely concentrically grooved; anterior side much the longer, and rounded at its extremity, its dorsal slope slight but convex; posterior side attenuated, the slightly convex ventral edge rising on that side, so as to form a beak in its junction with the moderately sloping and scarcely convex ligamental one; umbonal ridge blunt but conspicuous; inside white or yellowish, the lateral teeth distinct, and equidistant. The typical variety is extremely rare, I have only seen two specimens in the collection of William Walton, Esq., and the original one in the French Museum.

Catbalonga, Isle of Samar; &c.

5. *TELLINA ANTONII* (pl. lviii. f. 74, 75; and pl. lx. f. 131.) *Philippi*, Ab. und Bes. Conch. pt. v. t. 5. f. 3.

T. testâ oblongo-elongatâ, solidâ, convexâ, albidâ aut purpurâ, maculis lineisve nebulosis radiantibus fusco-purpureis rubro-roseis aut fulvo-aurantiis plerumque ornatâ, concentricè sulcatâ et obsolete radiatim striatâ; sulcis posticè sublamellosis; margine ventrali anticè crenato, posticè retuso et sursum acclinato; dorsali anticè vix declivi, posticè subrecto subincurvato satisque declivi; latere postico manifestè breviorè, rostrato; flexurâ costâque umbonali conspicuis; ligamento subdepresso; dentibus lateralibus validis, subæquidistantibus.

Elongated-oblong, solid, convex, somewhat glossy, variable in colouring being uniform whitish or purple (both within and without), but more commonly subradiated by purplish brown or rosy-red markings on a whitish ground; surface with very distinct concentric sulci, which become strong elevated striæ as they approach the prominent umbonal ridge, on and beyond which they become sublamellar; the interstices traversed by nearly obsolete radiating striæ; ventral edge swelling out in front and sloping upwards behind, so as to render the narrow and biangulated posterior extremity rostrated; anterior side decidedly the longer, its extremity rounded, its slope very slight; ligamental edge straightish

but somewhat incurved, its slope moderate : flexure conspicuous ; lateral teeth strong and nearly equidistant.

Guadaloupe.

6. *TELLINA VERRUCOSA* (pl. lxxviii. f. 77.) *Hanley*, Zool. Proc. 1844.

T. testâ oblongo-elongatâ, solidâ, compressâ ; aut flavidâ rubropurpureo radiatâ, aut pallidè roseâ radius subalbidis angustis zonisque saturatoribus ornatâ ; valvulâ alterâ undique (natibus excipiendis) verrucosâ ; alterâ medio concentricè sulcatâ, lateribus solùm verrucosâ ; verrucis ellipticis aut semilunatis, in seriebus vix interruptis, concentricè ordinatis ; margine ventrali anticè subarcuato, posticè sursum acclinato ; dorsali anticè convexiusculo et vix subdeclivi, posticè subrecto, subdeclivi ; latere antico longiore, rotundato ; extremitate posticâ attenuatâ, subrostatâ, obliquè subbiangulatâ ; ligamento posticâ haud prominente ; flexurâ ventrali costâque umbonali conspicuis ; dentibus lateralibus magnis, æquidistantibus.

Elongated-oblong, depressed, solid, yellowish with reddish crimson rays or pale pink with deeper concentric bands and paler narrow rays ; surface roughened in one valve by concentrically arranged and frequently connected compressed wart-like tubercles, which on the disc of the other valve are replaced by concentric elevated sulci ; anterior side the longer, rounded ; ventral edge arcuated anteriorly and curving upwards to the angulated posterior extremity so as to render it subrostrated and attenuated ; slopes gentle, ligament not prominent ; lateral teeth large.

Corrigidor, Bay of Manilla.

7. *TELLINA RASTELLUM* (pl. lxiv. f. 231. and pl. lxxv. f. 242) *Hanley*, Zool. Proc. (April) 1844 ; *Hanl.* Des. Cat. sup. t. xiv. f. 14. *T. Philippii*, *Anton* Ab. und Bes Conch. part v. t. 5.

T. testâ elongatâ, solidâ, convexiusculâ, æquilaterali, flavidâ, pallidè roseo radiatâ ; anticè sulcatâ, posticè squamosâ, squamis erectis, lamellosis, in iisdem seriebus cum sulcis concentricis, ordinatis ; margine ventrali vix convexiusculo ; dorsali utrinque paululum declivi, anticè vix convexiusculo, posticè recto ant subincurvato ; latere postico attenuato, subrostrato, ad extremitatem obliquè biangulato ; extremitate anticâ rotundatâ ; ligamento costâque umbonali, conspicuis ; superficie internâ albidâ, umbones versus aurantio utrinque fucatâ ; dentibus lateralibus validis, subæquidistantibus.

Elongated, solid, moderately convex, equilateral, yellowish with pale crimson rays, the concentric sulci posteriorly so interrupted as to form rows of thin tooth-like scales, and so arranged that the

teeth of one line are just beneath the interstices in the other; dorsal slopes gentle, the posterior one nearly straight, anterior extremity somewhat narrowed and rounded, posterior end much attenuated and subrostrated; ventral edge but little convex; umbonal ridge, ligament, and flexuosity, very prominent; inside white, disc tinged with orange, a short orange ray on either side near the beak; lateral teeth strong, and subequidistant.

Zanzibar.

A species sometimes confused with *Pulcherrima*, but much more elongated, the striæ stronger, and the scales entirely absent from the posterior side. It is not impossibly the *Aspera* of Gmelin, but that shell is too briefly described for recognition.

8. *TELLINA PULCHERRIMA* (pl. lxi. f. 165.) *Sowerby*, Tank. Cat. App. p. 3. t. 1. f. 1; Hanl. Des. Cat. p. 67. sup. t. 11. f. 14; Knorr. vol. iv. t. 25. f. 1.

T. testâ oblongâ, solidâ, convexiusculâ, subæquilaterali, flavo rubroque radiatâ, in medio concentricè et tenuiter sulcatâ; utrinque squamosâ, squamis erectis, dentiformibus aut sublunatis; margine ventrali convexiusculo; dorsali utrinque leviter declivi; extremitate anticâ rotundatâ; posticâ subattenuatâ, angulatâ; superficie internâ albidâ, umbones versus aurantio tinctâ; dentibus lateralibus validis, subæquidistantibus.

Oblong, solid, moderately convex, subequilateral, rayed with pale crimson red and yellow, concentrically sulcated; sulci fine and terminated at either extremity and near the little arcuated ventral margin, by sublunate scales, which in some specimens become more elevated and somewhat tongue-shaped; dorsal slopes gentle; anterior extremity rounded; posterior rather attenuated; lateral teeth very strong, and nearly equidistant.

Var. *a.* More elongated, sulci and scales stronger and more numerous. Var. *b.* Sulci very fine and rather oblique; scales visible along the ventral margin.

Singapore, and Straits of Banca (Cuming).

9. *TELLINA ASPERRIMA* (pl. lx. f. 135.) *Hanley*, Zool. Proc., 1844.

T. testâ oblongo-ellipticâ, convexiusculâ, solidiusculâ, flavidâ aut incarnatâ, roseo-radiatâ; totâ superficie externâ squamis asperimâ; squamis anticè sublunatis, posticè (et præsertim supra costam umbonalem distinctam) spinosis; margine ventrali medio subrecto, flexurâ distinctâ; dorsali utrinque subdeclivi, et paululum convexi-

usculo; latere antico brevior; extremitate posticâ subbiangulatâ, attenuatâ; dentibus lateralibus magnis, antico approximato.

Oblong-elliptic, moderately convex, rather solid, flesh-coloured, with dull crimson rays, the entire surface rough with tooth-like scales, which become more crowded at the rounded anterior extremity, and between the distinctly indicated umbonal ridge and the nearly straight posterior edge rise into rather distant elevated fangs; slopes gentle; posterior side the longer, its extremity angulated; ventral edge nearly straight; teeth very distinct, the lateral large, the anterior lateral approximate. The scales are replaced near the beaks by raised concentric striæ.

An unique specimen, in the cabinet of Mr. Cuming, was found by him at Sual, province of Sangasinan, Isle of Luzon (sandy mud, six fathoms). Allied to *Pulcherrima*.

10. *TELLINA SULCATA* (pl. 60. f. 138.) *Wood*, Gen. Conch. p. 178. t. 47. f. 1; *Hanl. Des. Cat.*, t. 4. f. 63; *Dilw.* p. 89.

T. testâ ovali-subtrigonâ, solidâ, subcompressâ, subæquilaterali, subimpolitâ, albidâ aut flavescente, concentricè sulcatâ et radiatim striatâ; sulcis elevatis, subconfertis, ad costam umbonalem prominentem rudioribus; flexurâ conspicuâ; margine ventrali convexo; dorsali utrinque (et præsertim posticè) declivi, anticè subrecto, posticè recto aut subincurvato; latere postico leviter subrostrato, attenuato; extremitate posticâ biangulatâ; superficie internâ flavidâ; dentibus lateralibus validis, subæquidistantibus.

Oval-subtriangular, solid, rather depressed, subequilateral, rather dull, white or pale straw coloured, with regular rather close raised concentric sulci, which become sublamellar on the prominent umbonal ridge; the interstices with radiating striæ; ventral edge convex; flexure conspicuous; anterior dorsal slope gentle, and nearly straight; posterior side attenuated, its dorsal slope straight and rather strong, its extremity biangulated; inside yellowish; lateral teeth strong and nearly equidistant.

A small specimen was procured by Mr. Cuming, at Catbalonga, Isle of Samar.

~ 11. *TELLINA INTERRUPTA* (pl. lxiii. f. 207.) *Wood*, Gen. Conch. (1815), p. 146. t. 36. f. 3; *Dilw.* 75. T. maculosa, *Lam.* (1818) no. 4; *Chem.* vol. vi. f. 73; *Enc. Mèth.* t. 288. f. 7.

T. testâ ovali aut ovali-oblongâ, æquilaterali, solidâ, subventricosâ, impolitâ, albâ, lineis undosis angularibus lividis brunneis fulvis aut purpureis nebulosâ aut subradiatâ; concentricè sulcatâ, sulcis posticè membranaceis; margine ventrali subarcuato; dorsali utrinque declivi et convexiusculo; latere postico attenuato, subros-

trato; extremitate posticâ angulatâ; costâ umbonali prominente; natibus plerumque purpureis; umbonibus flavis; ligamento magno; superficie internâ flavidâ; dentibus lateralibus magnis, subæquidistantibus.

Var. Testâ angustiore; sulcis remotioribus; margine ventrali rectiore, dorsali utrinque minus declivi.

Oval or oval-oblong, equilateral, solid, subventricose, not polished, white crowded with narrow zigzag linear streaks of fawn-colour brown chocolate or purple, beaks usually tipped with purple, umbones yellowish; surface with moderately distant raised concentric sulci, which become more elevated but *not* thickened on the conspicuous umbonal ridge; ventral edge moderately arcuated; anterior side rounded at its extremity, the dorsal slope gentle; posterior side attenuated, subrostrated, the slope strong and very slightly convex, the tip obtusely angulated; inside tinged with yellow; lateral teeth very strong, subequidistant.

Var. Narrower and more elongated; dorsal edges less sloping; ventral edge straighter; sulci rather more distant.

Surinam. The *T. Mexicana* of the 'Revue Cuverienne,' is now regarded by its author as an elongated variety of this species.

- 12. *TELLINA VIRGATA* (pl. lxiii. f. 212.) *Lin. Sys.*, p. 1116; *Chem.* vol. vi. f. 67; *Enc. Mèth.* t. 288. f. 2; *Mawe's Conch.* t. 6. f. 1; *Lam.* no. 5; *Hanl. Des. Cat.* p. 61. t. 3. f. 3; *Wood, Gen. Conc.* t. 35. f. 2, 3.

*T. testâ ovali, subtrigonâ, solidâ, convexiusculâ, subæquilaterali, flavâ, roseo-purpurascente radiatâ, concentricè sulcatâ,; sulcis anticè subimbricatis, posticè rudioribus magisque elevatis; margine ventrali posticè sursum acclinato, medio subrecto, anticè arcuato, carinâ umbonali prominente; latere antico brevior, rotundato; postico attenuato, subrostrato; margine dorsali postico elongato, declivi, subincurvato; ligamento magno; superficie internâ aurantio tinctâ; dentibus lateralibus validis, subæquidistantibus.*

Oval, subtriangular, solid, moderately convex, nearly equilateral, rich yellow, either uniform, or closely rayed with dull purplish crimson; the concentric sulci becoming closer and slightly imbricated anteriorly, and coarser more prominent and distant as they approach the projecting umbonal ridge; ventral edge subarcuated in front, nearly straight in the middle, and sloping up towards the attenuated and subrostrated posterior extremity; ligamental edge somewhat incurved, much sloping; ligament large and prominent; within usually tinged with orange, the lateral teeth strong and subequidistant.

Indian Ocean.



— 13. *TELLINA JUBAR* (pl. lxiii. f. 214.) *Hanley*, Zool. Proc. 1844. *T. virgata*, var. *Wood*, Gen. Conc. t. 36. f. 2. Enc. Mèth. t. 288. f. 4. (bad).

*T. testâ* specie priore magis triangulari, altiore, minus elongatâ; rubro-purpureâ, radiis albis aut albidis ornatâ; margine ventrali subarcuato; dorsali utrinque decliviore; superficie internâ albidâ, aut coloribus externis fucatâ.

Extremely like *Virgata*, but more triangular and less elongated, the dorsal slopes being more sudden; the ventral edge is more convex, and less straight in the middle; the external colouring of broader and narrower white rays on a dull rosy or reddish ground, is frequently visible on the otherwise white surface of the interior.

Mus. Cuming.

The ligamental margin inclines to convexity, which is the reverse of what we meet with in *Virgata* and *Marginalis*.

14. *TELLINA MARGINALIS* (pl. lxiii. f. 216.) Dilw. Des. Cat. p. 74; Chem. f. 68, (tolerable).

*T. testâ* *Virgatæ* et *Jubaris* maximè assimili, altiore autem, magisque rotundatâ; margine ventrali planè arcuato, posticè sursum acclivi; dorsali utrinque leviter declivi; superficie externâ albidâ, radiis rubescentibus ornatâ.

Closely allied to the two preceding, but less oblong than either; the ventral edge much more arcuated, and ascending abruptly on the posterior side; the ligamental edge sloping, but slightly and somewhat concavely; the colouring whitish, with pale rosy-red or deep flesh-coloured rays; interior nearly white.

Indian Ocean? &c.

15. *TELLINA STAURELLA* (pl. lx. f. 148. and pl. lxi. f. 171. and pl. lxv. f. 261.) *Lam.*, no. 6; *Hanl. Des. Cat.* p. 61. sup. t. 11. f. 33; *Delessert*, t. 6. f. 2; *Chem.* vol. vi. f. 70, 71; *Philip. Ab.* und *Bes. Conch.* part v. t. 5. f. 6. *T. scalaris*, *Lam.* no. 29.

*T. testâ* ovali aut oblongo-ovali, solidâ, convexâ, nitidâ, extus intusque albidâ aut flavidâ, radiis angustis roseis confertis plerumque ornatâ; concentricè sulcatâ, sulcis anticè subobliquis et subimbricatis, posticè submembranaceis et paucioribus (plerumque in valvulâ alterâ omnino obsoletis); margine ventrali convexo; dorsali anticè subdeclivi et vix convexiusculo, posticè declivi et subrecto; latere antico manifestè longiore, postico attenuato; extremitate posticâ obtusè angulatâ; natibus plerumque roseis, nonnunquam cruce roseo-rubrà notatis; ligamento magno, subconspicuo; dentibus lateralibus magnis, subæquidistantibus.

Oval or oblong-oval, solid, convex, glossy, white or yellowish, with narrow pink or rosy rays (which often become obsolete,) and concentric sulci, which, subimbricated in front, are almost lamellar towards the decidedly shorter posterior side, where in one valve they almost always become obsolete; dorsal slopes rather gentle; ventral edge convex; posterior extremity attenuated and angulated; umbonal ridge distinct; inside whitish or tinged with yellow, beaks tinged with rose-colour; lateral teeth strong and nearly equidistant.

Var. *a*, with two short rays at the beak in each valve, forming a small crimson cross; a smooth posterior area in one valve. Var. *b*, beaks with a very large crimson cross; no rays; sulci finer; area as in last. Var. *c*, rays distinct; no smooth area. Var. *d*, As last, but perfectly white, and oval-oblong.

*T. scalaris* is but an abbreviated variety of this very variable species, with the posterior side more strongly biangulated and less attenuated at the extremity. The typical specimen of the French Museum is yellow, (externally very pale) with the apical cross of a rich red and very small.

Philippines &c., very common.

16. *TELLINA GUILDINGII* (pl. lvi. f. 1.) *Hanley*, Zool. Proc. 1844.

*T. testâ oblongâ, tenui, compressâ, subæquilaterali, albidâ, radiis aurantio-roseis zonisque albido-roseis ornatâ; sulcis exilibus confertis concentricè exaratâ; margine ventrali subrecto, medio subretuso; dorsali utrinque declivi, et vix convexiusculo; latere postico paululum brevior, attenuato; extremitate posticâ infernè angulatâ, anticâ rotundatâ; natibus acutis; costâ umbonali et flexurâ ventrali inconspicuis; ligamento prominente; dentibus lateralibus magnis, æquidistantibus.*

Oblong, thin, compressed, subequilateral, whitish, with narrow roscate flesh-coloured rays, and fine close-set concentric sulci; ventral edge nearly straight, rather retuse in the middle; dorsal margin scarcely at all convex or sloping on either side; posterior side very slightly the shorter, attenuated, its extremity angulated below (biangulated in the young, and the ligamental edge retuse); anterior end rounded; umbonal ridge obsolete; flexure inconspicuous; ligament prominent; inside pale; lateral teeth large, and equidistant.

West Indies.

17. *TELLINA PULCHELLA* (pl. lvi. f. 4.) *Lam.*, no. 23; *Hanl.* Des. Cat. p. 64. sup. t. 1. f. 4; *Philip. En. Mol. Sicil.* vol. i. p. 24.

*T. rostrata*, *Born.* t. 2. f. 10; *Poli. Tes. Sic.* vol. xv. f. 8; *T. virgata*, var. *Chem.* vol. vi. p. 72.

*T. testâ oblongâ, angustâ, tenui, compressâ, nitidâ, radiis roseis vel roseo-aurantiis et albidis vel flavidis eleganter ornatâ, striis confertis exilibus concentricè sulcatâ; margine ventrali subrecto; latere antico producto, rotundato; postico brevior, valde attenuato, subrostrato; margine dorsali anticè paululum declivi, posticè valde declivi, convexo, intusque roseo; natibus albidis; superficie internâ albido-roseâ; dentibus lateralibus validis, antico subapproximato.*

Var. *testâ omnino roseâ.*

Elongated-oblong, very thin, compressed, glossy, rayed with rose-colour and white, the former often passing into orange, the latter tinged usually with yellow; more rarely of an uniform rose-colour, except the tips of the beaks which are always pale or whitish; most delicately but regularly and closely striated concentrically; ventral edge nearly straight; anterior side produced, its slope very gentle, its extremity rounded; posterior side shorter, much attenuated, subrostrated, the dorsal slope great but convex; inside pink, ligamental edge rosy; lateral teeth strong, anterior one subapproximate.

Mediterranean.

18. *TELLINA VIRGULATA* (pl. lvi. f. 5.) *Hanley, Zool. Proc.* 1844.

*T. testâ Donacinae* simillimâ, sed paululum angustiore, striisque exilioribus ornatâ; extus intusque albidâ, roseo-radiatâ; radiis latis, haud interruptis; margine dorsali anticè albido.

Extremely like *Donacina*, but the shape is slightly narrower, the striae more delicate, and the colouring consists of a few interrupted rose-coloured rays upon a white ground; the interior dorsal edge is white.

Mus. Cuming.

19. *TELLINA DISTORTA* (pl. lvi. f. 6.) *Poli, Test. Sic.* vol. i. t. 15. f. 11; *Philip. Mol. Sic.* vol. i. p. 25.

*T. testâ ovato-oblongâ, compressiusculâ, tenui, nitidâ, subpellucidâ, pallidè roseâ aut aurantio-roseâ, radiis saturatoribus plerumque pictâ; concentricè striatâ, striis minimis et posticè elevatis; margine ventrali convexiusculo; dorsali utrinque subrecto, anticè minimè declivi; latere antico longiore, ad extremitatem obtusè rotundato; extremitate posticâ cuneiformi, paululum subrostratâ; flexurâ costâque umbonali subinconspicuis; ligamento parvo, satis prominente; dentibus lateralibus distinctis, postico subremoto.*

Ovate-oblong, slightly compressed, thin, subpellucid, glossy, pink or rosy orange, with narrow deeper-coloured rays, crowded with minute concentric striæ which become more distinct and elevated posteriorly; ventral margin slightly convex; anterior side the longer, obtusely rounded at its extremity, its dorsal edge scarcely at all declining; posterior extremity wedge-shaped, the ligamental edge nearly straight; ligament small but not sunken; fold and flexure not conspicuous; lateral teeth distinct, posterior the more distant.

Mediterranean.

20. *TELLINA DONACINA* (pl. lvi. f. 12. and pl. lxvi. f. 259.) *Lin.*, Sys. 1118; *Lin. Trans.* vol. viii. t. 1. f. 1.; *Mont. Test. Brit.* t. 27. f. 3; *Turt. Biv. Brit.* p. 103. t. 8. f. 4; *Wood, Gen. Conc.* t. 45. f. 5; *Philip, En. Mol. Sic.* vol. i. p. 24; *Lam. no. 27*; *Hanl. Des. Cat.* p. 64. t. 4. f. 31. *T. variegata Poli*, vol. i. t. 15. f. 10. *T. Llantivyi, Payr. Cat. Cors.* p. 40. t. 1. f. 13, 14.

*T. testâ oblongâ, solidiusculâ, subconvexâ, impolitâ, albido-flavescente, radiis roseo-rubris angustis interruptè ornatâ, striisque subimbricatis concentricè sulcatâ; margine ventrali convexiusculo, et anticè arcuato; margine dorsali roseo, anticè minimè posticè subito, declivi; latere antico latitudinem postici duplo superante; extremitate anticâ rotundatâ, posticâ obtusissimè cuneiformi; natibus radio brevi recto roseo strigatis; superficie internâ flavescente, propenates roseo-tinctâ; dentibus lateralibus validis, subæquidistantibus.*

Var. *Penitus candidâ.*

Oblong, rather solid, moderately convex, rather dull, pale yellow, with crimson interrupted narrow rays, and concentric subimbricated striæ; ventral edge moderately convex, swelling out anteriorly; dorsal edge rosy; beaks with a short direct rosy streak; anterior side thrice as long as the posterior, its extremity rounded, its dorsal edge nearly horizontal; posterior side very short, most obtusely cuneiform, inside tinged with yellow, becoming pink near the beaks, teeth strong, subequidistant.

Var. *Perfectly white.*

*Britain, Europe.*

21. *TELLINA TULIPA* (pl. lvi. f. 25.) *Hanley, Zool. Proc.* 1844.

*T. testâ ovali-oblongâ, solidâ, subinæquilaterali, compressâ, nitidiusculâ, lineis elevatis concentricè et confertim striatâ; albidâ aut albido-roseâ, radiis roseis multis angustis subinterruptè ornatâ; margine ventrali medio retuso; dorsali utrinque subrecto, anticè vix paululum declivi, posticè satis declivi; latere antico paulo lon-*

giore, ad extremitatem obtusè rotundato; postico truncato-cuneiformi; ligamento prominente; costâ umbonali subobsoletâ; dentibus ut in *Donacina*.

Oval-oblong, solid, slightly inequilateral, compressed, somewhat glossy, with very close concentric striæ, which are subimbricated in front, and raised behind; white or pale pink, with numerous rosy somewhat interrupted rays; ventral edge subretuse in the middle; dorsal edges nearly straight, the front one scarcely, the hinder one moderately sloping; ligament prominent; anterior side rather the longer, and obtusely rounded at its termination; posterior side of a truncated wedge-shape; umbonal ridge not prominent; inside paler than the exterior but usually tinged with rose-colour; teeth as in *Donacina*.

Mus. Cuming, Walton.

22. *TELLINA LINEATA* (pl. lvi. f. 35, 36. and pl. lvii. f. 46, 47.)  
*Turton*, Conch. Dict. p. 168. f. 16; *Turt. Biv. Brit.* p. 99. t. 7. f. 1.  
*T. Brasiliana Lam.* no. 46; *Hanl. Des. Cat.* p. 70. sup. t. 9. f. 3.  
*T. striata Mont.* Test. Brit. p. 60. t. 27. f. 2.

T. testâ ovatâ, solidâ, convexâ, concentricè et confertissimè striatâ; margine ventrali medio subrecto; dorsali anticè magis minusve convexo et paululum declivi, posticè umbones versus subretuso, deinde convexo et declivi; latere antico longiore, ad extremitatem rotundato; extremitate posticâ cuneiformi; ligamento prominente; flexurâ costâque umbonali satis conspicuis; dentibus lateralibus validis, subæquidistantibus.

Var. *a*, Testâ sordidè albidâ, umbonibus nonnunquam aurantiis, radioque brevi obliquo roseo anticè ornatis. Var. *b*, Penitus candidâ. Var. *c*, Ovato-oblongâ, lævigatâ, nitidâ, albido-roseâ, radio obliquo ut in var. *a*; intus roseâ.

More or less ovate solid and convex, with concentric minute crowded striæ; the ventral margin nearly straight in the middle; anterior side the longer (sometimes scarcely so,) its extremity rounded, its dorsal edge but slightly sloping; posterior extremity wedge-shaped, the ligamental edge, which near the beaks is somewhat retuse, becoming convex at its termination; fold and flexure inconspicuous; lateral teeth strong, and subequidistant.

Var. *a*, Pale, with a short oblique anterior crimson ray at the umbones, which are sometimes tinged with orange. Var. *b*, Entirely white. Var. *c*, The ray as in var. *a*, but the shape more elongated, and the surface pink smooth and glossy; inside rosy.

Britain (?), West Indies, Florida, and Brazil.

There being a previous *Brasiliana*, the subsequent name of Turton must be adopted.

23. *TELLINA GELIDA*, *Hanley*.

T. testâ ellipticâ, æquilaterali, solidiusculâ, subinæquivalvi, convexiusculâ, subnitidâ, intus extusque candidâ, lineis elevatis concentricis subconfertim striatâ, striis anticè et posticè elevatioribus; margine ventrali convexo; dorsali utrinque subdeclivi, subrecto, anticè lamellis lanceolatis dentato; natibus haud incumbentibus; extremitate anticâ rotundatâ; posticâ subbiangulatâ, costâ umbonali distinctâ; ligamento infosso; dentibus lateralibus subæquidistantibus.

Elliptic, equilateral, slightly inequivalve, tolerably strong, moderately convex, scarcely glossy, white both within and without, with elevated and rather close concentric lines, which are stronger at the extremities, interstices quite smooth; ventral edge convex; dorsal but moderately sloping on either side, nearly straight, dentated in front by lanceolate lamellæ; beaks not inclining; front extremity rounded, hinder one somewhat biangulated; fold distinct; ligament sunken; lateral teeth nearly equidistant.

Mus. Belcher.

Unfortunately a figure of this distinct species cannot be given, as the possessor of the unique specimen is not in England.

24. *TELLINA SERRATA* (pl. lxxv. f. 256.) *Costa*, p. xvi. no. 8; Philip. Er. Mol. Sic. p. 25; Ab. und Bes. Conch. part 5. t. 5 f. 5.

T. testâ ovali-trigonâ, solidiusculâ, satis convexâ, inæquilaterali, nitidiusculâ, intus extusque albidâ, (umbonibus plerumque flavis;) concentricè striatâ, striis exilibus, confertissimis, anticè posticè et infernè subelevatis; margine ventrali convexiusculo; dorsali anticè convexiusculo vixque declivi, postico subrecto et valdè declivi; latere antico longiore, ad extremitatem rotundato; posticè cunei-formi; costâ umbonali satis conspicuâ; natibus acutis; ligamento infosso; dentibus lateralibus, magnis, haud remotis, subæquidistantibus.

Subtriangularly oval, tolerably solid and convex, inequilateral, slightly glossy, within and without whitish, with the umbones usually bright yellow, most closely and delicately striated; the striæ raised at the extremities and towards the moderately convex ventral margin; front dorsal edge moderately convex and not much sloping, hinder one straightish and considerably sloping; front side the longer, its extremity rounded; hinder side wedge-shaped; fold

distinct; beaks acute; ligament sunken; lateral teeth large, tolerably and almost equally close.

Mediterranean.

25. *TELLINA OWENII* (pl. lx. f. 133.) *Hanley, Zool. Proc.* 1844.

T. testâ ovato-oblongâ, solidiusculâ, subimpolitâ, compressâ, æquilaterali, albidâ, concentricè et confertissimè striatâ; margine ventrali valdè arcuato; dorsali utrinque subdeclivi, anticè subrecto, posticè incurvato et lamellis subdentato; extremitate anticâ rotundatâ; latere postico acuminato, subrostrato; costâ umbonali conspicuâ; flexurâ satis distinctâ; natibus acutis; ligamento infosso; disco interno aurantio; dentibus lateralibus, subremotis, subæquidistantibus.

Ovate-oblong, tolerably strong, rather dull, whitish, (with often a tinge of orange in the front part of the umbones, and the internal disc always of that colour,) equilateral, compressed, with crowded concentric striæ; ventral edge much arcuated; dorsal moderately sloping on either side of the acute beaks, nearly straight in front, incurved behind and somewhat toothed by the lamellæ, (which in some specimens commence before the conspicuous fold); ligament sunken; lateral teeth distinct, and nearly equally remote.

Africa.

26. *TELLINA PHARAONIS* (pl. lxiii. f. 215.) *Hanley, Zool. Proc.* 1844.

T. testâ *T. Vulsellæ* simillimâ, sed solidâ lineisque elevatis concentricè striatâ, umbonibus aurantio-roseis; flexurâ sinuatâ.

Extremely like *T. Vulsellæ*, but solid, and rough with concentric elevated striæ; fold sinuated.

In the only specimen I have ever seen, one of the valves is of a delicate pink, the other white tinged with the same hue, the beaks of a very rosy orange and the ventral flexure so deep as to form a sinus; the lower margin is not so convex in front as in *Rostrata*; the front lateral tooth is distinctly crenated, but this is probably accidental.

Red Sea.

27. *TELLINA VULSELLA* (pl. lxi. f. 162, 163.) *Chem.* T. rostrata, *Gmel.* p. 3233 (in part); *Dil. Des. Cat.* p. 84; *Lam.* no. 9; *Chem.* vol. vi. f. 105; *Enc. Mèth.* t. 289. f. 1; *Wood, Gen. Conc.* t. 37. f. 3; *Knorr.* vol. iv. t. 2. f. 2, 5; *Hanl. Des. Cat.* p. 62. t. 4. f. 28.

T. testâ valdè clongatâ, compressâ, tenui, subpellucidâ, nitidis-

simâ, sublævigatâ, roseâ, radio pallidiore postico plerumque notatâ; margine ventrali anticè arcuato, posticè flexuoso et sursum acclinato; dorsali utrinque vix minimè declivi, prope nates acutas subretuso, deinde subrecto aut paululum convexiusculo; latere antico longiore, ad extremitatem rotundato; postico angusto, acuminato, rostrato; ligamento infosso; flexurâ costâque umbonali conspicuis; dentibus lateralibus parvis, tenuibus, æquidistantibus.

Var. Flavo-aurantiâ.

Much elongated, compressed, thin, subpellucid, polished, rose-coloured, with usually a paler posterior ray, nearly smooth, or with minute radiating and concentric striæ; ventral edge arcuated in front, flexuous and greatly elevated behind; dorsal edge scarcely at all sloping in front, rather more so posteriorly, subretuse near the acute beaks, and then straightish or very slightly convex; anterior side the longer, rounded at its extremity; posterior side forming a narrow acuminated beak; ligament sunken; fold and flexure conspicuous; lateral teeth thin, small, and equidistant.

Var. Rich yellow.

Yellow var. Isle of Zebu.

As this is not the *Rostrata* of Linnæus, I have adopted the other name proposed by Chemnitz "*Rostrata seu Vulsella*".

28. *TELLINA PERNA* (pl. lxiii. f. 202, 217, 219.) *Spengler*, *Schriфт. Naturhis. Gelstav. Kiobenha. 4.* (1798). *T. pallescens Dil. Des. Cat. p. 84.* *T. sulphurea* and *latirostra Lam. no. 10, 11. Hanl. Des. Cat. p. 62.* *T. incarnata Born p. 36. t. 2. f. 12.* (erroneously quoted as 11 in text). *T. rostrata flavescens Chem. vol. vi. f. 104.* *T. latirostra, Swain. Zool. Ill. 2nd ser. vol. i. t. 20.*

*T. testâ oblongo-elongatâ, tenuiusculâ, convexâ, subpellucidâ, nitidâ, sublævigatâ, albâ, flavâ, aurantiâ aut roseâ, radio pallidiore postico plerumque ornatâ; margine ventrali anticè arcuato, posticè sursum declivi; dorsali anticè vix declivi, convexiusculo, posticè paululum declivi, prope nates conspicuas subincurvato, mox convexiusculo; latere antico longiore, subproducto; postico rostrato; extremitate posticâ obtusè angulatâ; umbonibus subtumidis; ligamento infosso; margine cardinalis interno, angusto, callo ligamentali munito; dentibus lateralibus tenuibus, inconspicuis, æquidistantibus.*

Elongated-oblong, rather thin, semitransparent, convex, glossy, very variable in colouring, being sulphur-yellow, white, pink with or without paler rays, orange with white rays, or orange flesh-coloured, &c., nearly smooth, often with obsolete radiating striulæ, which are more evident posteriorly; ventral edge arcuated in front,



curving up posteriorly; anterior side rather the longer, its extremity obtusely rounded, scarcely any dorsal slope, but the edge convex; posterior side beaked, its dorsal edge moderately sloping and convex at its termination, its extremity obtusely subangulated; umbones rather tumid; ligament sunken; hinge-margin narrow, a callus near the teeth; lateral teeth thin, equidistant, indistinct.

Singapore, Isle of Burias, &c.

29. *TELLINA ALBINELLA* (pl. lxi. f. 164, 166.) *Lam.*, no. 17; *Hanl. Des. Cat.* p. 63. sup. t. 14. f. 3.

T. testâ oblongo-ovali, subæquilaterali, tenuiusculâ, complanatâ, nitidiusculâ, extus intusque albidâ; alterâ valvulâ sublævigatâ, alterius arêâ posticâ concentricè striatâ; margine ventrali vix convexiusculo; dorsali anticè arcuato et declivi, posticè declivi et subincurvato; ligamento infosso; flexurâ subinconspicuâ; costâ umbonali carinatâ; dentibus lateralibus perparvis, inconspicuis, antico approximato. Var. Roseâ.

Oval-oblong, rather thin, flat, subequilateral, slightly glossy, whitish, (usually with a fulvous stain towards the beaks,) nearly smooth, except on the posterior half of one of the valves which is deeply and distinctly striated concentrically; ventral edge nearly straight in the middle and rising behind; front dorsal edge arched and deeply sloping; posterior side acuminate, angulated below, and slightly beaked, the ligamental edge moderately sloping, and rather incurved; ligament sunken; inside whitish, the lateral teeth very small, indistinct, the anterior one closely approximate.

Var. Rose-coloured.

Australia.

30. *TELLINA ROSEA* (pl. lxi. f. 170) *Spengler*, *Schriiv. Natur. Gelsta. Kioben.* vol. 4. 1798; *T. pallidè rosea*, *Chem.* vol. 6. f. 96.

T. testâ oblongâ, crassissimâ, subæquilaterali, convexiusculâ, nitidâ, roseâ, sublævigatâ, striis concentricis tenuissimis, lineisque radiantibus obsoletis tantum signatâ; striis posticis prope nates acutas remotis, subelevatis; margine ventrali convexo, posticè acclivi; dorsali utrinque declivi, anticè convexo, posticè subrecto plerumque subincurvato; latere postico subcuneiformi; extremitate posticâ subacuminatâ; costâ umbonali angustâ; flexurâ inconspicuâ; ligamento infosso; superficie internâ roseâ, politâ, impressionibus muscularibus fortiter impressis; margine cardinali intus lato, valido; dentibus lateralibus parvis, subæquidistantibus, ætate obsoletis.

Oblong, very solid, subequilateral, rather convex, glossy, pink,

with delicate concentric and obsolete radiating striae, the posterior concentric more distant raised and distinct near the beaks; ventral edge scarcely arcuated, ascending posteriorly; anterior side rounded, its dorsal slope strong; posterior side wedge-shaped, ligamental edge nearly straight, often slightly incurved, moderately sloping, point acuminate; within rosy, polished, scars deeply impressed; hinge margin broad and very strong; lateral teeth subequidistant, rather small, by age obsolete.

Indian Ocean (?).

31. *TELLINA INÆQUISTRIATA* (pl. lvii. f. 58. and pl. lviii. f. 80.) *Donovan*, Brit. Shells, (1802), vol. 4. t. 123; *T. sanguinea*, *Wood*, Gen. Conch. p. 159. t. 44. f. 2; *Hanl. Desc. Cat.* p. 67. t. 4. f. 27.

*T.* testâ oblongo-subtrigonâ, solidiusculâ, subæquilaterali, nitidâ, compressiusculâ, pallidè rubro-sanguineâ, concentricè confertim et exilissimè striatâ; striis posticis alterâ in valvulâ remotis, rudioribus; margine ventrali subrecto; dorsali anticè convexiusculo et subdeclivi, posticè declivi et prope nates retuso; latere antico paulo longiore, ad extremitatem rotundato; postico truncato-cuneiformi; colore interno saturatiore; ligamento prominulo; flexurâ costâque umbonali obsoletis; dentibus lateralibus magnis, antico approximato, postico haud remoto.

Var. *a*, striis posticis in utrâque valvulâ remotis, rudioribus.

Var. *b*, testâ albidâ, pallidè aurantio ad nates utrinque radiatâ.

Oblong, very slightly triangular, rather solid, somewhat compressed; glossy, subequilateral, of a pale blood red, with fine and crowded concentric striae, which however in one of the valves are posteriorly replaced by coarser and more distant ones; ventral edge nearly straight; anterior side rather the longer, its end rounded, its dorsal slope moderate and tolerably convex; posterior end truncated-cuneiform, its dorsal slope strong, nearly straight, or somewhat concave near its acute beaks; ligament but slightly projecting; fold and flexure obsolete; internal colours deeper; lateral teeth large, the posterior not very distant, the anterior subapproximate.

Var. *a*, the posterior striae distant in both valves. Var. *b*, white, with a short pale orange ray on either side of the beaks.

In sculpture resembling *Nitida*.

Bay of Guayaquil; (Cuming).

32. *TELLINA PRINCEPS* (pl. lxiii. f. 206.) *Hanley*, Zool. Proc. 1844.

*T.* testâ oblongo-ellipticâ, æquilaterali, solidâ, compressâ, nitidiusculâ, intus extusque roseo-purpurascente, concentricè sulcatâ; striis exilibus, profundis, radiantibus, sulcos confertos, utrinque (et

alterâ in valvulâ undique) decussantibus; margine ventrali subrecto; dorsali utrinque leviter et æqualiter subdeclivi; extremitate posticâ obtusissimè angulatâ; flexurâ ventrali costâque umbonali, obsolete; ligamento prominente; dente laterali antico approximato et conspicuo, postico parvo, remoto, inconspicuo.

Oblong elliptic, equilateral, solid, compressed, rather glossy, both internally and externally of an uniform reddish lilac colour, the surface decussated by close-set concentric sulci and fine but very distinct radiating striæ, which are only visible at the extreme sides in one of the valves; the ventral edge nearly straight; the dorsal slopes gentle and all but equal; the posterior end most obtusely angulated; umbonal ridge and flexure obsolete; the anterior lateral tooth extremely close and distinct, the posterior very distant and inconspicuous.

Not readily confused with any in this division, but very closely allied to *Purpurascens*, from which the teeth distinguish it.

Tumbez, Peru; (Cuming).

33. *TELLINA PUNICEA* (pl. lviii. f. 89. and pl. lx. f. 154.)  
*Born*, p. 33. t. 2. f. 2; *Lam.* no. 21; *Enc. Mèth.* t. 291. f. 2;  
*Gmel.* 3239; *Dil. Des. Cat.* p. 90.

T. testâ ovali-trigonâ, solidâ, convexiusculâ, subæquilaterali, nitidâ, extus intusque albido-roseâ, (rarius candidâ) prope nates albas colore saturatius tinctâ; concentricè sulcatâ, sulcis confertis, nonnunquam alterâ in valvulâ posticè obsolete; margine ventrali convexiuseulo; dorsali utrinque declivi, anticè convexiuseulo, posticè subrecto; extremitate anticâ obtusâ, et supernè subemarginatâ; latere postico truncato-cuneiformi: flexurâ costâque umbonali inconspicuis; ligamento prominulo aut prominente; dentibus lateralibus distinctis, antico valdè approximato, postico remoto.

Var.? Magis trigonâ, tenuiore, saturatius pictâ, sulcis alterius valvulæ posticè obsolete, emarginatione subobsolete.

Subtriangularly oval, solid, moderately convex, nearly equilateral, glossy, of a pale rose-colour, increasing in intensity near the white-tipped beaks; surface closely and deeply grooved (the sulci rarely obsolete behind); ventral edge slightly convex; anterior side (if either) the shorter, the extremity obtusely emarginated or indented near the dorsal slope, which is slight and rather convex; posterior side of a truncate wedge shape, its dorsal edge nearly straight and much sloping; fold and flexure obsolete; ligament always more or less prominent; lateral teeth distinct, the front one approximate, the hinder remote.

Var.? more triangular, thinner, deeper coloured; the sulci pos-

teriorly quite obsolete in one of the valves; emargination almost obsolete.

Xipixapi, W. Columbia, in sandy mud, 10 fathoms (Cuming); and Trinidad, W. Indies; Britain (?).

Easily distinguished from *Striata*, by its closer and more delicate sulci, its less prominent beaks, and the absence of all radiation. It is also thicker and more opaque, and remarkable for the slight indentation which terminates the rather flattened space adjacent to the front dorsal edge, a character which is not possessed by any of the allied species.

34. *TELLINA STRIATA* (pl. lxi. f. 161.) *Chemn.*, vol. x. p. 349. f. 1654-5; *T. angulosa*, *Gmel.* 3244; *T. læta*, *Montagu*, *Test.* Br. p. 57; *T. punicea*, in part, *Turt. Dic.*; *Donax Martinicensis. Lam.* (fide Deles. *Rec. t. 6. f. 15.*)

*T. testâ* ovali-subtrigonâ, solidiusculâ, subpellucidâ, convexiusculâ, nitidâ, aut albido-roseâ radio roseo utrinque ornatâ, aut albidâ radio flavescente; concentricè sulcatâ, sulcis remotis, alterâ in valvulâ posticè evanescentibus; margine ventrali paululum convexiusculo; dorsali utrinque declivi, anticè convexiusculo, posticè subrecto; latere antico longiore, postico truncato-cuneiformi; extremitate anticâ simplici, rotundatâ; flexurâ costâque umbonali inconspicuis; natibus acutis, pallidis; ligamento prominente; dentibus ut in *Puniceâ*.

Subangularly oval, moderately solid and convex, either pale pink with a roseate ray down the obsolete umbonal ridge (preceded by a paler streak) and a short roseate ray in front of the acute and prominent beaks, or white with the rays yellowish; surface grooved by subimbricated distant sulci, which disappear on one of the valves at the truncate wedge-shaped hinder extremity; ventral edge but slightly convex; anterior side the longer, its extremity simply rounded and never indented, its dorsal edge but moderately sloping and slightly convex; hinder dorsal edge sloping and nearly straight; flexure nearly obsolete; teeth as in *Punicea*.

W. Indies; Britain (?).

The rays are never absent, and the anterior side is always the longer. Mr. Sowerby informs me, that this is the *Tellinides rosacea* of King (*Zool. Journ.* vol. 5.), which is too briefly defined for recognition.

*TELLINA REGIA* (pl. lx. f. 140.) *Hanley*, *Zool. Proc.* 1844.

*T. testâ* oblongâ, tenui, compressiusculâ, subinæquivalvi, subæquilaterali, nitidissimâ, pellucidâ, intus extusque roseo-purpuras-

cente; concentricè sulcatâ, sulcis remotis, alterâ in valvulâ posticè evanescentibus; margine ventrali subrecto, medio subretuso; dorsali utrinque subæqualiter declivi, posticè subrecto; latere antico paululum brevior, ad extremitatem obtusè rotundato; extremitate posticâ supernè angulatâ, attenuatâ; costâ umbonali et flexurâ ventrali obsoletis; ligamento prominulo; dentibus ut in *Puniceâ*.

Oblong, thin, rather compressed, almost inequivalve, subequilateral; extremely glossy, both externally and internally of a deep subpellucid purplish crimson; the surface marked with distant concentric grooves, which posteriorly become obsolete in one or both of the valves; the ventral edge nearly straight, subretuse in the middle; anterior side slightly the shorter, its extremity obtusely rounded; posterior extremity almost biangulated; dorsal moderately and almost equally sloping on either side of the beaks, nearly straight posteriorly; umbonal ridge and flexure nearly obsolete; the ligament rather prominent; teeth as in *Punicea*.

Most closely allied to *Punicea*, but its deep purplish red, and its more distant sulci suffice to distinguish it. From *Princeps* it differs by the absence of radiating striæ; from *Striata*, by its slightly longer posterior side; from *Rubescens*, by its colouring, and the lesser slope of the dorsal edges.

Real Llejios, Central America.

36. *TELLINA EBURNEA* (pl. lviii, f. 91.) *Hanley*, Zool. Proc. 1844. p. 67.

T. testâ oblongâ, opacâ, solidâ, inæquivalvi, convexâ, nitidâ, albidâ, inæquilaterali; concentricè sulcatâ, sulcis profundis, remotis, (alterâ in valvulâ, nonnunquam posticè evanescentibus); margine ventrali convexiusculo, anticè sursum acclinato; dorsali anticè subrecto, leviterque declivi, posticè recto subitoque declivi; latere postico multum brevior, subcuneiformi; lineis erectis obliquis aream nymphalem et analem rugantibus; ligamento brevî, prominente; flexurâ ventrali costâque umbonali subobsoletis; superficie internâ candidâ; dentibus lateralibus, ut in *Puniceâ*.

Oblong, solid, opaque, rather inequivalve, convex, whitish, glossy, clearly inequilateral, with strong deep concentric sulci, which usually become obsolete in one of the valves, and which diverge and become elevated on passing the flattened space at the upper edge of the more convex valve; ventral edge very slightly convex, curving obliquely upward anteriorly; posterior side much the shorter, subcuneiform; the ligamental edge straight, and abruptly sloping; ligament short and prominent; fold and umbonal ridge almost obsolete; inside pure white, teeth as in *Punicea*.

The peculiar elevated sulci on the dorsal areas, at once distinguish it from all the allied species.

Tumbez, Peru.

37. *TELLINA ALTERNATA* (pl. lxi. f. 159.) *Say*, Jour. Ac. Phil. vol. ii. p. 275; Amer. Conch. t. 65. f. 1; Hanl. Des. Cat. p. 66. sup. t. 14. f. 9.

T. testâ oblongo-elongatâ, solidiusculâ, nitidâ, compressiusculâ, albâ aut albo-flavescente, (intus flavâ); concentricè sulcatâ, sulcis remotis, profundis, ad costam umbonalem obsoletissimam alternatim evanescentibus; margine ventrali subrecto; dorsali anticè declivi convexiusculo, posticè recto et multum declivi; latere postico manifestè longiore, ad extremitatem rotundato; antico cuneiformi; ligamento prominente; flexurâ inconspicuâ; dentibus iisdem ut in *Puniceâ*.

Elongated-oblong, rather solid, and somewhat compressed, of a glossy opaque white or yellowish white, grooved concentrically with rather distant deeply impressed sulci, which alternately disappear in crossing the nearly obsolete umbonal ridge; ventral edge nearly straight; anterior side much the longer, rounded at its extremity, its dorsal edge moderately sloping in a convex line; posterior side cuneiform, its dorsal line being straight and greatly sloping; ligament prominent; fold indistinct; inside white, the teeth as in *Punicea*.

In the only specimen I have ever seen of this shell, the striæ are closer in one valve than in the other. In general appearance the shell is extremely like *Striata*.

South Carolina.

38. *TELLINA RUBESCENS* (pl. lx. f. 153.) *Hanley*, Zool. Proc. 1844.

T. testâ *T. Striata* et *Punicea* simillimâ, ab hâc autem differt, testâ tenuiore anticæ emarginationis experte, ab illâ natibus obtusioribus et extremitate posticâ minus attenuatâ; ab utrâque, superficie nitidissimâ, ligamento infosso, margineque dorsali convexiore et utrinque subæqualiter declivi; sulcis in utrâque valvulâ posticè obsoletis; dentibus ut in *Puniceâ*, sed minimis, inconspicuis.

Most closely allied to *Striata* and *Punicea*, but differing from the former by the greater obtuseness of its beaks, and the lesser attenuation of its hinder extremity; from the latter by its more delicate texture and the absence of any anterior emargination; from both, in the sunken ligament, the greater convexity and more equal slope of the dorsal edge. The sulci are obsolete posteriorly

in both the valves, and the whole surface is peculiarly glossy; the teeth are as in *Punicea*, but very small and indistinct.

Tumbez and Panama (Cuming).

39. *TELLINA PRORA* (pl. lx. f. 152.) *Hanley*, Zool. Proc. 1844.

T. testâ subtrigonâ, ovali, solidâ, convexiusculâ, subæquilaterali, nitidâ, extus intusque pallidè roseâ; aut sublævigatâ, aut concentricè et tenuissimè striatâ; margine ventrali convexo; dorsali utrinque valdè et subæqualiter declivi, subrecto; extremitate posticâ acuminatâ; flexurâ ventrali costâque umbonali subobsoletis; ligamento prominulo; dentibus lateralibus magnis, antico approximato.

Subtriangular, oval, tolerably convex, solid, glossy, nearly equilateral, uniform pink both within and without, becoming deeper towards the beaks, most delicately and regularly striated concentrically; ventral edge convex; dorsal edges nearly equally sloping, slopes strong and scarcely at all convex; posterior end acuminate; fold and flexure nearly obsolete; ligament but little prominent, both slopes being slightly excavated; lateral teeth large, the anterior approximate.

Porto St. Elena, and Salango, West Columbia.

40. *TELLINA LACERIDENS* (pl. lxi. f. 168 and 176. and pl. lxvi. f. 258.) *Hanley*, Zool. Proc. 1844.

T. testâ oblongo-subtrigonâ, solidâ, opacâ, subæquilaterali, nitidissimâ, convexiusculâ, albâ (intus candidâ), supernè lævigatâ, infernè concentricè et confertim striatâ; margine ventrali convexiusculo; dorsali utrinque convexiusculo, et subæqualiter declivi; extremitate anticâ rotundatâ (plerumque subattenuatâ), posticâ rotundato-acuminatâ; costâ umbonali et flexurâ ventrali inconspicuis; ligamento magno, elongato, prominente; dentibus primariis laceratis, antico laterali subapproximato, postico laterali minore, remoto.

Var. Testâ magis trigonâ, undique sulcis confertis excavatâ.

Oblong subtriangular, solid, subequilateral, of an extremely glossy enamel-like opaque white surface, which is smooth near the beaks, and grooved closely towards the but slightly convex ventral margin; dorsal line nearly equally sloping on each side, the edges very slightly convex; anterior extremity rounded (and in the typical examples rather attenuated), posterior end obtusely acuminate; fold and umbonal ridge obsolete; ligament large, elongated, and prominent; inside white; the primary teeth minutely jagged, and

one of them rather large, anterior lateral rather approximate, posterior lateral smaller and distant.

Tumbez, Peru.

Var. More triangular, the close concentric sulci occupying the entire surface. Chiriqui, West Columbia.

Its extraordinary teeth, large and elongated ligament, and either smooth or *closely* sulcated surface, suffice for its distinguishment.

41. TELLINA MADAGASCARIENSIS (pl. lxiii. f. 218.) *Gmelin*, 3237; Wood Gen. Conch. 169. t. 39. f. 2, 3; List. 386. f. 233; *T. rosea*, *Lam.* no. 14; *Hanl. Des. Cat.* p. 63. t. 4. f. 46.

T. testâ ovatâ, crassissimâ, convexâ, subimpolitâ, pallidè roseâ, lævigatâ; margine ventrali vix convexiusculo; dorsali utrinque valde declivi; costâ umbonali latâ, obtusâ; latere antico manifestè breviorè, postico subcuneiformi; extremitate anticâ obtusâ; superficie internâ roseâ, margine cardinali latissimo; dentibus lateralibus subobsoletis, parvis, antico approximato, postico remoto; impressionibus muscularibus fortiter impressis.

Ovate, very thick, convex, rather dull, darker or paler pink, smooth, ventral edge but little convex; anterior side somewhat the shorter, its extremity rounded, its slope moderate; posterior side subcuneiform, its extremity obtuse, its dorsal edge much sloping and slightly convex; umbonal ridge very obtuse; ligament prominent; inside rosy, scars deeply impressed; hinge-margin very broad; lateral teeth subobsolete, small, anterior one approximate, posterior distant.

Madagascar.

The figure in Lister is referred to by Spengler for his *T. rudis*, but neither the locality he gives, nor the expression "with coarse transverse striæ," will apply to this species.

42. TELLINA TONGANA (pl. lxiii. f. 209.) *Quoy* and *Gaimard*, *Astrolabe Zool.* vol. iii. p. 498. t. 81. f. 11, 12, 13.

T. testâ oblongâ, solidâ, subpellucidâ, subæquivalvi, convexâ, subæquilaterali, nitidâ, sublævigatâ, albâ (intus albidâ), roseo ad nates radiatâ, eodemque colore ad laterum extremitates ornatissimè pictâ; margine ventrali subarcuato, posticè sursum acclivi; dorsali anticè vix declivi, posticè leviter declivi convexo; latere postico paululum breviorè, attenuato, subrostrato; flexurâ ventrali valde conspicuâ; costâ umbonali prominente; umbonibus tumidis; ligamento infosso; dentibus lateralibus haud magnis, subæquidistantibus.

Oblong, solid, subequivalve, convex, subequilateral, rather pel-



lucid, nearly smooth, glossy, white, the beaks rayed with rose-colour, the extremities of the sides pink; ventral edge subarcuated, ascending posteriorly; anterior side very slightly the longer, its dorsal edge scarcely sloping; posterior edge attenuated and somewhat beaked, its dorsal side convex and moderately sloping; the flexure and umbonal ridge very conspicuous; umbones tumid; ligament sunken; inside whitish, a ligamental callus, and rather small and nearly equidistant lateral teeth.

Tonga.

43. *TELLINA ELEGANS* (pl. lxxv. f. 241.) *Gray* in *Wood's Ind. Test. sup. t. 1. f. 5*; *Hanl. Des. Cat. p. 62. t. 1. f. 5.*

T. testâ oblongo-elongatâ, valdè tortuosâ, subinæquivalvi, subæquilaterali, solidâ, politâ, subventricosâ, lævigatâ; nivêâ, (intus albidâ) roseo radiatâ, radiis latis versus marginem ventralem arcuatum et posticè acclinatum evanescentibus; margine dorsali anticè convexusculo et vix declivi, posticè leviter declivi et prope nates roseas subincurvato deinde convexo; latere postico paululum longiore, attenuato, leviter subrostrato; extremitate posticâ rotundato-angulatâ; ligamento brevi, pergrandi, prominente; costâ umbonali obsoletissimâ; dentibus parvis, antico subapproximato.

Twisted, elongated-oblong, solid, rather inequivalve, subequilateral, subventricose, smooth, highly polished, white with broad rosy rays, which disappear towards the arcuated and posteriorly ascending ventral margin; extremity of the anterior side rounded, its dorsal edge but slightly sloping; posterior side very slightly the longer, attenuated, obtuse at the extremity; umbonal ridge quite obsolete; ligament peculiarly large, short, and prominent; inside white, the lateral teeth small and rather approximate.

Gulf of Mexico.

— 44. *TELLINA RADIATA* (pl. lxxiii. f. 220, 221.) *Lin. Syst. 1117*; *Lam. no. 1*; *Chem. 6. f. 102*; *Wood, Gen. Con. t. 38. f. 2, 3*; *Ency. Mèt. t. 289. f. 2, 3*; *Hanl. Des. Cat. p. 61. t. 4. f. 26*; *Knorr. vol. iv. t. 2. f. 2, 4*; T. unimaculata, *Lam. no. 2.*

T. testâ oblongo-elongatâ, subinæquilaterali, haud tortuosâ, subventricosâ, solidâ, lævi, politâ, albâ, (intus flavescente) radiis roseis latis plerumque ornatâ; margine ventrali vix convexusculo; dorsali anticè vix declivi, posticè subrecto et leviter declivi; extremitate lateris antici longioris rotundatâ, lateris postici subattenuatâ, obtusè angulatâ; ligamento brevi, magno, prominente; costâ umbonali obsoletâ; flexurâ ventrali conspicuâ; lunulâ distinctâ, angusto-lanceolatâ; dentibus lateralibus validis, subæquidistantibus.

Var. testâ radiorum experte; natibus roseis.

Elongated-oblong, more or less ventricose, solid, not twisted, of a highly polished smooth white, with zones of very pale yellow, and generally with broad rosy rays; ventral margin scarcely convex; anterior side rather the longer, its dorsal edge scarcely sloping, its extremity rounded; posterior side slightly attenuated, its extremity obtusely subangulated, its dorsal edge nearly straight and but slightly sloping; ligament short, large, and prominent; lunule distinct, elongated-lanceolate; inside yellowish; lateral teeth strong and nearly equidistant.

West Indies.

45. *TELLINA BRASILIANA* (pl. lxii. f. 179.) *Spengler*, *Skriv. Nat. Selskab. Kiobenh.* 1798; *T. semizonalis*, *Lam.* no. 3; *Deles.* t. 6. f. 1; *Hanl. Des. Cat.* p. 61. sup. t. 11. f. 60.

*T. testâ oblongo-elongatâ, inæquilaterali, solidâ, convexiusculâ, nitidâ, pallidè roseo-violaceâ, lævigatâ; margine ventrali subrecto; dorsali anticè vix declivi et paululum convexiusculo, posticè declivi et subrecto; latere antico longiore, producto; extremitate posticâ obtusè angulatâ; costâ umbonali obtusâ; lunulâ lanceolatâ angustâ; superficie internâ roseo-purpurascente, radiis duobus pallidioribus aut albis ad extremitatem posticam ornatâ; dentibus lateralibus remotis, postico parvo.*

Elongated-oblong, inequilateral, solid, slightly convex, glossy, pale lilac, smooth; ventral margin nearly straight, slightly ascending in front, its flexure distinct; the umbonal ridge obtuse; anterior side much the longer, its dorsal slope very gentle; posterior extremity obtusely angulated, the dorsal edge nearly straight and moderately sloping; a narrow lanceolate lunule; the inside of a purplish crimson with two pale or white rays at the posterior extremity; teeth remote, equidistant, the posterior one small and rather indistinct.

Brazil (fide Spengler).

46. *TELLINA PUDICA* (pl. lvi. f. 7.) *Hanley*, *Zool. Proc.* 1844.

*T. testâ parvâ, solidâ, ovato-subtrigonâ, subæquilaterali, nitidâ, candidâ, sublævigatâ, tenuissimè concentricè striatâ; margine ventrali anticè arcuato, posticè subito sursum acclinato; dorsali utrinque valdè declivi, anticè convexo, posticè elongato et subrecto; latere antico paululum brevior, subventricoso, rotundato; postico compresso, rostrato; extremitate posticâ subacuminatâ; natibus prominentibus; flexurâ ventrali costâque umbonali subinconsipi-*

cuis; ligamento minimo, infosso; superficie internâ politâ; dentibus lateralibus distinctis, subapproximatis, subæquidistantibus.

Small, ovate, subtrigonal, nearly equilateral but rather longer posteriorly, rather ventricose in front but compressed at the posterior end, tolerably thick, of a very glossy pure uniform white, with scarcely impressed concentric striæ; the ventral edge bulging out towards the rounder anterior extremity, but rising in rather a concave line towards the posterior end, which is strongly beaked and has the tips rounded but much attenuated; front dorsal edge convex and strongly sloping, ligamental edge elongated, nearly straight and strongly sloping; beaks prominent; umbonal ridge and fold not conspicuous; ligament sunken and minute; inside polished, the lateral teeth distinct and nearly equally subapproximate.

Catbalonga, Isle of Samar.

47. *TELLINA SOWERBII* (pl. lxiii. f. 205.) *Hanley*, Zool. Proc. 1844.

T. testâ subellipticâ, subæquivalvi, tenui, compressâ, lævigatâ, politâ, candidâ; margine ventrali arcuato, posticè sursum acclinato; dorsali anticè vix paululum declivi, convexiusculo, posticè paululum declivi et prope nates subincurvato, deinde convexo; latere antico longiore, subproducto; postico rotundato-acuminato; ligamento parvo, angustissimo, infosso; flexurâ ventrali costâque umbonali subinconspicuis; superficie internâ albidâ, plerumque aurantio tinctâ; dentibus lateralibus tenuibus, subæquidistantibus.

Nearly elliptical, subinequivalve, compressed, highly polished, of an uniform smooth white (in some specimens with a tinge of rosy orange which increases in intensity towards the beaks); ventral edge arcuated, and rising at the posterior extremity; anterior side decidedly the longer, its extremity rounded, its dorsal edge nearly horizontal; posterior side attenuated, but rounded at its termination, its dorsal edge a little sloping at first, slightly incurved near the beaks, and then decidedly convex; ligament sunken, and very narrow; inside usually tinged with orange; lateral teeth thin, and nearly equidistant.

Australia.

48. *TELLINA LATA* (pl. lxiv. f. 237.) *Quoy et Gai*. Astrol. Zool. vol. 3. p. 497. t. 81. f. 8, 9, 10; Lam. 2nd. ed. vol. 6. p. 108; Hanl. Des. Cat. p. 71. sup. t. 14. f. 13.

T. testâ subovali, inæquilaterali, solidâ, subæquivalvi, compressiusculâ, nitidâ, albâ, sublævigatâ, rubro eleganter radiatâ; radiis

interruptis; margine ventrali convexo, utrinque (maximè autem posticè) sursum acclinato; dorsali posticè valdè declivi prope umbones subincurvato moxque subrecto, anticè convexiusculo et paulo declivi; latere postico brevi, acuminato; antico producto, ad extremitatem rotundato-obtuso; natibus acutis, rubro-roseis; ligamento infosso; superficie internâ albâ aut flavescente; dentibus lateralibus modicis, haud remotis.

Large, suboval, subequivalve, inequilateral, strong, rather compressed, nearly smooth or marked with concentric striulæ; of a glossy white with brilliant rays of a rosy red; ventral edge convex, rising at each extremity, but more particularly so at the posterior end which is short and acuminated; anterior side produced, its extremity obtusely rounded, its dorsal edge but slightly convex and little sloping; posterior dorsal edge greatly sloping, somewhat incurved near the rather acute beaks which are always rosy-red, and then nearly straight or but slightly convex; ligament sunken; inside usually tinged with yellow, lateral teeth moderately large and distant.

South Seas.

49. *TELLINA CHLOROLEUCA* (pl. lxiii. f. 208, 210.) *Lam.* no. 15; *Rumphius* t. 45. f. 1.

*T. testâ* ovatâ, subtenui, convexâ, nitidâ, albidâ aut albido-flavescente, radiis multis angustis pallidè roseis ornatâ, sublævigatâ, (lineis obsoletis radiantibus sæpe versus marginem ventralem strias concentricas confertissimas decussantibus); margine ventrali arcuato, posticè sursum acclivi; dorsali anticè convexo aut convexiusculo satisque declivi, posticè satis brevi convexiusculo et valdè declivi; latere antico longiore, ad extremitatem rotundato; postico brevi, acuminato; natibus roseis, prominulis; costâ umbonali et flexurâ ventrali conspicuis; ligamento angusto infosso; superficie internâ albido-flavescente; dentibus lateralibus, distinctis, subæquidistantibus.

Var. Radiis albidis et rubro-aurantiis pictâ; natibus vix roseis.

Ovate, convex, rather thin, of a glossy yellowish white, with numerous interrupted narrow rays of a pale rose colour, nearly smooth, (often decussated towards the arcuated and posteriorly rising ventral margin with concentric striulæ and obsolete radiating lines); anterior side the longer, its extremity rounded, its dorsal edge convex and moderately sloping; posterior side acuminated, very slightly beaked and rather short, the dorsal edge but slightly convex and much sloping; beaks of a deep rose-colour and rather prominent; ligament narrow and sunken; umbonal ridge and

flexure distinct; inside yellowish white; lateral teeth distinct and nearly equidistant.

Var. Rayed with orange-red and white; beaks but slightly if at all rosy.

Isle of Negros, Philippines (Cuming).

— 50. *TELLINA LÆVIGATA* (pl. lxiv. f. 225, 227.) *Lin. Sys.* 1117; *Lin. Mus. Ulr.* 480; *Schroet. einl. t. 7. f. 10*; *List. t. 387. f. 234*; *Lam. no. 36*; *Hanl. Des. Cat. p. 68*.

T. testâ ovato-rotundatâ, solidâ, subæquilaterali, nitidâ, convexâ, aut subventricosâ, sublævigatâ; margine ventrali anticè arcuato, posticè sursum acclinato; dorsali, anticè subdeclivi magis minusve convexo, posticè declivi et supernè subretuso aut subrecto et infernè convexo; latere antico paululum longiore, rotundato, dilatato; postico magis minusve angulato; natibus prominentibus; pube striis lamellosis rugosâ; ligamento infosso; flexurâ costâque umbonali conspicuis; cardine, dentibus lateralibus validis, subapproximatis, subæquidistantibus.

Var. *a.* Subalbidâ, radiis roseo-aurantiis ornatâ. Var. *b.* (T. concinna, *Phil. Ab. und Bes. Conch. part. 5. t. 5. f. 1.*) Albidâ, marginibus intus extusque rubro-aurantiis.

Rounded-ovate, solid, subequilateral, glossy, convex or subventricose, nearly smooth (there are obsolete crowded radiating lines, and frequently but not always minute very close raised concentric striulæ towards the lower margin and at either extremity); ventral edge arcuated in front, greatly rising behind; anterior side rather the longer, rounded and dilated, its dorsal edge more or less convex and not much sloping; posterior side more or less angulated, its dorsal edge sloping and nearly straight or slightly retuse above, but convex at its termination; beaks prominent; ligament sunken; fold and flexure conspicuous; lateral teeth strong, subapproximate, subequidistant.

Var. *a.* Whitish, or flesh-coloured with rosy orange rays. Var. *b.* White with the edges alone of a bright orange red, and the umbones and interior usually tinged with yellow.

An extremely variable species, with the beaks usually worn at the touching point.

St. Thomas, West Indies.

51. *TELLINA SEMEN* (pl. lvi. f. 8.) *Hanley, Zool. Proc.* 1844.

T. testâ ovali, crassâ, inæquilaterali, subventricosâ, nitidâ, albidâ (intus submargaritaccâ), anticè rotundatâ, posticè obtusâ; concentricè striatâ, striis anticè subimbricatis confertissimisque, posticè

remotioribus et elevatis; margine ventrali convexo; dorsali utrinque convexo aut convexiusculo, anticè subdeclivi, posticè declivi; latere antico multo longiore; natibus obtusis; ligamento minimo, prominulo; lunulâ angustissimâ; costâ umbonali nullâ, flexurâ subobsoletâ; dentibus lateralibus conspicuis, postico magis approximato.

Var. Ovatâ aut obovatâ, margine dorsali utrinque magis declivi.

Oval, thick, inequilateral, subventricose, glossy, whitish, with concentric striæ, which in front are crowded and subimbricated, but behind are rather distant and elevated; ventral edge convex; dorsal edges more or less convex, the hinder one sloping, the front one but moderately so, anterior side much the longer, rounded (and sometimes attenuated) at its extremity; posterior extremity obtuse; ligament very small and rather prominent; a very narrow lunule; beaks obtuse; flexure nearly obsolete; inside subnacreous; lateral teeth distinct, the hinder much the closer.

Var. Ovate or obovate, the dorsal edges more sloping.

Were it not for the slight fold, it might easily be mistaken for a *Donax*.

52. TELLINA BIMACULATA (pl. lvi. f. 16, 19, 20, 21, 22.) *Lin.*, Sys. 1120; Chem. vol. vi. f. 127, 132. Dillwyn. D. Cat. p. 101. Encyc. Méth. t. 290. f. 9, 10. Donovan, Brit. Shells, t. 19. f. 1. Turt. Biv. Brit. t. 8. f. 5; Lam. no. 52; Hanl. Des. Cat. p. 71. t. 4. f. 83; T. sexradiata *Lam.* no. 53.; Hanl. Des. Cat. p. 71. sup. t. 11. f. 37.

T. testâ ovatâ aut ovato-rotundatâ, crassâ, convexâ, nitidiusculâ, lævigatâ, variis coloribus pictâ; margine ventrali convexo aut arcuato (sæpe posticè paululum subincurvato); dorsali utrinque paululum convexiusculo, anticè valde declivi, posticè brevi leviterque declivi; latere antico longiore, attenuato, ad extremitatem rotundato; postico brevi, obtusè biangulato; costâ umbonali et flexurâ ventrali obsoletis; ligamento brevi, prominente; natibus acutis, nymphis prominentibus, incumbentibus; dentibus primariis permagnis; lateralibus brevibus, crassis, subæquidistantibus, (plerumque obsoletis).

Varying in shape from ovate to rounded ovate; thick, convex, slightly glossy, smooth; ventral edge varying from convex to arcuated, often slightly incurved posteriorly; hinder side much the shorter, obtusely biangulated, its dorsal edge very short and but slightly sloping; anterior side much attenuated, its extremity rounded, its dorsal edge much sloping and little convex, beaks inclined; ligament short but very prominent; primary teeth large,

the lateral (which however are usually obsolete), short, thick, and nearly equidistant.

The varieties in colouring are too numerous to specify, the principal however, are as follows; violet, orange-red, white, yellow or pinkish red, (either uniform, or rayed with purple, the white specimens with usually two lateral crimson internal spots); whitish or yellowish with narrow interrupted violet rays often assuming the appearance of arrow-heads; violet, rayed with white; whitish, with broad fawn-coloured and narrower purplish rays.

Britain, and West Indies.

53. *TELLINA LUCINOIDES* (pl. lvi. f. 32.) *Hanley*, Zool. Proc. 1844.

T. testâ suborbiculari, tenui, subæquilaterali, subventricosâ, impolitâ, intus extusque albâ, striis concentricis tenuissimis lineisque radiantibus minutissimis obsoletè decussatâ, utrinque rotundatâ; margine ventrali arcuato; dorsali utrinque convexiusculo et subæqualiter declivi; latere antico paululum longiore; flexurâ costâque umbonali obsoletis; ligamento angustissimo, infosso; dentibus lateralibus magnis, æquidistantibus.

Suborbicular, thin, subequilateral, rather ventricose, dull, both externally and internally white, with very delicate concentric striae and most minute radiating lineoles; ventral margin strongly arcuated; extremities rounded; dorsal edges nearly equally sloping and slightly convex; anterior side very slightly the longer; umbonal ridge and fold obsolete; ligament very narrow, sunken; lateral teeth distinct, equidistant.

Not unlike the shell which is figured without a name in the 292nd plate (f. 8.) of the *Encyclopédie Méthodique*.

Chain Island, South Seas.

54. *TELLINA NUX* (pl. lvi. f. 33.) *Hanley*, Zool. Proc. 1844.

T. testâ obovali, subinæquivalvi, subæquilaterali, tenui, convexâ aut subventricosâ, nitidâ, sublævigatâ aut internè concentricè substriatâ, intus extusque albidâ, umbonibus hyalinis et fulvis; margine ventrali subarcuato; dorsali utrinque convexo, satis et æqualiter declivi; extremitate posticâ obtusâ, attenuatâ; ligamento angustissimo, infosso; natibus obtusis; flexurâ distinctâ; costâ umbonali subinconspicuâ; dentibus lateralibus distinctis, antico paululum propinquiore.

Oboval, thin, slightly inequivalve, almost equilateral, from convex to subventricose, rather glossy, whitish both externally and

internally, with the umbones diaphanous and of a tawny orange-colour; nearly smooth or merely substriated concentrically towards the somewhat arcuated ventral margin; dorsal edges convex, equally and moderately sloping on either side; beaks obtuse; fold distinct; umbonal ridge almost obsolete; posterior side obtuse and attenuated; ligament extremely narrow, sunken; lateral teeth distinct, the anterior slightly the nearer.

Cagayan, Misamis, Mindanao; St. Nicolas, Zebu.

More oval than the three succeeding allied species.

55. *TELLINA PINGUIS* (pl. lvi. f. 34.) *Hanley*, Zool. Proc. 1844.

T. testâ parvâ, rotundato-ovali, tenui, subinæquivalvi, subæquilaterali, nitidâ, convexâ aut subventricosâ, intus extusque albidâ (rarius incarnatâ), lævigatâ (nonnunquam concentricè substriatâ); margine ventrali arcuato, dorsali utrinque convexo et subæqualiter declivi; extremitate posticâ obtusâ; flexurâ distinctâ; natibus minimis; costâ umbonali ferè obsoletâ; ligamento angustissimo, infosso; dentibus ut in *Nuce*.

Rounded-oval, thin, rather inequivalve, nearly equilateral, from convex to subventricose, both externally and internally whitish, or more rarely flesh-coloured, smooth, (sometimes concentrically substriated); the ventral edge arcuated; the dorsal on each side nearly equally sloping and convex; beaks minute; fold distinct; umbonal ridge almost obsolete; posterior end obtuse; ligament extremely narrow and sunken; teeth as in *Nux*.

St. Nicolas, Zebu.

By its more orbicular outline it may be distinguished from *Nux* and *Casta*. It is still more closely allied to *Robusta*, but its tenuity, more obtuse hinder extremity, and the greater convexity and less sudden slope of the dorsal edges, suffice for its separation.

56. *TELLINA ROBUSTA* (pl. lvi. f. 2, 3.) *Hanley*, Zool. Proc. 1844.

T. testâ parvâ, rotundato-subtrigonâ, solidâ, subinæquivalvi, subæquilaterali, ventricosâ, nitidâ, aut albo-flavescente, aut pallidè rosacèâ, sublævigatâ (plerumque inferius remotè substriatâ); margine ventrali valdè arcuato, dorsali utrinque subrecto et valdè declivi; latere postico attenuato, ferè subrostrato, ad extremitatem acuminato; ligamento angustissimo, infosso; natibus distinctis, prominentibus, et rectè incurvatis; umbonibus tumidis; flexurâ distinctâ; costâ umbonali ferè obsoletâ; lunulâ parvâ; superficie internâ, plerumque sub umbonibus, flavidâ aut rosèâ; dentibus ut in *Nuce*.



Rounded-subtrigonal, solid, rather inequivalve, nearly equilateral, ventricose, of a glossy yellowish white, or pale rose-colour, nearly smooth, or marked in fresh specimens with distant striulæ towards the extremely arched ventral margin; dorsal edges nearly straight and strongly sloping on either side; posterior side attenuated, almost beaked, and obtusely acuminate at the extremity; ligament extremely narrow and sunken; umbones tumid; beaks distinct, prominent, and incurved; fold distinct; umbonal ridge almost obsolete; inside usually yellow or rose-coloured at the umbones; a small lunule; teeth as in *Nux*.

Isle of Annaa, South Seas, and Isle of Burias, Philippines.

Is closely allied to *Nux*, *Pinguis* and *Casta*, but of a stouter texture even in the youngest individuals. The ligamental edge being nearly straight, easily separates it from *Pinguis*, where it is decidedly convex.

57. *TELLINA CASTA* (pl. lvi. f. 9,) *Hanley*, Zool. Proc. 1844.

T. testâ obovali, tenuissimâ, subinæquivalvi, subinæquilaterali, pellucidâ, convexâ, nitidâ, extus intusque candidâ, lævigatâ; margine ventrali maximè arcuato; dorsali utrinque subrecto, subæqualiter satisque declivi; extremitate posticâ angustâ, biangulatâ; ligamento angustissimo, subin fosso; natibus acutis; flexurâ ventrali costâque umbonali distinctis; dentibus ut in *Nuce*.

Oboval, extremely thin, slightly inequivalve, nearly equilateral, convex, glossy, pellucid, white, smooth; ventral margin extremely arcuated; the dorsal on either side nearly straight, moderately and nearly equally sloping; beaks acute; fold and umbonal ridge distinct; posterior extremity narrow and biangulated; ligament extremely narrow and scarcely, if at all, projecting; teeth as in *Nux*.

Singapore.

Bears some resemblance to *Nux*, but the shape is less broad, the ventral edge decidedly arcuated, the dorsal edges less convex and shorter, and the umbones colourless.

58. *TELLINA BALAUSTINA* (pl. lvi. f. 10.) *Lin.* Sys. 1119. *Poli.* Test. Sic. vol. i. t. 14. f. 17.; *Lam.* 2nd ed. vol. vi. p. 209; *Phil.* En. Mol. Sic. vol. i. p. 25; *Hanl.* Des. Cat. p. 72. sup. t. 9. f. 17; *Lucina balaustina*, *Payr.* Cat. Cors. t. 1. f. 21, 22.

T. testâ obovatâ, aut suborbiculari, solidiusculâ, convexâ, nitidiusculâ, albo-flavescente, radiis angustis rubris ornatâ, sublævigatâ, nonnunquam lineis elevatis concentricè striatâ; margine ventrali arcuato; dorsali utrinque convexo aut convexiusculo, satisque de-

clivi; latere antico vix paululum longiore, ad extremitatem rotundato; extremitate posticâ obtusissimè subangulatâ; natibus acutis, prominentibus, rectè incurvatis; ligamento angustissimo, infosso; umbonibus plerumque flavis; flexurâ ventrali costâque umbonali satis conspicuis; superficie internâ flavâ; dentibus lateralibus magnis, subæquidistantibus.

Shape varying from obovate to suborbicular, moderately strong, decidedly convex, when quite perfect dull and covered with raised membranaceous concentric striæ, but usually worn smooth and rather glossy; whitish or yellowish, with numerous narrow red rays and the umbones tinged with yellow; ventral edge arcuated; both edges of the dorsal line convex and moderately sloping; front extremity rounded; that of the scarcely shorter posterior side most obtusely subangulated; beaks prominent and incurvated; ligament extremely narrow and sunken; fold and umbonal ridge tolerably manifest; inside yellow; lateral teeth large and nearly equidistant.

Mediterranean.

59. *TELLINA OBLIQUILINEATA* (pl. lvi. f. 11, and pl. lix. f. 127.)  
*Conrad, Jour. Ac. N. S. Philad. 7.*

T. testâ parvâ, ovatâ, tenuiusculâ, inæquilaterali, convexâ, nitidâ, intus extusque flavescens aut rosaceo-incarnatâ, tenuissimè striatâ; striis obliquis et anticè flexuosis; margine ventrali subarcuato; dorsali anticè arcuato et declivi, posticè longiore convexiusculo et angulum acutum cum margine ventrali formante; latere antico brevi, rotundato; postico producto, acuminato; natibus ligamento que parvo prominulis; flexurâ costâque umbonali inconspicuis; dentibus lateralibus satis distinctis, subæquidistantibus, antico majore.

Small, ovate, rather thin, very inequilateral, glossy, convex, tinged both externally and internally with yellow or rosy flesh-colour, and most delicately but distinctly striated in oblique lines which become flexuous in front; the ligament small and scarcely projecting; the beaks rather prominent; the umbonal ridge and the fold not conspicuous; anterior side short and rounded, the dorsal slope rather sudden and very convex; posterior side produced, and rendered acuminated by the elongated and rather convex ligamental edge forming an acute angle with the rather arcuated ventral; lateral teeth distinct, neither very distant, the anterior decidedly larger.

Sandwich Isles.

60. *TELLINA AMPULLACEA* (pl. lvi. f. 23, 24.) *Philippi, (June*

1844) Ab. und Bes. Conch. part 5. t. 5. f. 7; *T. puella*, *Hanley*, Zool. Proc. (November) 1844.

*T. testâ inæquilaterali, tenui, ventricosâ, lævi, nitidiusculâ, extus intusque albido-roseâ aut pallidè carneâ; margine ventrali anticè arcuato, posticè sursum acclinato; dorsali anticè primum retuso et vix declivi deinde arcuato satisque declivi, posticè primum subrecto deinde convexiusculo et valde declivi; latere antico longiore, rotundato; postico brevi, et obtusè angulato; flexurâ costâque umbonali haud conspicuis sed satis distinctis; lunulâ parvâ, impressâ; natibus obtusis; ligamento angusto, prominulo; dentibus parvis; lateralibus remotis, subæquidistantibus.*

Ovate, thin, inequilateral, ventricose, smooth, rather glossy, pink or flesh-coloured both externally and internally; ventral edge arcuated in front, rising posteriorly; dorsal edge convex and but little sloping in front, slightly convex and much sloping behind; anterior side the longer, rounded; posterior side short, narrowed and angulated; umbonal ridge nearly obsolete, ventral flexure distinct; beaks obtuse; ligament rather prominent; teeth small, the lateral distant and nearly equally remote.

Senegal.

Dr. Philippi's publication had not reached me at the date of my describing this very distinct species.

- 61. *TELLINA STRIATULA* (pl. lxi. f. 175.) *Lam.* no. 33; *Enc. Mèth.* t. 292. f. 1; *T. hippoidea*, *Jonas*, Ab. und Bes. Conch. part 3. t. 5. f. 1.

*T. testâ sublenticulari, subobliquâ, paululum subinæquivalvi, solidiusculâ, subnitidâ, candidâ, sublævigatâ; margine ventrali valdè acuto, posticè sursum acclinato; dorsali anticè convexo, satisque declivi, posticè subrecto et valdè declivi; natibus acutis, prominentibus; flexurâ ventrali subinconspicuâ; areâ posticâ dorsali marginibus acutis costæ umbonalis circumseptâ, planulatâ aut subconçavâ; ligamento infosso; latere antico longiore, rotundato; extremitate posticâ breviter acuminatâ; superficie internâ niveâ; dentibus lateralibus magnis, antico approximato.*

Sublenticular, rather oblique, very slightly inequivalve, not very strong, convex, slightly glossy, white, nearly smooth, (being only marked with most indistinct minute radiating lines near the arcuated ventral margin which rising upward posteriorly almost forms a rectangle with the nearly straight and strongly sloping ligamental edge; front side the longer, extremity rounded, dorsal edge convex and moderately sloping; beaks acute and prominent, slightly curving

forwards; ligament sunken; edges of the almost terminal umbonal ridge acute, the intervening space flattened or concave; fold not conspicuous; inside pure glossy white; lateral teeth large, the anterior approximate.

St. Nicholas, Zebu.

The *Inflata* of Chemnitz (6. f. 86.) is very like this species, but I do not venture to assert its identity, as the figure certainly differs in some respects, and neither Lamarck nor Philippi, although both conversant with the figure, have quoted it.

62. *TELLINA DELTOIDALIS* (pl. lix. f. 128 and pl. lxiv. f. 229.)  
*Iam.* no. 49; *Hanl. Des. Cat.* p. 70. sup. t. 13. f. 3; *T. lactea*,  
*Quoy et Gai.* *Voy. Astrol. Zool.* vol. 3. p. 501. t. 81. f. 14, 15, 16;  
*Deles.* t. 6. f. 7.

*T.* testâ subtrigonâ, ovatâ, obovatâ, aut suborbiculari, solidâ, subinæquivalvi, convexâ, subæquilaterali, impolitâ aut vix nitidiusculâ, intus extusque albidâ, sublævigatâ; margine ventrali in medio subrecto, utrinque sursum acclivi; dorsali anticè vix declivi, posticè valdè declinente incurvato; latere postico subcuneiformi, ad extremitatem subangulato; extremitate anticâ obtusâ; natibus prominentibus, rectè incurvatis; flexurâ costâque umbonali distinctis; ligamento magno, infosso; dentibus lateralibus parvis, antico approximato, postico remoto plerumque subobsoleto.

Var. Latere antico haud dilatato; margine dorsali anticè magis declivi, posticè recto haudque incurvato.

Shell varying in shape from ovate-triangular to suborbicular-triangular, solid, slightly inequivalve, convex, subequilateral, scarcely if at all glossy and usually quite dull chalky white, nearly smooth or only marked with rather strong lines of growth; ventral margin sloping upward at both ends, nearly straight in the middle; fold and umbonal ridge distinct; beaks prominent and incurved; ligament large but sunken; anterior side dilated, obtuse, dorsal slope slight; posterior side somewhat wedge-shaped (at times almost subrostrated), its dorsal edge incurved and very strongly sloping, its extremity subangulated and much attenuated; lateral teeth small, the anterior approximate, the posterior distant and often almost obsolete.

Var. Ligamental edge not incurved but straight; anterior side not dilated, slope stronger, and extremity rounded.

Australia.

63. *TELLINA FAUSTA* (pl. lxiv. f. 230, 234) *Donovan*, vol. 3.  
t. 98; *Dor. Cat.* t. 5. f. 5; *Dillwyn D. Cat.* p. 94; *Hanl. Des. Cat.*

p. 68. t. 4. f. 68 ; *T. lævis*, *Wood*, Gen. Con. p. 181. t. 37. f. 1 ; *T. remies*, *Born*, p. 36. t. 2. f. 11 ; Lam. no. 31 ; List. t. 266. f. 102.

*T. testâ* ovato-suborbiculari, solidâ, convexâ aut subventricosâ, impolitâ, albidâ ; concentricè striatâ, striis tenuibus, confertissimis, ad umbones posticèque obsoletis ; lineis verticalibus interruptis fissuræformibus notatâ ; margine ventrali convexo vel subarcuato ; dorsali posticè subrecto (plerumque subincurvato) declivi, anticè prope umbones retuso, deinde convexo, satisque declivi ; latere postico brevior, subbiangulato ; extremitate anticâ rotundatâ ; natibus prominentibus, acutis ; ligamento maximo, prominente ; flexurâ ventrali satis distinctâ ; superficie internâ nitidâ, plerumque flavescente ; dentibus lateralibus magnis, antico approximato.

Var. Obovatâ, extus flavescente, intus flavâ.

Rounded-ovate, solid, varying from convex to subventricose, dull whitish, with vertical fissure-like interrupted lines, and crowded concentric fine striæ, which latter become obsolete on the umbones and posteriorly ; ventral edge varying from convex to subarcuated ; beaks prominent and acute, with a forward inclination ; fold tolerably distinct ; ligament very large, projecting ; anterior side rounded, rather the longer, the dorsal edge at first slightly concave and then convex and moderately sloping ; posterior end nearly biangulated, the dorsal edge nearly straight (usually incurved) and rather strongly sloping ; inside glossy, white, or tinged with yellow ; lateral teeth large, anterior approximate.

Var. Obovate, yellowish ; within deep yellow.

West Indies.

64. *TELLINA DISCUS* (pl. lxiv. f. 228, 232) *Hanley*, Zool. Proc. 1844.

*T. testâ* *T. remiei* simillimâ, sed subobliquâ et sulcis concentricis valdè irregularibus vixque continuis ; natibus haud rectè incurvatis ; margine dorsali posticè elevatiore, anticè prope nates prominulas subincurvato ; umbonibus lævigatis ; extremitate posticâ angulatâ.

Var. Testâ magis compressâ ; sulcis concentricis haud rudibus, et versus marginem ventralem solum conspicuis.

In sculpture, colour, and general shape extremely like *Remies*, but appearing oblique from the beaks having a forward inclination, and the anterior dorsal edge being slightly incurved ; posterior dorsal edge the more elevated, longer than in the last species ; posterior extremity distinctly angulated ; ventral edge slightly ascending posteriorly ; beaks rather prominent.

Var. More depressed, the sulci less coarse and only visible near the ventral margin.

Isle of Ticao, Philippines.

65. *TELLINA REMIES* (pl. lxiv. f. 226.) *Lin. Sys.* 1119; *Lin. Mus. Ulr.* p. 482; *Chem.* vol. 6. f. 113; *Wood, Gen. Conch.* p. 182. t. 38. f. 1; *Dillw. D. Cat.* p. 94; *Hanl. Des. Cat.* p. 67. sup. t. 1. f. 2; *T. sulcata, Lam.* no. 32; *T. convexa, Wood, Ind. Test. Sup.* t. 1. f. 2.

*T. testâ* suborbiculari, solidissimâ, inæquivalvi, æquilaterali, ventricosâ, subimpolitâ, extus intusque albidâ, epidermide tenui fulvo-cinereâ aut fulvo-fuscâ indutâ; supernè concentricè striolatâ, infernè concentricè sulcatâ, sulcis latis, profundis, rudibus; margine ventrali valdè arcuato; dorsali utrinque subrecto, anticè paululum, posticè satis declivi; latere antico rotundato; postico supernè subangulato; ligamento satis magno, prominente; lunulâ nullâ; natibus obtusis et rectè incurvatis; flexurâ ventrali conspicuâ; costâ umbonali subobsoletâ; dentibus magnis, validis, antico laterali quamplurimum approximato, postico permagno subremoto.

Suborbicular, very solid, inequivalve, equilateral, ventricose, scarcely glossy, whitish both within and without (covered when fresh with a dull yellowish brown epidermis), most delicately striated concentrically near the obtuse and incurved beaks, but very coarsely grooved towards the regularly arcuated ventral edge; anterior side rounded, its dorsal edge nearly straight or very slightly convex and but little sloping; posterior side subangulated above, its dorsal edge moderately declining and straight or subretuse; fold obscure, flexure conspicuous; ligament large and prominent; no lunule; teeth large and strong, the front lateral as close as possible, the hinder very large and rather distant.

Baclyon, Isle of Bohol (Cuming); Amboyna; Australia.

66. *TELLINA CYRENOIDEA* (pl. lvi. f. 31) *Hanley, Zool. Proc.* 1844.

*T. testâ* suborbiculari, solidiusculâ, inæquivalvi, valdè inæquilaterali, subventricosâ, sordidè albâ (intus sub umbonibus purpureâ), concentricè costellatâ; striis minutis obliquè radiantibus costellas confertas posticè decussantibus; margine ventrali arcuato; dorsali utrinque valdè et subrectè declivi, anticè brevi, posticè longiore; latere antico breviorè, obtusè rotundato; extremitate posticâ obtusâ, paululum attenuatâ; natibus prominentibus; lunulâ distinctâ; ligamento infosso; flexurâ costâque umbonali

obsoletissimis: dentibus lateralibus parvis, distinctis, subæquidistantibus.

Suborbicular, tolerably solid, inequivalve, very inequilateral, rather ventricose, hardly at all glossy, dirty white, with regular close-set narrow concentric ribs, which posteriorly are decussated by minute rather oblique radiating striulæ; ventral edge arcuated; anterior side the longer, its dorsal edge strongly sloping, rather long, nearly straight, its extremity rounded; posterior end obtusely rounded, slightly subangulated above, by the short straight and much sloping dorsal edge; beaks prominent; fold and umbonal ridge obsolete; ligament sunken; a purple internal stain beneath the umbones; lateral teeth small, but distinct, subequidistant and rather close.

St. Nicolas, Isle of Zebu.

67. *TELLINA SPLENDIDA* (pl. lvi. f. 39.) *Anton, Verz. Conch.* p. 5.

T. testâ suborbiculari, solidiusculâ, subinæquivalvi, inæquilaterali, convexâ, nitidâ, albido-flavescente, prope umbones subcærulescente aut purpureo tinctâ; anticè profundè et concentricè sulcatâ, dein striis subradiantibus remotis obliquè exaratâ, et in valvâ sinistrâ posticè concentricè striatâ; margine ventrali subarcuato; dorsali utrinque subrectè declivi, anticè brevi; extremitate anticâ obtusè rotundatâ, supernè angulatâ; latere postico longiore, ad extremitatem rotundatam subattenuato; ligamento infosso; costâ umbonali obsoletâ; flexurâ ventrali subinconspicuâ; superficie internâ purpureâ; dentibus lateralibus parvis, distinctis, subapproximatis, antico magis approximato.

Suborbicular, tolerably strong, rather inequivalve, inequilateral, glossy (usually, as in the succeeding species, coated here and there with a dark rust-like substance), yellowish white assuming a pale blueish or purplish cast towards the beaks; with strong rather distant concentric anterior grooves, followed by deep and not very close oblique radiating ones, which terminate posteriorly in one valve in fine close concentric striæ; ventral edge rather arcuated; anterior extremity obtusely rounded and rather angulated by the very short straight and much sloping dorsal edge; posterior side the longer, rather attenuated at its rounded extremity, the dorsal slope strong and nearly straight; ligament sunken and small; flexure not conspicuous; umbonal ridge obsolete; inside purple; lateral teeth small but distinct, close, anterior one the more so.

Basay, Isle of Samar; coarse sandy mud, four fathoms; (Cuming).

68. *TELLINA SENEGALENSIS* (pl. lvi. f. 17.) *Hanley, Zool. Proc.* 1844.

T. testâ *T. splendida* simillimâ, sed striis sulcisque exilioribus et magis confertis; utrâque valvulâ striis arcuatis obliquis ad extremitatem posticam notatâ; superficie internâ purpureâ, albo posticè biradiatâ.

Extremely like the last, but in addition to the superior delicacy and more crowded arrangement of both sulci and striæ, the posterior end is distinguished by the possession of those curved oblique striæ which we perceive less distinctly in *Carnaria*; the inner surface is adorned posteriorly with two white or pale rays on a purplish ground.

Senegal.

69. TELLINA CARNARIA (pl. lvi. f. 37, 38.) *Lin.* 1119; *Chem.* vol. 6. f. 126; *Wood, Gen. Conch.* t. 40. f. 4; *List.* t. 339. f. 176. *Hanl. Des. Cat.* p. 72. t. 4. f. 79; *Lucina carnaria, Lam.* no. 8; *Strigilla carnaria, Turt. Brit. Biv.* p. 117. t. 7. f. 15.

T. testâ suborbiculari, convexâ, subinæquivalvi, inæquilaterali, solidiusculâ, nitidâ, albido-roseâ, (intus roseâ), variè et exiliter striatâ, striis medio obliquis, radiantibus, ad latus posticum spectantibus; anticè flexuosis et cum striis mediis angulum formantibus; posticè tenuissimis, confertissimis, arcuatis (nonnunquam subobsoletis, et cum striis mediis angulum acutissimum formantibus); margine ventrali subarcuato; dorsali utrinque subrecto, declivi, anticè brevi, posticè elongato; extremitate anticâ obtusâ, supernè subangulatâ; latere postico valdè longiore, attenuato, ad extremitatem rotundato; flexurâ costâque umbonali obsoletis; dentibus lateralibus distinctis, subæquidistantibus.

Suborbicular, convex, moderately strong, glossy, pink, rose-coloured, or whitish more or less tinged with pink, most delicately striated in various directions; striæ on the anterior area very flexuous and forming an angle with those on the central area, which obliquely radiate posteriorward; striæ on the posterior slope extremely fine, curved, and meeting at a very acute angle; ventral edge arcuated; anterior side much the shorter, its extremity obtuse, and subangulated above, its dorsal edge short and moderately sloping, posterior extremity rounded and narrowed, the dorsal edge greatly sloping and nearly straight; a small lunule; ligament sunken; umbonal ridge and flexure nearly obsolete; beaks incurved; inside rosy; lateral teeth distinct and nearly equidistant.

W. Indies; W. Columbia.

The shell is deeper from the umbones to the ventral margin, than in *Splendida*; the posterior and central striæ are often divided by a smooth radiating space.



70. *TELLINA SINCERA* (pl. lxx. f. 144.) *Hanley*, Zool. Proc. 1844. 8

T. testâ *T. Carnariae* simillimâ, sed majore, latiore, compressâ et albidâ; striis tenuioribus; ligamento valdè angusto; natibus paululum ad latus anticum spectantibus; margine ventrali solum subarcuato; dentibus lateralibus conspicuis, subæquidistantibus.

Extremely like *Carnaria*, but larger, broader, white and depressed; the ligament extremely narrow; sculpture still finer, beaks slightly curved forward, ventral edge less arcuated; lateral teeth conspicuous and subequidistant.

N. W. Coast of America, (Dr. Sinclair).

✓ 71. *TELLINA PISIFORMIS* (pl. lvi. f. 30.) *Lin.* Syst. 1120.; *Dillwyn*. D. Cat. p. 102.; *Wood*. Gen. Con. p. 194.

T. testâ parvâ, suborbiculari, solidiusculâ, ventricosâ, nitidiusculâ, albidò-roseâ; obliquè substriatâ, striis posticè angulo acuto reflexis; margine ventrali convexo; dorsali anticè brevi subrecto et paululum declivi, posticè convexo et valdè declivi; extremitate anticâ obtusâ; latere postico longiore, rotundato; natibus prominentibus; umbonibus extus intusque roseis; flexurâ costâque umbonali obsoletis; dentibus lateralibus distinctis, subæquidistantibus.

Var. Testâ majore, pallidè luteâ.

Small, subglobose, tolerably strong, but slightly glossy, both within and without of a pinkish white with the umbones rosy, obliquely substriated, the striæ curving posteriorwards, except over the umbonal slope, (which, as well as the fold, is quite obsolete) where they are met at acute angles by another set with a slight anterior inclination; ventral edge simply convex; front dorsal edge very short, nearly straight, and scarcely sloping, the hinder one convex, and greatly sloping; anterior extremity obtuse; posterior side the longer and narrower, tip rounded; beaks very prominent; lateral teeth distinct, and nearly equidistant.

Var. Larger, and of an uniform pale yellow.

W. Indies.

The *Cardium discors* of Montagu, which has erroneously been inserted in the British Fauna, is evidently this or the succeeding species.

72. *TELLINA FLEXUOSA* (pl. lvi. f. 28, 29.) *Say*. Journ. Ac. N.S.; *Philad.* vol. ii. p. 303.; *Hanl.* Des. Cat. p. 2; *T. mirabilis*, *Philippi*, *Wieg.* Archiv. f. nat. 1841. p. 260.

T. testâ *T. pisiformi* simillimâ, pallidiore autem et margine ventrali magis arcuato; striis obliquis, fortioribus, flexuosis, posticè bis terve in ziczac flexis; ligamento minimo,

Extremely like *T. pisiformis*, but paler and more arcuated at the ventral margin; the oblique striæ are flexuous and more deeply impressed than in that species, and at the posterior extremity several times change their direction at acute angles, thus forming numerous zigzag lines; the ligament is minute and not prominent.

United States.

73. TELLINA FIMBRIATA (pl. lx. f. 132.) *Hanley*, Zool. Proc. 1844.

T. testâ obovato-suborbiculari, solidâ, convexâ, impolitâ, intus extusque candidâ, striis concentricis confertissimis lamellosis fimbriatis et lineis radiantibus confertis decussatâ; margine ventrali arcuato, posticè sursum acclinato; dorsali utrinque convexiusculo, anticè subdeclivi, posticè valdè declivi; latere antico longiore, rotundato; extremitate posticâ brevi, angulatâ; costâ umbonali valdè conspicuâ; ligamento infosso; lunulâ distinctâ; dentibus laterilibus subremotis, subæquidistantibus.

Rounded-obovate, solid, convex, very inequilateral, dull white, with extremely delicate crowded lamellar striæ, which are fringed by the intersection of radiating lines; one of the valves projecting beyond the other at the beaks; ventral margin arcuated and rising posteriorly; anterior side the longer and rounded, its dorsal edge moderately convex and sloping; posterior side short and angulated, the ligamental edge strongly sloping and tolerably convex; fold very prominent; ligament sunken; a distant lanceolate lunule; inside white; lateral teeth rather remote and subequidistant.

Mus. Cuming.

74. TELLINA DECUSSATA (pl. lx. f. 184.) *Lam.*, no. 45.; *Hanl. Des. Cat.* p. 70.; *Encyc. Mèth.* t. 292. f. 2.?

T. testâ suborbiculari, subæquilaterali, solidâ, convexâ aut subventricosâ, impolitâ, sordidè albâ, striis concentricis costellas radiantes decussantibus; striis elevatis, confertissimis, tenuissimis; costellis numerosis, paululum obsoletis; margine ventrali arcuato; dorsali utrinque declivi, anticè convexo, posticè convexiusculo; latere antico rotundato; postico obtusè angulato; ligamento infosso; natibus prominulis; flexurâ costâque umbonali inconspicuis; dentibus lateralibus magnis, antico propinquiore.

Var. Testâ majore, candidâ; striis magis elevatis et minus confertis; margine dorsali postico elevatiore.

Suborbicular, solid, nearly equilateral, from convex to subventricose, dull dirty white, sculptured with numerous somewhat obsolete radiating costellæ, which are decussated by very fine raised crowded concentric striæ; ventral edge arcuated; the dorsal sloping from the rather prominent beaks on either side, and rather the more convex in front; anterior side rounded, posterior obtusely subangulated; ligament sunken, fold and umbonal ridge not conspicuous; inside white; lateral teeth large, the front one the nearer.

Var. Larger, white; the hinder (and not as in the type,) the front), dorsal edge the more elevated; the striæ more elevated and rather more distant.

Australia; the variety from Ceylon.

75. *TELLINA CARNICOLOR*, (pl. lvi. f. 15.); *T. incarnata* Hanley, Zool. Proc. 1844.

*T. testâ obovatâ, subobliquâ, inæquilaterali, ventricosâ, solidâ, incarnatâ aut albido-roseâ, impolitâ; striis elevatis concentricis tenuissimis strias radiantes elevatas confertissimè decussantibus; margine ventrali arcuato, posticè sursum accliviore; dorsali anticè declivi et prope nates paululum incurvato, posticè elevatiore subarcuato et subito declivi; latero antico longiore, rotundato; extremitate posticâ rotundato-angulatâ; ligamento angusto, parvo, infosso; natibus acutis, prominentibus; flexurâ costâque umbonali subobsoletis; lunulâ lanceolatâ; superficie internâ flavescente, margines versus subroseâ; dentibus lateralibus maximis, subæquidistantibus.*

Obovate, rather oblique, dull, solid, ventricose, pinkish white, or flesh-coloured, closely decussated by elevated radiating, and by fine concentric lamellar striæ; ventral edge arcuated, and ascending posteriorly; anterior side the longer, its extremity rounded and slightly attenuated, its dorsal edge sloping, and near the acute and prominent beaks, somewhat incurved; hinder extremity subangulated, the more elevated ligamental edge sloping rather suddenly, and in an arcuated line; ligament small, narrow, and sunken; umbonal ridge and fold nearly obsolete; a lanceolate lunule; inside yellowish, pink near the margins; lateral teeth very large and nearly equidistant.

San Nicolas, Zebu.

In almost every adult specimen the tips of the beaks are chalky white, the umbones yellow, and the ligamental edge rosy.

I had inadvertently designated this shell with a pre-occupied name.

76. *TELLINA GARGADIA* (pl. lxi. f. 156 and pl. lxii. f. 181.)

*Lin. Sys.* 1116; *Lin. Mus. Ulr.* 476. *Chemn.* vol. vi. f. 63, 64; *Lam.* no. 40; *Hanl. Des. Cat.* p. 69. t. 3. f. 1.

T. testâ ovatâ, solidâ, compressiusculâ, nitidiusculâ, candidâ; margine ventrali arcuato; dorsali anticè subrecto et valdè declivi, posticè elevatiore valdè declivi paululum convexusculo dentato; latere antico longiore, angustato, rotundato; extremitate posticâ, obtusè biangulatâ; lunulâ areâque ligamentali excavatis; natibus acutis, prominentibus, ligamento infosso; dentibus lateralibus validis, æquidistantibus.

Var. *a.* Sulcis irregularibus, rudioribus, subobliquis, concentricis, posticè exarâtâ; lineis subobliquis, exilibus, concentricis, anticè striatâ. Var. *b.* Sculpturâ obsoletissimâ.

Ovate, solid, rather compressed, and not very glossy, uniform white; ventral edge arcuated; anterior side the longer, attenuated but rounded at its extremity; its dorsal slope sudden, and nearly straight; hinder side obtusely subangulated, its dorsal edge toothed, rather short and scarcely convex, the slope sudden; ligament sunken; lunule and lozenge excavated; beaks acute and prominent; fold and umbonal ridge by no means conspicuous; inside white, or yellowish white; the lateral teeth strong and subequidistant.

Var. *a.* Hinder side roughened by coarse irregular rather oblique concentric grooves; anterior surface traversed by more or less delicate rather obliquely concentric striæ. Var. *b.* Sculpture almost entirely obsolete, but usually discernible on one of the areas.

Siquijor, Philippines, (Cuming), &c.

77. *TELLINA SPINOSA* (pl. lxxii. f. 40.) *Hanley*, *Zool. Proc.* 1844.

T. testâ ovatâ, solidiusculâ, subimpolitâ, inæquilaterali, convexusculâ, extus intusque albidâ, striis minutis elevatis confertissimis concentricè asperatâ; margine ventrali arcuato, posticè sursum acclinato, dorsali posticè elevatiore convexo et declivi, antice prope nates acutas subincurvato, deinde subrecto subdeclivi; latere postico brevi; extremitate anticâ rotundatâ; extremitate posticâ spinarum duabus vel tribus seriebus radiantibus serratâ; lunulâ parvâ distinctâ; ligamento infosso; dente laterali antico subapproximato, postico remoto.

Ovate, moderately strong, inequilateral, not very convex, dull whitish, with minute elevated crowded concentric striæ; ventral edge rather arcuated, and rising behind; anterior side the longer, its extremity rounded, its dorsal edge slightly incurved near the

acute beaks and then straightish, (or slightly convex), and moderately sloping; posterior side short, and armed at its extremity with two or three rows of radiating serrated spines, its dorsal edge convex and sloping; ligament sunken; a small distinct lunule; inside white, the anterior lateral tooth slightly the closer, posterior distant.

Island of Ticao.

78. *TELLINA CRASSA* (pl. lxi. f. 169, 173.) *Pennant*, Brit. Zool. vol. iv. p. 73. t. 48. f. 28; Wood Gen. Conc. p. 186. t. 40. f. 1; Turt. Biv. Brit. p. 109. t. 7 f. 2; Lam. no. 35; Hanl. Des. Cat. p. 68. t. 4. f. 75; List. t. 299. f. 136; *T. rigida* *Don*. Br. Sh. vol. 3. t. 103.

T. testâ rotundato-ovali, subobliquâ, solidâ, inæquivalvi, subventricosâ, impolitâ, albidâ, radiis angustis rubris plerumque ornatâ, sulcis rudibus concentricè exaratâ; margine ventrali arcuato; dorsali utrinque declivi, convexo, posticè elevatiore, anticè convexiore; latere antico longiore, rotundato, prominente; postico abbreviato, ad extremitatem obtuso; costâ umbonali obsoletâ; natibus incumbentibus; lunulâ parvâ, excavatâ; ligamento infosso; flexurâ distinctâ; superficie internâ plerumque aurantiâ, (rarius albidâ); dentibus lateralibus magnis, antico subapproximato; cicatricibus muscularibus profundè impressis.

Roundish oval, slightly oblique, inequivalve, very inequilateral, rather ventricose, dull white, almost always adorned with narrow red rays, and concentrically and coarsely grooved; the ventral edge arcuated; the anterior side the longer, its dorsal edge arcuated, its extremity rounded and projecting; the posterior side rather short, its dorsal slope very sudden and very convex, its tips most obtusely subangulated; the umbonal ridge obsolete; ligament sunken; fold distinct; the beaks recurved anteriorwards; the lunule small and excavated; inside usually stained with orange, very rarely white; the lateral teeth large, the anterior one rather approximate; muscular impressions deeply impressed.

Britain; Europe.

The *T. maculata* of Turton (Biv. Brit.) is a young discoloured specimen of this species.

79. *TELLINA PLECTRUM* (pl. lxi. f. 177.) *Hanley*, Zool. Proc. 1845.

T. testâ obovatâ, solidâ, convexiusculâ, subæquilaterali, impolitâ, intus extusque albidâ, lamellis confertis concentricis undique operatâ; interstitiis lævibus; margine ventrali arcuato; dorsali utrinque subrecto, subdeclivi (in junioribus declivi); extremitate anticâ

rotundatâ; latere postico vix paululum longiore, biangulato; natibus acutis; costâ umbonali prominulâ; ligamento infosso (?); lunulâ distinctâ; dente laterali antico majore postico parvo et remotiore.

Obovate, solid, moderately convex, nearly equilateral, not polished, white, the entire surface covered with regular, distinct, short and crowded lamellæ, the interstices smooth; ventral edge arcuated; anterior side very slightly the shorter, extremity rounded; posterior extremity biangulated; dorsal edges straightish, and not greatly sloping; beaks acute; ligament sunken (?); fold not prominent; a distinct lunule; inside white, the front lateral tooth distinct, the hinder small and remote.

A single valve in the collection of J. E. Gray, Esq. is all I have ever seen of this distinctly characterized species, I am consequently unable to state whether it be equivalve or not.

17

80. *TELLINA SCOBINATA* (pl. lxxv. f. 235.) *Lin. Syst.* 1119; *Lin. Mus. Ulr.* 482; *Chem.* vol. 6. f. 122-124; *Wood, Gen. Conch.* p. 180. t. 35. f. 1; *Lam.* no 34; *Hanl. Des. Cat.* p. 68. t. 4. f. 67.

T. testâ suborbiculari, inæquivalvi, solidâ, æquilaterali, ventricosâ aut subventricosâ, impolitâ, albidâ, maculis brunneis undatis radiatâ aut nebulosâ, tuberculis, squamosis confertis undique asperatâ; margine ventrali arcuato; dorsali utrinque subdeclivi et leviter convexiusculo; extremitate anticâ rotundatâ, posticâ obtusè subangulatâ; costâ umbonali distinctâ, obtusâ; lunulâ excavatâ, lanceolatâ; flexurâ ventrali conspicuâ; ligamento depresso; disco interno albido-flavescente; dentibus lateralibus validdissimis, subæquidistantibus.

Suborbicular, solid, inequivalve, equilateral, more or less ventricose, dull whitish clouded or rayed with wavy brown markings, and rough with small crowded rasp-like tubercles which become almost spinous beyond the distinct but obtuse umbonal ridge; fold conspicuous; ligament depressed; a lanceolate excavated lunule; anterior side rounded, the dorsal edge scarcely convex and little sloping; hinder side with the extremity obtusely subangulated, the dorsal edge nearly straight and moderately sloping; inside yellowish white; the lateral teeth very strong and subequidistant.

Society Islands and Philippines; (Cuming).

82. *TELLINA LINGUA-FELIS* (pl. lxxiv. f. 236.) *Lin. Sys.* 1116; *Lin. Mus. Ulr.* p. 476; *Knorr, vol. 2. t. 2. f. 1*; *Wood, Gen. Conch.* t. 40. f. 2, 3; *Lam.* no. 37; *Hanl. Des. Cat.* p. 69. t. 3. f. 2.

T. testâ obovatâ, subventricosâ, solidâ, subæquilaterali, albidâ,

roseo radiatâ, impolitâ, squamis tubercularibus confertis undique asperrimâ; margine ventrali posticè arcuato, anticè sursum acclivi; dorsali anticè convexiusculo et paululum declivi, posticè elevatiore convexo et valde declivi; latere postico breviorè, subrostrato; extremitate anticâ rotundatâ; natibus acutis, roseis aut roseo-purpureis, incumbentibus; lunulâ excavatâ, lanceolatâ; costâ umbonali et flexurâ conspicuis; ligamento infosso; dentibus lateralibus validis, subæquidistantibus.

Obovate, rather ventricose, crowded with rasp-like scales, solid, white with crimson rays and the beaks of the same colour; anterior side rather the longer, its extremity rounded, its dorsal edge slightly convex and little sloping; hinder side slightly beaked, the ligamental edge the more elevated convex and deeply sloping; ventral edge much arcuated in front, ascending posteriorly; flexuosity and umbonal ridge conspicuous; beaks acute, inclined forwards; lunule deeply impressed, lanceolate; lateral teeth strong, subequidistant.

Isle of Negros, Philippines; (Cuming).

82. *TELLINA RUGOSA* (pl. lxiv. f. 233, 238.) *Born*, p. 29. t. 2. f. 3, 4; Wood, *Gen. Conch.* t. 41. f. 2, 3; Lam. no. 38; Hanl. *Des. Cat.* p. 69. t. 3. f. 8.

T. testâ obovatâ, inæquivalvi, subæquilaterali, solidâ, subventricosâ, intus extusque albâ (rarius roseo-purpurascente aut aurantiâ); subconcentricè rugosâ, rugis erectis flexuosis interruptis, striis radiantibus minutissimis decussantibus; margine ventrali anticè arcuato; dorsali anticè paululum declivi, posticè subrecto et valdè declivi; latere postico obtusè subcuneiformi; lunulâ elongatâ, excavatâ; flexurâ costâque umbonali conspicuis; dentibus lateralibus magnis, subæquidistantibus.

Obovate, subequilateral, inequivalve, solid, rather ventricose, white both within and without (rarely orange or purplish crimson), with raised interrupted irregularly but concentrically disposed wavy wrinkles which become still rougher from the minute radiating striæ; anterior side rounded, its dorsal edge scarcely sloping; posterior side obtusely subcuneiform; ligamental edge strongly sloping and nearly straight; ventral edge much arcuated in front; lunule very elongated, excavated; fold and flexure conspicuous; lateral teeth large, subequidistant.

Isle of Oparâ (Cuming), New California.

83. *TELLINA SUBTRUNCATA* (pl. lvii. f. 48.) *Hanley*, *Zool. Proc.* 1844.

T. testâ obovatâ, solidiusculâ, convexâ, impolitâ, valdè inæquilaterali, intus extusque striis lamellosis confertissimis fimbriatis ornatâ; margine ventrali anticè arcuato, posticè subrecto et sursum acclinato; dorsali utrinque magis minusve convexo, anticè declivi, posticè maximè declivi; extremitate anticâ rotundatâ; latere postico brevissimo, subtruncato, angulato; ligamento infosso; natibus prominentibus; costâ umbonali subinconspicuâ, terminali; dentibus lateralibus subæquidistantibus.

Obovate, moderately thick, convex, very inæquilateral, dull whitish, with very crowded fringed lamellar striæ; ventral margin arcuated in front, straightish and ascending behind; dorsal edges moderately convex; anterior side the longer, rounded, its dorsal edge sloping; posterior side subtruncated, its extremity angulated; beaks prominent; fold terminal and not prominent; ligament sunken; inside white, sometimes with two orange rays; lateral teeth subequidistant.

Isle of Bohol, Philippines.

84. *TELLINA PRISTIS* (pl. lxi. f. 160.) *Lam.* no. 41; *Hanl. Des. Cat.* p. 69. sup. t. 9. f. 4. (young); *Enc. Mèth.* t. 287. f. 1.

T. testâ rotundato-ovatâ, crassâ, subinæquilaterali, convexâ aut subventricosâ, extus intusque albâ, impolitâ; concentricè striatâ, striis elevatis, confertis; interstitiis, striulis minimis radiantibus, decussatis; margine ventrali magis minusve convexo; dorsali anticè brevi, subretuso et paulo declivi, posticè subrecto et valdè declivi; latere antico paululum longiore, ad extremitatem rotundato-obtuso; latere postico angulato; natibus acutis; lunulâ ligamentoque infosso magnis; flexurâ costâque umbonali satis conspicuis; dentibus lateralibus subremotis, subæquidistantibus.

Rounded-ovate, thick, rather inequilateral, convex or subventricose, white both within and without, not glossy, with concentric elevated crowded striæ decussated by minute interstitial radiating striulæ; ventral edge more or less convex; dorsal short subretuse and but little sloping in front, nearly straight and considerably sloping behind; anterior side very slightly the longer, obtusely rounded at its termination; posterior side angulated; beaks acute; ligament sunken, and as well as the lunule, large; fold and flexure tolerably strong; lateral teeth rather and nearly equally distant.

Australia, &c.

85. *TELLINA CAPSIDES* (pl. lxii. f. 185.) *Lam.* no. 44; *Hanl. Des. Cat.* p. 70.

T. testâ obovato-subtrigonâ, solidâ, subæquilaterali, convexius-



culâ, vix nitidâ, intus extusque albâ aut albidâ; concentricè striatâ, striis posticè elevatioribus, anticè a lineis radiantibus distinctis decussatis; margine ventrali subarcuato; dorsali utrinque valdè declivi, anticè convexo, posticè recto aut subrecto; extremitate anticâ rotundatâ; latere postico angulato; flexurâ costâque umbonali satis distinctis; ligamento magno, infosso; dentibus lateralibus distinctis, remotis, validis, subæquidistantibus.

Obovate-subtriangular, solid, subequilateral, moderately convex, scarcely glossy, white or whitish, anteriorly decussated by closely radiating and concentric striæ, posteriorly with the concentric striæ more distant and more elevated; ventral edge subarcuated; dorsal slopes strong, the posterior nearly straight and roughened by the edges of the concentric sulci; posterior tip angulated; beaks acute; anterior extremity rounded; ligament sunken, and large; fold and flexure moderate; inside white; lateral teeth strong, subequidistant.

Jimaimalan, Isle of Negros, &c.; (Cuming).

86. *TELLINA PERPLEXA* (pl. lx. f. 139.) *Hanley*, Zool. Proc. 1844; *T. concinna*, (?) *Ab. und Bes. Conch.* part 5. t. 5. f. 1.

*T. testâ T. ostracea* affinis, ovatâ autem, striisque ejus concentricis confertioribus, et supernè haud lamellosis; margine dorsali antico paululum declivi.

Very like *T. ostracea* but subovate and not triangular; the concentric striæ more crowded and not lamellar, except near the less arcuated ventral margin; the anterior dorsal edge but little sloping.

Bay of Manila and Isle of Annaa; (Cuming).

87. *TELLINA OSTRACEA* (pl. lvii. f. 45.) *Lam.* no. 54; *Hanl. Des. Cat.* p. 71. sup. t. 14. f. 11; *Enc. Mèth.* t. 290. f. 13.

*T. testâ* obovato-subtrigonâ, solidiusculâ, satis convexâ, extus intusque albidâ, lineis elevatis subremotis concentricè ornatâ; interstitiis lævibus; margine ventrali subarcuato; dorsali utrinque declivi, anticè convexo, posticè recto aut subretuso; extremitate anticâ rotundatâ; latere postico paulo breviorè cuneiformi; natibus acutis, prominentibus; ligamento infosso; costâ umbonali prominente; dentibus lateralibus, distinctis, postico subremoto.

Obovate-subtriangular, tolerably strong and convex, uniform white, with regular raised rather distant concentric striæ, the interstices smooth; ventral edge arcuated, sloping upward posteriorly; anterior side rather the longer, its dorsal edge convex and sloping, its extremity rounded; posterior side subrostrated and wedge-

shaped, its dorsal edge straight (or subretuse) and much sloping; beaks prominent, acute; ligament sunken; umbonal ridge very distinct; inside white; lateral teeth distinct, the hinder slightly the more distant.

Ceylon.

88. *TELLINA PLICATA* (pl. lxii. f. 191.) *Valenciennes, Enc. Mèth.* t. 287. f. 3.

T. testâ obovali, tenuiusculâ, subventricosâ, tortuosâ, æquilaterali, impolitâ, intus extusque candidâ, striis elevatis lamellosis concentricè ornatâ; margine ventrali arcuato, posticè sursum acclinato; dorsali utrinque paululum declivi, anticè convexo (sed juxta lunulam brevem distinctam incurvato), posticè convexiusculo et elevatiore; extremitate anticâ rotundatâ, obtusè biangulatâ; costâ umbonali subcarinatâ flexurâque ventrali conspicuis; ligamento longo, infosso; dentibus lateralibus parvis, antico subapproximato.

Oboval, rather thin, somewhat ventricose, tortuous, nearly equilateral, dull white, with regular raised and moderately distant concentric sublamellar striæ; the ventral margin arcuated, and rising behind; anterior side very slightly the longer, its extremity rounded, its dorsal edge near the short but distinct lunule incurved and scarcely sloping; extremity of the posterior side obtusely biangulated, its dorsal slope the more elevated slightly convex and moderately sloping; ligament elongated and sunken; umbonal ridge subcarinated, and as well as the ventral fold, very distinct; inside white; lateral teeth small, the anterior the more approximate.

Hab.? Mus. Cuming, Metcalfe.

89. *TELLINA CRYSTALLINA* (pl. lvii. f. 43.) *Chem.* vol. 11. p. 210. f. 1947, 1948; *Hanl. Des. Cat.* p. 66. t. 3. f. 10; *Wood, Gen. Conch.* p. 149.

T. testâ obovatâ, tenui, inæquivalvi, inæquilaterali, convexiusculâ, nitidâ, submargaritaceâ, candidâ, subpellucidâ, lamellis subremotis concentricis undique ornatâ; interstitiis lævibus; margine ventrali valdè arcuato, posticè brevi et incurvato; latere antico longiore, rotundato; postico brevi, rostrato; natibus acutis; ligamento minimo, infosso; costâ umbonali distinctâ; dentibus lateralibus distinctis, subæquidistantibus.

Obovate, thin, inequivalve, inequilateral, moderately convex, glossy, pure white, subnacreous, with distant regular raised concentric striæ, the interstices quite smooth; ventral edge greatly

arcuated; anterior side much the longer, rather dilated, rounded, its dorsal edge convex and not greatly sloping; posterior side terminating in a short beak, which is placed very high up owing to the shortness and small slope of the incurved dorsal edge; fold prominent; ligament minute and sunken; beaks acute; inside white; lateral teeth distinct, subequidistant.

St. Elena, West Columbia. *W. J.*

90. *TELLINA BURNETI* (pl. lviii. f. 99.) *Broderip* and *Sowerby*, *Zool. Jour.* vol. 4. p. 362. t. 9. f. 2; *Hanl. Des. Cat.* p. 72. t. 13. f. 51.

T. testâ trigonâ, suborbiculari, subtenui, subæquilaterali, inæquivalvi, compressâ, candidâ, submargaritaceâ; valvulâ planulatâ lineis elevatis concentricè striatâ; valvulâ convexiore sulcis dilatatis concentricis obsolete anticè et supernè ornatâ; margine ventrali arcuato; dorsali utrinque valde declivi et dentato, anticè valdè incurvato, posticè subrecto; natibus acutissimis, anticè incumbentibus; ligamento minimo lunulâque maximâ infossis; flexurâ distinctâ; dentibus lateralibus distinctis, subæquidistantibus.

Triangular, rather thin, inequivalve, subequilateral, compressed, subnacreous, white; the convex valve with obsolete distant dilated concentric grooves, which are most visible in front and towards the elevated acute and curving beaks (one of which projects over the other); the flat valve with elevated fine rather irregular and close concentric striæ, which become obsolete posteriorly; ventral edge arcuated; dorsal edges strongly sloping on either side, and armed with large tooth-like projections, the front one greatly incurved, the hinder nearly straight; ligament minute; dorsal slopes excavated; fold and flexure distinct; lateral teeth distinct, rather remote, and subequidistant.

Salango, W. Columbia.

91. *TELLINA LYRA* (pl. lxii. f. 187.) *Hanley*, *Zool. Proc.* 1844.

T. testâ ellipticâ, tenui, compressâ, nitidiusculâ, intus extusque albâ, striis concentricis elevatis ornatâ; striis anticè et infernè confertioribus, interstitiis lævigatis; margine ventrali medio convexiusculo, utrinque arcuato; dorsali posticè elevatiore convexo satisque declivi, anticè prope lunulam excavatam, aut horizontali aut leviter acclivi; latere antico paululum longiore, rotundato; extremitate posticâ obtusâ; ligamento infosso; natibus acutis, prominentibus, incumbentibus; flexurâ obsoletâ; dentibus lateralibus distinctis, antico approximato, postico parvo, remotiore.

Elliptic, thin, depressed, scarcely glossy, white both within and without (sometimes subnacreous), with regular elevated concentric striæ, which are distant towards the very prominent and acute beaks, but become closer towards the ventral edge (which is rounded at both extremities, but is little convex in the middle); the interstices of the striæ quite smooth; anterior side very slightly the longer, the extremity rounded; situation of the umbonal ridge indicated by a linear carina almost adjacent to the convex and moderately sloping dorsal edge and separated from it by a narrow concavity; the beaks curved forwards; the anterior dorsal edge the less elevated, and at first running almost horizontally or even slightly ascending; lunule and nymphal area excavated; front lateral tooth distinct and subapproximate, the hinder one small and distant.

Probably inequivalve, but as yet I have only seen one of the valves.

Tumbez, Peru.

92. *TELLINA GOULDII* (pl. lvi. f. 26.) *Hanley*.

T. testâ parvâ, obliquè subovatâ, solidâ, convexiusculâ; nitidissimâ, extus intusque niveâ, lævigatâ, valdè inæquilaterali; margine ventrali anticè fortiter arcuato, posticè sursum acclinato; dorsali antico subdeclivi, prope nates acutas retuso, deinde arcuato; dorsali postico subrecto, subitoe declivi; latere postico brevissimo, obtusè angulato; extremitate anticâ rotundatâ; lunulâ parvâ; flexurâ costâque umbonali subobsoletis; ligamento minimo infosso; dente laterali antico parvo et haud approximato, postico rudimentali.

Obliquely subovate, solid, small, moderately convex, very glossy, quite smooth and both within and without of a pure white; ventral margin much arcuated in front, ascending posteriorly; front dorsal edge moderately sloping, retuse near the acute beaks and then arcuated or convex; hinder dorsal edge nearly straight and suddenly sloping; sides very unequal, the hinder peculiarly short and obtusely angulated; front extremity rounded; fold and flexure almost obsolete; ligament minute and sunken; a small lunule; front lateral tooth small and not approximate, hinder almost rudimentary.

West Indies; (Guilding).

Named in honour of the talented author of the 'Report on the Invertebrata of Massachusetts'.

93. *TELLINA RUBICUNDA* (pl. lxvi. f. 255.) *Gould*, Proc. Boston Nat. Hist. Soc. 1845.

T. testâ oblongo-ovali, inæquivalvi, compressâ, tenui, subnitidâ, inæquilaterali, pallidè aurantio-rosçâ, radiis subroseis ornatâ; sublævigatâ, concentricè exiliter striolatâ, valvulâ autem alterâ sulcis remotis concentricis posticè exaratâ; margine ventrali medio subrecto; dorsali anticè convexiusculo paululumque declivi, posticè subrecto et subdeclivi; latere postico brevi, angulato; extremitate anticâ rotundato-obtusâ; ligamento prominulo; costâ umbonali subinconspicuâ; dentibus lateralibus distinctis, subæquidistantibus.

Oblong-oval, inequivalve, compressed, thin, not very glossy, inequilateral, of a very delicate salmon-colour, with light pink rays; smoothish, being merely striolated concentrically, with a posterior space grooved concentrically and distantly in one of the valves only; ventral edge straightish in the middle; dorsal slightly convex and very little sloping in front, moderately sloping and nearly straight behind; posterior side short and angulated; anterior extremity obtusely rounded; ligament moderately projecting; fold by no means conspicuous; lateral teeth distinct, and tolerably equidistant.

\*\* With a single anterior lateral tooth.

Africa, (Dr. Bates).

94. *TELLINA MARS* (pl. lxii. f. 180.) *Hanley*.

T. testâ oblongâ, solidâ, subæquilaterali, compressiusculâ, extus intusque rosçâ, concentricè et confertissimè striatâ, lineisque obsolete radiatim impressâ; margine ventrali in medio subrecto, utrinque convexo, posticè incurvato; dorsali anticè convexiusculo et vix paululum declivi, posticè prope nates obtusas excavato deinde convexo et subdeclivi; extremitate anticâ rotundatâ; latere postico obtusè acuminato, subrostrato; flexurâ costâque umbonali valdè conspicuis; ligamento infosso; dente laterali quamplurimum approximato.

Oblong, solid, nearly equilateral, rather compressed, rose-coloured within and without, decussated by very crowded concentric striæ and obsolete radiating lines; ventral edge nearly straight in the middle, convex at the sides, incurved posteriorly; dorsal edge but slightly convex and scarcely at all sloping in front, posteriorly excavated near the obtuse beaks and then convex and moderately sloping; front extremity rounded; posterior side subrostrated and obtusely acuminated; fold and flexure very conspicuous; ligament sunken; lateral tooth as approximate as possible.

The only specimen I have ever seen, (now in the possession of Mr. Cuming,) is stated to have come from Guinea.

95. *TELLINA FOLIACEA* (pl. lxiv. f. 253.) *Lin. Sys.* 1117; *Lin. Mus. Ulr.* p. 479; *Chem.* vol. 6. f. 95; *Wood, Gen. Conch.* t. 36. f. 1; *Lam.* no. 12; *Hanl. Des. Cat.* p. 62. t. 3. f. 20; *Knorr.* vol. v. t. 29. f. 2.

*T. testâ oblongâ, tenui, planulatâ, nitidissimâ, subæquilaterali, aurantiâ, sublævigatâ, concentricè et obsoletè substriatâ, et supra carinam umbonalem alterâ in valvulâ punctis squamosis radiatâ; margine ventrali subrecto; latere antico subattenuato, rotundato; postico biangulato; margine ligamentali convexo, paululum declivi, et spinis lanceolatis armato; umbonibus valdè compressis; natibus inconspicuis; ligamento infosso; dente laterali approximato, parvo.*

Oblong, thin, and flattened, very glossy, subequilateral, bright orange-yellow, nearly smooth but striated in front with very fine concentric lines, and roughened in one valve beyond the subcarinated umbonal ridge by radiating rows of scaly dots; ventral edge nearly straight; anterior side rounded, and rather attenuated; posterior side obtusely biangulated; ligamental edge convex, but little sloping, elongated, and armed with lanceolate spines; ligament elongated and sunken; beaks not prominent; umbones much flattened, plicated; inside somewhat tinged with purple; lateral tooth small and approximate.

San Nicolas, Zebu (Cuming); Amboyna; Moluccas.

96. *TELLINA MAGNA* (pl. lxiv. f. 239, and pl. lxiii. f. 201. young.) *Spengler*, loc. cit. (1798); *T. acuta*, *Wood*, *Gen. Conch.* p. 157. t. 44. f. 1; *Dillw. Des. Cat.* p. 81; *Hanl. Des. Cat.* p. 63. t. 3. f. 25; *T. elliptica*, *Lam.* no. 16.

*T. testâ oblongâ, solidâ, inæquilaterali, compressâ, nitidissimâ, lævigatâ, albidâ, pallidè aurantio supernè radiatâ (in junioribus aurantiâ, radiis angustis pallidioribus ornatâ); margine ventrali convexo, posticè sursum acclinato; dorsali anticè convexo et paululum declivi, posticè supernè subretuso satisque declivi; latere antico longiore, rotundato; postico angustato, angulato; natibus roseis, compressis; ligamento infosso; flexurâ costâque umbonali satis conspicuis; dente laterali subapproximato.*

Oblong, solid, inequilateral, compressed, smooth and polished, white with pale orange rays near the umbones; the beaks compressed and rosy (in the young the surface is deep orange with narrow paler rays); ventral edge convex and rising posteriorly; anterior side the longer, rounded at the extremity, its dorsal edge more or less convex and but little sloping; posterior side narrow and angu-

lated, its dorsal edge (in the adult) retuse above and moderately sloping; ligament sunken; fold and flexure distinct; interior whitish tinged with orange; lateral tooth tolerably close.

West Indies; (Wood).

97. *TELLINA SOL* (pl. lxiv. f. 240.) *Hanley*, Zool. Proc. 1844.

T. testâ oblongo-ellipticâ, solidiusculâ, compressâ, nitidâ, rubro-aurantiâ, alterâ valvulâ concentricè substriatâ, alterâ sublævigatâ; margine ventrali convexo, posticè sursum acclinato; dorsali utrinque subdeclivi, convexiusculo; latere antico longiore, ad extremitatem rotundato; extremitate posticâ in junioribus subacuminatâ, in adultis obtusè angulatâ; natibus planulatis; ligamento infosso; dente laterali distincto.

Oblong-elliptic, tolerably strong, compressed, glossy, orange red (rayed in some specimens with paler streaks); one valve with concentric striulæ, the other nearly smooth; ventral edge simply convex, ascending posteriorly; dorsal edges moderately sloping, and slightly convex; anterior side the longer, its extremity rounded; hinder extremity somewhat acuminate in the young, obtusely angulated in the adult; beaks flattened and devoid of colour; ligament sunken; lateral tooth distinct.

Hab. ?

The adult is in the cabinet of Mr. Cuming, the young in that of Mr. Metcalfe.

98. *TELLINA STRIGOSA* (pl. lxiii. f. 211.) *Gmelin*, 3239; *Dillw. Des. Cat.* 82; *Lam. no.* 19; *Hanl. Des. Cat.* p. 63. t. 4. f. 61; *Le Vagal, Adanson*, Seneg. t. 17. f. 19.

T. testâ ovato-oblongâ, solidâ, inæquivalvi, convexiusculâ, nitidâ, candidâ, subpellucidâ, sublævigatâ; margine ventrali subrecto, medio paululum subretuso, posticè sursum acclinato; dorsali anticè arcuato declivi et elevatiore, posticè prope nates subretuso et vix declivi, deinde convexiusculo et obliquo; latere postico longiore, ad extremitatem attenuatam subrostrato; extremitate anticâ rotundatâ; flexurâ costâque umbonali satis conspicuis; ligamento infosso; superficie internâ candidâ, plerumque roseo subradiatim tinctâ; dente laterali approximato, plerumque obsoleto.

Ovate-oblong, solid, inequivalve, moderately convex, of a glossy subpellucid white, with the marks of growth tinged with zones of violet grey, nearly smooth, but the convex valve posteriorly grooved near the beaks with most delicate but distinct crowded striæ; ventral margin somewhat retuse in the middle, rising behind; anterior side rather the shorter, its extremity rounded, its dorsal edge

arcuated; hinder extremity attenuated and subrostrated; fold and flexure distinct; the ligamental edge at first incurved, and then oblique and slightly convex; within often stained with rose colour; ligament sunken; the approximate lateral tooth almost or entirely obsolete.

Senegal.

99. *TELLINA PLANATA* (pl. lxi. f. 174.) *Lin. Sys.* 1117; Born, t. 2. f. 9; Lam. no. 20; Philip. En. Mol. Sic. vol. 1. p. 26; Hanl. Des. Cat. p. 63. t. 3. f. 24; Poli. vol. 1. t. 14. f. 1 to 5.

VAR. (?) *T. flavescens operculata*, *Chem.* vol. 6. f. 98.

*T. testâ ovali, solidiusculâ, inæquivalvi, æquilaterali, convexiusculâ, nitidâ extus intusque albido-carneâ, sublævigatâ; margine ventrali convexiusculo, medio subrecto; dorsali anticè convexiusculo paululumque declivi, posticè subdeclivi et prope ligamentum infossum subretuso; latere antico rotundato; postico subangulato; flexurâ costâque umbonali satis conspicuis; dente laterali parvo, approximato.*

Oval, tolerably strong, inequivalve, equilateral, moderately convex, nearly smooth, pale flesh colour; ventral margin moderately convex, but nearly straight in the middle; anterior side rounded, its dorsal edge convex and scarcely at all sloping; posterior extremity subangulated; the ligamental margin moderately sloping, retuse near the sunken ligament; fold and flexure tolerably distinct; inside flesh-coloured; lateral tooth small, approximate.

Mediterranean.

The shell represented by Chemnitz (vol. vi. f. 98) is probably a variety. It is slightly more elongated, with the radiating lines more distinct.

100. *TELLINA IMBELLIS* (pl. lx. f. 155.) *Hanley, Zool. Proc.* 1844.

*T. testâ oblongo-ovali, solidâ, inæquivalvi, inæquilaterali, convexâ, nitidâ, extus intusque albidâ, valvulâ convexiore striis elevatis concentricis posticè notatâ; valvulâ alterâ lævigatâ; margine ventrali convexo, posticè sursum acclinato; dorsali anticè convexiusculo paululumque declivi, posticè magis minusve declivi et subretuso; latere antico producto, rotundato; postico brevi, obtusè angulato; flexurâ costâque umbonali obsoletis; ligamento prominente; dente laterali parvo, approximato.*

Oblong-oval, solid, inequivalve, inequilateral, convex, rather glossy, whitish (inside white), smooth, excepting upon the posterior side of the more convex valve, which is finely striated by raised



concentric lines; ventral margin convex and rising behind; anterior side produced, its dorsal edge tolerably convex and but little sloping; posterior side short and obtusely angulated; ligamental edge straightish or subretuse, more or less sloping; ligament prominent; fold and flexure obsolete; lateral tooth small, approximate.

Hab.?

The unique specimen which I purchased without a locality, is now in the possession of H. Cuming, Esq.

101. *TELLINA MERA* (pl. lxvi. f. 268.) *Say*, Amer. Conch. pl. 64. f. 2.

T. testâ ovato-suborbiculari, inæquilaterali, convexâ aut convexiusculâ, vix nitidâ, haud solidâ, extus intusque albâ, lineis exilibus elevatis concentricè asperatâ; margine ventrali convexo et posticè sursum subacclinato; dorsali utrinque declivi, anticè convexo, posticè subrecto; latere postico breviorè, et inferne obtusissimè angulato; extremitate anticâ rotundatâ; natibus acutis; ligamento subinfosso; flexurâ costâque umbonali haud conspicuis; dente laterali haud magno, haud approximato.

Rounded ovate, moderately inequilateral, more or less convex, not solid nor highly polished, white both within and without, most delicately striated by raised concentric lines, (which however are scarcely visible in the young); ventral margin convex, and somewhat rising posteriorly; dorsal edges sloping, the front one more or less straight, the hinder decidedly convex; posterior side the shorter, very obtusely angulated below; anterior side rounded; beaks acute; ligament rather sunken; fold and flexure just evident; lateral tooth neither large nor approximate.

Guayaquil; south coast of U. States.

102. *TELLINA CYCLADIFORMIS* (pl. lvii. f. 61.) *Hanley*, Zool. Proc. 1844.

T. testâ rotundato-subtrigonâ, tenui, ventricosâ, intus extusque incarnatâ aut pallidè roseâ, sublævigatâ, aut concentricè tenuissimè striolatâ; margine ventrali convexo; dorsali utrinque declivi, convexiusculo; latere antico rotundato, paulo breviorè; extremitate posticâ obtusè subangulatâ; ligamento prominulo; flexurâ costâque umbonali obsoletis; dente laterali parvo, approximato.

Small, subtriangularly orbicular, thin, ventricose, nearly smooth, (or marked with most minute concentric striæ), both within and without of a pale rose or flesh-colour; ventral edge convex, both dorsal edges sloping and moderately curved; anterior side rather the shorter, rounded; hinder extremity obtusely subangulated;

ligament tolerably large, and somewhat prominent; fold and flexure obsolete; lateral tooth small, approximate.

San Nicolas, Zebu.

103. *TELLINA LISTERI* (pl. lxxv. f. 251.) *Hanley, Zool. Proc.* 1844; *Lister, Hist. Conc. t. 288. f. 235*; *T. scutra Gould, Proc. Boston Nat. H. Soc.*

*T. testâ obovatâ, solidâ, ventricosâ, (anticè subventricosâ), æquilaterali, glabrâ, extus intusque candidâ; margine ventrali medio subrecto; dorsali anticè arcuato paululumque declivi, posticè recto et declivi; latere antico dilatato, obtusè rotundato; posticè obtusissimè biangulato; ligamento magno, infosso; natibus obtusis; umbonibus plerumque subplanulatis; flexurâ costâque umbonali satis distinctis; dente laterali parvo subapproximato.*

Obovate, solid, more or less ventricose, equilateral, white both externally and internally, almost smooth, (being merely marked with the concentric wrinkles of growth), rather glossy; the ventral edge nearly straight in the middle; the anterior side dilated, its extremity obtusely rounded, its dorsal edge scarcely sloping and much arcuated; hinder extremity most obtusely biangulated, the dorsal edge straight and much sloping; umbonal ridge and ventral flexure not very conspicuous; ligament large and sunken; umbones rather flattened; beaks obtuse; lateral tooth small, and nearly approximate.

Senegal.

104. *TELLINA INÆQUALIS* (pl. lvii. f. 44.) *Hanley, Zool. Proc.* 1844.

*T. testâ subovatâ, valdè inæquilaterali, solidâ, convexâ, candidâ, tenuiter striatâ; striis supernè obliquis, infernè concentricis, flexuosis; supra costam umbonalem inconspicuam, rugis erectis flexuosis asperatâ; margine ventrali convexissimo; dorsali anticè subincurvato et valdè declivi, posticè brevi recto subdeclivi; latere antico producto, ad extremitatem attenuato, rotundato; extremitate posticâ obtusâ; natibus acutis; lunulâ distinctâ; superficie internâ candidâ, aut flavescente; dente laterali magno, subremoto.*

Subovate, very inequilateral, solid, convex, white with fine striæ, which are oblique near the acute beaks, but become concentric and flexuous towards the very convex ventral margin; the surface beyond the umbonal ridge (which is not prominent), covered with elevated flexuous oblique wrinkles, which project so as to form three radiating lines; anterior side produced, attenuated but rounded at the extremity, its dorsal line rather concave and greatly

sloping; hinder side very short, its extremity obtuse but slightly angulated above, its dorsal edge moderately sloping; beaks acute; a distinct lunule; inside yellowish; lateral tooth moderately distant.

Ceylon.

105. *TELLINA PUMILA* (pl. lvii. f. 41.) *Hanley*, Zool. Proc. 1844.

T. testâ *T. Philippinarum* simillimâ, sed angustiore; margine ventrali medio subrecto; dorsali utrinque recto aut subconcavo, anticè paululum declivi; margine antico recto, verticali; latere postico cuneiformi.

Bearing a very close resemblance to *T. Philippinarum*, but decidedly narrower; both dorsal edges (except in the young) straight or even concave, and the front one which is longer and less sloping than in the next species, forming an angle with the straight and direct anterior margin; the hinder extremity wedge-shaped, the ventral margin (which is nearly straight in the middle) not being elevated posteriorly.

Valparaiso.

106. *TELLINA PHILIPPINARUM* (pl. lvii. f. 55, 56.) *Hanley*, Zool. Proc. 1844.

T. testâ ovatâ aut subovatâ, tenui, subæquilaterali, intus extusque candidâ, (sæpe umbones versus incarnatâ aut pallidè aurantiâ), submargaritaceâ, concentricè et tenuissimè striatâ; margine ventrali arcuato, posticè sursum acclivi; dorsali anticè brevi recto subdeclivi, posticè declivi et convexiusculo; extremitate anticâ obtusâ; latere postico subcuneiformi, aut obsoletè subangulato; ligamento prominulo; concavitate dorsali anticâ lunulam simulantè; dente laterali subapproximato, (in junioribus subobsoleto).

Ovate or subovate, thin, subequilateral, rather depressed, whitish (with a pearly gloss), often tinged near the beaks with flesh-colour or pale orange, most delicately striated concentrically; ventral edge arcuated, inclining upward towards the posterior side, which varies from subcuneiform to obsoletely subangulated; anterior side very slightly the longer, its dorsal edge slightly sloping and not convex, its extremity obtuse; an obsolete lunule; a distinct subapproximate lateral tooth which is nearly obsolete in the young.

Philippines.

107. *TELLINA CULTER* (pl. lvii. f. 51.) *Hanley*, Zool. Proc. 1844.

T. testâ parvâ, ovatâ, inæquilaterali, tenuiusculâ, convexâ, nitidâ, intus extusque aurantio-roseâ, lævigatâ; margine ventrali anticè arcuato, posticè sursum acclivi; dorsali anticè magis minusve convexo satisque declivi; latere antico producto, ad extremitatem obtusè rotundato; postico acuminato; natibus acutis; ligamento vix prominulo; costâ umbonali ferè et flexurâ ventrali omnino obsoletis; dente laterali parvo, approximato.

Small, ovate, inequilateral, thinnish, tolerably convex, both externally and internally of a rosy-orange, nearly smooth, in young specimens there are indications of concentric striæ near the arcuated front of the ventral edge, which latter ascending posteriorly in nearly a straight line, acuminates the short anterior side at its junction with the straight and much sloping ligamental margin; front dorsal edge always convex and not greatly sloping, extremity obtusely rounded; ligament scarcely prominent; ventral flexure entirely obsolete, umbonal ridge nearly so; beaks acute; lateral tooth small, approximate.

Mindanao, Philippines.

108. *TELLINA CORBULOIDES* (pl. lvii. f. 50, 57.) *Hanley*, Zool. Proc. 1844.

T. testâ subovali, inæquivalvi, solidâ, subventricosâ, sublævigatâ, roseo-incarnatâ, epidermide tenuissimâ et opali coloribus nitente indutâ; margine ventrali sinistræ valvulæ ultra marginem convexiusculum alterius prominente; margine dorsali anticè convexiusculo et declivi, posticè subrecto et valdè declivi; latere antico brevior, obtusè acuminato; extremitate posticâ rotundatâ; ligamento prominulo; areâ dorsali posticâ in adultis planulatâ; natibus obtusis; flexurâ ventrali distinctâ; superficie internâ plerumque aurantio-rubrâ; dente laterali parvo, approximato.

Var. *b.* Oblongâ; margine ventrali subrecto; flexurâ obsoletâ.  
Var. *c.* Extus intusque candidâ.

Suboval, inequivalve, solid, nearly smooth, more or less ventricose, covered when fresh with a very thin opalescent epidermis, under which the shell is dull, and of a rosy flesh colour; ventral edge moderately convex and in the left valve projecting beyond that of the right; anterior side shorter, and obtusely acuminated, its dorsal edge nearly or quite straight and much sloping; hinder side rounded, its dorsal edge convex and moderately sloping; hinder dorsal area flattened in the adult; beaks obtuse; ventral flexure distinct; inside usually orange-red; a single small approximate lateral tooth.

Var. *b.* Oblong, the ventral edge nearly straight; flexure obsolete. Var. *c.* Entirely white.

Catbalonga, Philippines, and Bay of Manila (var. *b.*).

109. *TELLINA RHODORA* (pl. lix. f. 105.) *Hinds*, Zool. Sulphur, vol. 2. t. 21. f. 3. p. 67.

T. testâ oblongo-triangulari, solidâ, convexâ aut subventricosâ, subæquilaterali, nitidissimâ, lævi, extus intusque albido-roseâ, marginibus roseis; margine ventrali convexiusculo, et anticè convexo; dorsali utrinque subrecto, anticè subdeclivi; latere postico brevior, cuneiformi; extremitate anticâ obtusè rotundatâ; flexurâ subobsoletâ; ligamento depresso; dentibus primariis parvis; laterali antico maximo et subapproximato, postico rudimentali et subremoto.

Oblong-triangular, solid, rather ventricose, nearly equilateral, perfectly smooth and glossy, within and without of a pale pink, becoming rose-coloured near the margins; ventral edge convex in front, dorsal edges nearly straight, the anterior one scarcely sloping; hinder side the shorter, wedge-shaped; front extremity obtusely rounded; ligament depressed; primary teeth small, front lateral close and very large, a rudimentary remote hinder one.

Straits of Macassar.

110. *TELLINA FELIX* (pl. lviii. f. 52.) *Hanley*, Zool. Proc. 1844.

T. testâ subovali, solidiusculâ, valdè inæquilaterali, convexiusculâ, nitidâ, intus extusque roseâ, lævigatâ; margine ventrali vix convexiusculo; dorsali anticè vix declivi convexo, posticè valdè declivi; latere postico brevissimo, obtusè subtruncato, infernè obtusè angulato; extremitate anticâ rotundatâ; costâ umbonali et flexurâ ventrali subobsoletis; ligamento brevi, prominulo; dente laterali magno, approximato.

Suboval, tolerably strong, very inequilateral, slightly convex, both within and without of a glossy rose-colour, quite smooth; the ventral edge nearly straight; the anterior side produced and rounded, its dorsal edge convex and little sloping; hinder side extremely short, and obtusely but rather obliquely subtruncated; umbonal ridge and fold, nearly obsolete; ligament rather prominent; a single large approximate lateral tooth.

Panama.

111. *TELLINA HILARIS* (pl. lvii. f. 54.) *Hanley*, Zool. Proc. 1844.

*T. testâ oblongo-cuneiformi, tenui, compressiusculâ, inæquilaterali, nitidâ, lævi, intus extusque roseâ, albo biradiatâ; radiis latis submediis; margine ventrali convexiusculo (flexurâ obsoletâ); dorsali anticè vix convexiusculo et subdeclivi, posticè subrecto et valdè declivi; extremitate lateris antici longioris rotundatâ; extremitate posticâ brevi, cuneiformi; dente laterali parvo, subapproximato.*

Oblong-wedge-shaped, thin, rather compressed, inequilateral, glossy, smooth, rosy both within and without, with an anterior and posterior subcentral broad white ray; ventral margin slightly convex and devoid of flexure; anterior side the longer, its extremity rounded, its dorsal edge scarcely convex and little sloping; hinder extremity short and wedge-shaped, the ligamental edge nearly straight; the lateral tooth small and nearly approximate.

Red Sea (M. Clery.)

112. *TELLINA HIBERNA* (pl. lvii. f. 53.) *Hanley, Zool. Proc.* 1844. ~~p. 445~~

*T. testâ oblongâ, solidâ, compressiusculâ, valdè inæquilaterali, subnitidâ, extus intusque candidâ, lævigatâ; margine ventrali subrecto; dorsali anticè convexiusculo et paululum declivi, posticè primùm convexo deinde subincurvato; latere antico producto rotundato; postico brevi, cuneiformi; flexurâ subobsoletâ; ligamento prominulo; dente laterali, magno, approximato.*

Oblong, solid, rather compressed, very inequilateral, rather glossy, white both within and without, smooth; ventral edge straightish; front dorsal margin slightly convex and scarcely sloping, hinder one first convex and then somewhat incurved; anterior side produced and rounded; posterior short and wedge-shaped; flexure nearly obsolete; ligament rather prominent; lateral tooth large, approximate.

Panama; Bay of Guayaquil.

113. *TELLINA POLITA* (pl. lvii. f. 60.) *Say, Jour. Ac. N. S. Philad.* 2. p. 276.; *Say, Amer. Conch.* t. 65. f. 2.; *Hanl. Des. Cat.* p. 65. *Sup. t.* 9. f. 39.

*T. testâ T. Teneræ affini, minus autem inæquilaterali; extremitate posticâ acutâ, cuneiformi; margine ligamentali, recto.*

Allied to *T. Tenera*, but less inequilateral, the ligamental edge straight, and the hinder extremity acute and wedge-shaped.

Massachusetts to Georgia, North America.

114. *TELLINA TENERA* (pl. lvii. f. 59.) *Say, Jour. Ac. N. S. Philad.* 2. p. 303.; *Hanl. Des. Cat.* p. 65. *sup. t.* 9. f. 38.; *Gould Invert. Massa.* f. 44. p. 68.

T. testâ oblongo-ovali, tenui, inæquilaterali, compressiusculâ, nitidâ, intus extusque candidâ, submargaritaceâ, sublævigatâ; margine ventrali convexiusculo; dorsali anticè subrecto paululumque declivi, posticè declivi et vix convexiusculo; latere antico producto, rotundato; extremitate posticâ obtusè cuneiformi; flexurâ obsoletâ; ligamento prominulo; dente laterali distincto, approximato.

Oblong-oval, thin, inequilateral, rather compressed, and both within and without of a nacreous white, nearly smooth or with delicate and almost obsolete concentric striæ; ventral edge moderately convex; dorsal straightish and but little sloping in front, slightly convex and decidedly sloping behind; anterior side produced and rounded; posterior extremity very obtusely wedge-shaped; flexure obsolete; ligament slightly prominent; lateral tooth distinct and approximate.

North America.

— 115. *TELLINA INCARNATA* (pl. lx. f. 142. pl. lxvi. f. 265.) *Lin. Syst.* 1118; (from his collection); *Poli. 1. t. 15. f. 1.*; *T. depressa Lam. no. 22.*; *Wood Gen. Con. t. 45. f. 3.*; *Philip. En. Moll. Sic. 1. p. 27.*; *Hanl. Des. Cat. p. 63. t. 4. f. 48.*; *Turt. Biv. Brit. p. 105. t. 8. f. 6.*; *T. squalida Mont. Test. Brit. p. 56.*

T. testâ oblongo-ovali, subtenui aut solidiusculâ, subcompressâ, subæquilaterali, subnitidâ, albido-aurantiâ and roseo-aurantiâ, concentricè substriatâ, striis alterâ in valvulâ obsoletis; margine ventrali convexo; dorsali anticè convexiusculo paululumque (in junioribus vix minimè) declivi, posticè subrecto aut subretuso et subdeclivi; latere antico longiore, rotundato; extremitate anticâ angulatâ subrostratâ; flexurâ satis distinctâ; ligamento infosso; coloribus superfiei internæ plerumque saturatioribus; dente laterali subapproximato.

Oblong-oval, rather thin or but moderately strong, somewhat compressed, subequilateral, more or less glossy, of a pale orange or deep flesh colour, with (in the young chiefly,) two whitish approximate posterior rays, concentrically substriated in one of the valves; ventral edge convex; anterior side rounded and rather the longer its dorsal edge slightly convex and but little sloping; posterior extremity angulated and subrostrated; the ligamental edge straightish or subretuse and moderately sloping; ligament sunken; fold and flexure tolerably distinct; internal colouring usually darker; lateral tooth subapproximate.

Britain, and Mediterranean.

116. *TELLINA VALTONIS* (pl. lvii. f. 68.) *Hanley, Zool. Proc.* 1844.

T. testâ subovali, tenui, compressâ, subæquilaterali, nitidâ, albido-roseâ radio angusto pallidiore (nonnunquam duobus) posticè ornatâ, sublævigatâ, pellucidâ, anticè rotundatâ; margine ventrali convexiusculo; dorsali utrinque convexiusculo, posticè declivi, anticè subdeclivi; latere postico paululum breviorè, obtusè cuneiformi; flexurâ costâque umbonali obsoletis; ligamento prominulo; dente laterali parvo, approximato.

Suboval, thin, compressed, subequilateral, pink with one or two narrow posterior whitish rays, nearly smooth, glossy, pellucid, rounded in front, rather shorter and obtusely wedge-shaped behind; ventral and dorsal edges tolerably convex, very moderately sloping in front, but more so behind; fold and flexure quite obsolete; ligament moderately prominent; lateral tooth small, approximate.

Hab. ? Mus. Metcalfe.

117. *TELLINA EXILIS* (pl. lix. f. 104.) *Lam.* no. 26; Deshayes, *Ency. Mèth.* vol. iii. p. 1013; *Hanl. Des. Cat.* p. 64. sup. t. 13. f. 2; *Deles.* t. 6. f. 6.

T. testâ subovali, tenuissimâ, compressâ, nitidâ, inæquilaterali, pellucidâ, roseâ, radio uno vel altero posticè ornatâ, minutissimè striatâ; striis anticè concentricis, medio obliquis; margine ventrali subrecto; dorsali anticè convexiusculo et subdeclivi, posticè subrecto et declivi; latere postico cuneiformi et paulo breviorè; extremitate anticâ rotundatâ; flexurâ costâque umbonali obsoletis; ligamento prominulo; dente laterali parvo, subapproximato.

Suboval, very thin, pellucid, compressed, rose-coloured, with two narrow white posterior rays, nearly smooth being only marked with very delicate and almost imperceptible regular striæ, which, concentric in front, become oblique on reaching the middle of the shell; ventral margin nearly straight; anterior side rather the longer, rounded, its dorsal edge not much sloping and slightly convex; posterior end wedge-shaped, the ligamental edge nearly straight; fold and flexure obsolete; ligament moderately prominent; lateral tooth small, approximate.

Hab. ? Mus. Walton.

118. *TELLINA VIRGO* (pl. lvii. f. 42.) *Hanley*, *Zool. Proc.* 1844.

T. testâ subovali, tenuissimâ, compressiusculâ, inæquilaterali, nitidâ, pellucidâ, candidâ, striis obliquis subremotis regularibus ornatâ; margine ventrali subrecto; dorsali anticè convexo et subdeclivi, posticè subrecto et declivi; latere antico longiore,



rotundato; extremitate posticâ obtusè cuneiformi; flexurâ costâque umbonali subobsoletis; ligamento satis prominente; dente laterali distincto, subapproximato.

Suboval, inequilateral, rather compressed, very thin, pellucid, white, with regular and not very close oblique striæ; ventral edge nearly straight; dorsal margin more or less convex and not greatly sloping in front, nearly straight and moderately sloping behind; anterior side the longer, rounded; posterior extremity obtusely wedge-shaped; ligament moderately prominent; fold and flexure obsolete; lateral tooth distinct, subapproximate.

Chiriqui, West Columbia.

119. *TELLINA SIMILIS* (pl. lvii. f. 65.) *Sowerby*, Brit. Miscel. pl. 75; Mont. Test. Brit. Sup. p. 167; Turt. Biv. Brit. p. 102; Philip. Ab. und Bes. Conch. part 9. pl. 5. f. 7.

T. testâ subovali, solidiusculâ, inæquilaterali, convexâ, nitidâ, extus intusque albidâ, concentricè et oblique striatâ; striis obliquis, minutis, confertissimis; ultimâ parte posticâ alterâ in valvulâ sublævigatâ, in alterâ autem striis subimbricatis haudque confertis concentricè exaratâ; margine ventrali subrecto; dorsali anticè convexiusculo, posticè brevi subretuso, utrinque paululum declivi; latere antico longiore, rotundato; extremitate posticâ obtusè biangulatâ; costâ umbonali satis conspicuâ; ligamento prominente; dente laterali magno, approximato.

Suboval, moderately strong, inequilateral, convex, glossy, whitish both within and without, with a few concentric and crowded minute oblique striæ, which do not extend to the obtuse hinder extremity, but are replaced before reaching the distinct umbonal ridge, by not very close subimbricated striæ (obsolete in one of the valves); ventral edge straightish; dorsal not greatly sloping on either side, convex in front, short and retuse behind; anterior side the longer, rounded; ligament prominent; lateral tooth large, approximate.

West Indies.

120. *TELLINA DECORA* (pl. lvi. f. 27., <sup>not</sup>lix. f. 127., and <sup>not</sup>lxvi. f. 260.) *Say*, Jour. Acad. Philad. vol. v.

T. testâ *T. simili* simillimâ, sed minore, magis convexâ, magis inæquilaterali, radiis angustis roseis ornatâ; margine cardinali roseo sub ligamento.

Very like a young *Similis* (of which it may possibly prove a variety), but decidedly more convex than specimens in that stage, and when adult much more inequilateral than full grown shells of

the preceding species; delicately coloured by narrow lighter or darker tinged crimson rays, and the hinge margin stained with rose-colour beneath the ligament; ventral edge decidedly convex.

Honduras (Dyson); Florida.

121. *TELLINA IRIS* (pl. lxvi. f. 267.) Say, Jour. Ac. Nat. Sc. Philad. vol. ii.

*T. testâ subovatâ, solidiusculâ, valdè inæquilaterali, nitidâ, satis convexâ, extus intusque albidâ, anticè radio subluteo obsoletè pictâ, striisque obliquis confertis utrâque in valvulâ insculptâ; margine ventrali anticè subarcuato, posticè sursum acclinato; dorsali utrinque convexiusculo, anticè declivi, posticè valdè declivi; extremitate anticâ rotundatâ; latere postico brevi, cuneiformi; flexurâ costâque umbonali subinconspicuis; dentium lateralium antico subapproximato, postico subobsoleto.*

Subovate, tolerably strong, very inequilateral, moderately convex, glossy, white both within and without, with a yellowish anterior indistinct ray at the umbones, and very distinct and numerous oblique striæ, which in both of the valves reach even to the but little prominent umbonal ridge; ventral edge subarcuated in front, and rising posteriorly; dorsal edges moderately convex, sloping, the hinder one particularly so; front extremity rounded; posterior side short and wedge-shaped; front lateral tooth distinct and rather close, hinder almost obsolete.

Carolina to Georgia, United States.

Very closely resembling *T. similis* from which it may be distinguished by the much greater slope of its dorsal edges, its pointed posterior termination, the greater extension of its oblique striæ, and the presence of a second (though indistinct) lateral tooth. Mr. Say adds, "it has *generally* a rosaceous disk and one or two posterior (*i. e.* anterior) rays."

122. *TELLINA IRIDESCENS* (pl. lviii. f. 88.); Sanguinolaria, Benson, Annals Nat. Hist. 1842. vol. ix.

*T. testâ oblongâ, solidiusculâ, nitidâ, subæquilaterali, convexiusculâ, lævigatâ, albido-aurantiâ (intus aurantiâ), epidermide tenuissimâ subiridescente indutâ; margine ventrali subrecto; dorsali anticè subrecto paululumque declivi, posticè vix convexo et valdè declivi; extremitate anticâ rotundatâ; latere postico, vix brevior, cuneiformi; ligamento haud prominente; flexurâ costâque umbonali obsoletis; dente laterali parvo, approximato.*

Oblong, moderately strong, tolerably convex, subequilateral, smooth, glossy, pale orange, covered by a very thin slightly irri-

descent epidermis; ventral edge nearly straight; anterior side slightly the longer, rounded at its extremity, its dorsal edge scarcely convex and strongly sloping; ligament not prominent; fold and flexure obsolete; inside orange, a single small approximate lateral tooth.

Chusan, China.

I have received the species from Dr. Philippi with the proposed name of *T. carnea*, but know not whether he has subsequently published it.

- 123. *TELLINA FABULA* (pl. lvii. f. 62.) *Gmelin*, 3239; *Donov. Brit. Shells. vol. iii. t. 97*; *Wood, Gen. Conch. p. 156. t. 45. f. 4*; *Lam. no. 24*; *Philip. En. Mol. Sic. p. 26*; *Hanl. Des. Cat. p. 64. t. 3. f. 23.*

*T. testâ ovato-oblongâ, tenui, compressâ, nitidâ, extus intusque candidâ, subæquilateralî, anticè rotundatâ, posticè cuneiformi; valvulâ alterâ lævigatâ; margine ventrali subrecto, posticè sursum acclinato; dorsali anticè subdeclivi et convexo, posticè declivi et juxta ligamentum prominens retuso, deinde magis minusve convexo; flexurâ obsoletâ; dente laterali approximato, distincto.*

Ovate-oblong, thin, compressed, glossy, white within and without, nearly equilateral, rounded in front, wedge-shaped behind; one valve smooth, the other with crowded oblique striæ; ventral edge nearly straight, and rising posteriorly; front dorsal margin moderately sloping and convex, posterior one declining and near the prominent ligament retuse, but afterwards more or less convex; lateral tooth approximate, distinct.

Britain, Europe.

*T. fragillissima* of Chemnitz, vol. vi. f. 101, (*T. vitrea* of *Gmelin*), is usually considered as a bad representation of this species.

- 124. *TELLINA TENUIS* (pl. lviii. f. 81, 82.) *Da Costa*, *Brit. Conch. p. 210*. *Wood, Gen. Conch. p. 155. t. 44. f. 3, 4*; *Mont. Tes. Brit. 1. p. 59*; *Lam. no. 25*. *Hanl. Des. Cat. p. 64. t. 3. f. 22*; *Philip. En. Mol. Sic. 1. p. 27*; *T. balaustina, Dillw. D. Cat. p. 93.*

*T. testâ ovali, tenuiusculâ, compressâ, nitidissimâ, subæquilateralî, intus extusque albâ, aurantiâ aurantio-rubrâ aut rosâ, lævigatâ; margine ventrali anticè convexo, posticè sursum acclinato; dorsali anticè convexo subdeclivi, posticè declivi et juxta ligamentum prominens retuso; latere antico paulo longiore, rotundato; antico obtusè angulato; flexurâ costaque umbonali subinconspicuis; dente laterali minimo, approximato.*

Var. (*T. exigua*, *Poli*. Test. Sicil. l. t. 15. f. 15, 17.) testâ abbreviatâ et idcirco margine dorsali utrinque magis declivi; flexurâ omnino obsoletâ.

Oval, thinnish, compressed, extremely glossy, subequilateral, smooth both within and without, of an orange, white, orange-red, yellow or rose colour; the ventral edge convex in front, straightish and ascending behind; anterior side rather the longer, rounded, its dorsal edge moderately sloping; posterior extremity obtusely angulated; ligamental edge sloping, at first retuse, then rather convex; ligament prominent; fold and flexure not quite obsolete; lateral tooth very small, approximate.

Var. Shortened, the dorsal edges consequently more sloping; the flexure quite obsolete.

Britain, Europe.

I agree with Dr. Philippi, in regarding the *T. hyalina* of Deshayes (Exped. Morea, t. 18. f. 12, 13, 14.), as a narrow variety of this very variable shell.

125. *TELLINA JUVENILIS* (pl. lvii. f. 63.) *Hanley*, Zool. Proc. 1844.

*T. testâ* ovatâ, subtrigonâ, tenui, pellucidâ. compressiusculâ, nitidâ, rubro-aurantiâ, lævigatâ, inæquilaterali; margine ventrali convexo aut convexiusculo; dorsali anticè subrecto et declivi, posticè convexo et valdè declivi; latere antico longiore, subattenuato, rotundato; postico brevi, et obtusè subcuneiformi; flexurâ costâque umbonali subobsoletis; ligamento satis conspicuo; dente laterali parvo.

Subtriangularly ovate, inequilateral, thin, pellucid, slightly compressed, smooth, glossy, orange-red; ventral margin more or less convex; anterior side the narrower, rounded at the extremity, its dorsal edge nearly straight; posterior extremity short, and obtusely wedge-shaped, the ligamental edge being convex and abruptly sloping; ligament distinct; fold and flexure almost entirely obsolete; lateral tooth small.

Philippines.

126. *TELLINA LUX* (pl. lvii. f. 71.) *Hanley*, Zool. Proc. 1844.

*T. testâ* subovali, tenui, pellucidâ, compressâ, nitidâ, aurantiâ, sublævigatâ, inæquilaterali; margine ventrali convexiusculo; dorsali utrinque subdeclivi, anticè convexiusculo, posticè brevi et incurvato; latere postico brevior, obtusissimè biangulato; extremitate anticâ obtusè rotundatâ; flexurâ costâque umbonali obsoletis; ligamento prominente; dente laterali approximato, distincto.

Somewhat oval, thin, pellucid, compressed, inequilateral, nearly smooth, and of an uniform glossy orange-flesh colour; the ventral margin moderately convex; the dorsal but little sloping on either side, but moderately convex in front, incurved and short behind; anterior extremity obtusely rounded; the posterior side much the shorter, and very obtusely biangulated; fold and flexure obsolete; ligament prominent; lateral tooth distinct and approximate.

Philippines.

127. *TELLINA VERNALIS* (pl. lviii. f. 84.) *Hanley*, Zool. Proc. 1844.

T. testâ subovali, tenuissimâ, compressiusculâ, pellucidâ, nitidissimâ, lævi, valdè inæquilaterali, albido-rosâ, utrinque rotundatâ; margine ventrali convexiusculo, anticè sursum acclinato; dorsali utrinque paululum convexiusculo, posticè declivi, anticè vix minimè declivi; latere antico producto, postico brevi; flexurâ obsoletâ; dente laterali minimo, subapproximato.

Somewhat oval, extremely thin and pellucid, slightly compressed, very inequilateral, uniform glossy pink, rounded at both extremities; margins slightly convex, the ventral rising in front, the anterior dorsal scarcely sloping; flexure obsolete; anterior side short and cuneiform; lateral tooth minute and nearly approximate.

Singapore.

128. *TELLINA INSCULPTA* (pl. lx. f. 136.) *Hanley*, Zool. Proc. 1844.

T. testâ oblongo-elongatâ, solidiusculâ, compressâ, æquilaterali, extus intusque candidâ; sulcis confertis concentricè exaratâ, striisque tenuissimis radiantibus (præsertim posticè) decussatâ; ultra costam umbonalem subobsoletam subsquamosâ; margine ventrali elongato, subrecto; dorsali utrinque subrecto, subdeclivi; extremitate posticâ subbiangulatâ; flexurâ ventrali distinctâ; dente laterali quamplurimum approximato.

Elongated-oblong, tolerably solid, compressed, equilateral, externally and internally white, with concentric crowded sulci, and most minute posterior radiating striulæ; beyond the almost obsolete umbonal ridge, the surface roughened by small scales or interrupted delicate lamellæ; ventral edge nearly straight, and much elongated; dorsal margins nearly straight and but moderately sloping; hinder extremity somewhat biangulated; ventral flexure distinct; lateral tooth extremely approximate.

Chiriqui, W. Columbia.

129. *TELLINA PSAMMOTELLA* (pl. lx. f. 134.) *Lamarck*, no. 30.  
*Hanl. Des. Cat.* p. 65. sup. t. 14. f. 6.

T. testâ oblongo-ovali, subtenui, convexiusculâ, nitidâ, extus intusque pallidâ, lævi, inæquilaterali; margine ventrali convexiusculo; dorsali utrinque subdeclivi, subrecto; latere antico producto, subattenuato, rotundato; extremitate posticâ obtusè biangulatâ; ligamento satis prominente; flexurâ obsoletâ; dente laterali distincto, subapproximato.

Oblong-oval, inequilateral, rather thin, moderately convex, glossy, smooth, both within and without of a very pale dull orange yellow; ventral margin moderately convex; dorsal edges but moderately sloping, and nearly straight; anterior side produced, rounded, and rather attenuated; ligament moderately prominent; flexure obsolete; lateral tooth distinct, approximated.

Hab. (?) Mus. Cuming and Walton.

130. *TELLINA (TELLINIDES) EMARGINATA* (pl. lvii. f. 70.) *Sow.*  
*Tankerv. Cat. App.* p. 3; *Hanl. Des. Cat.* p. 74. sup. t. 13, f. 55.

T. testâ oblongâ, tenui, subinæquivalvi, convexiusculâ, nitidâ, pallidâ, radio brevi rubro-aurantio utrinque ad umbones notatâ, lævigatâ, valdè inæquilaterali; margine ventrali subrecto, anticè sursum acclinato; dorsali anticè recto et subdeclivi, posticè convexo et valdè declivi; latere antico producto, rotundato, attenuato; extremitate posticâ brevi, obtusâ, emarginatâ; flexurâ obsoletâ; dente laterali distincto, approximato.

Oblong, rather inequivalve, tolerably convex, glossy, very pale orange, with a short deeper tinged ray on either side of the beaks, smooth, subpellucid, very inequilateral, produced rounded and attenuated in front, obtuse short and emarginated behind; ventral and anterior dorsal edges nearly straight, the latter not much sloping; posterior dorsal much sloping and convex beyond the ligament; flexure obsolete; lateral tooth distinct, approximate.

Singapore.

The Oblonga of Gmelin, (p. 2314. from Chemnitz, vol. 6. f. 87.) may possibly prove to be this or the preceding, but the resemblance is not sufficiently exact for identification, the figure bearing a greater resemblance to Cumana, or Aurora, but the description of the teeth by no means coinciding with the hinge of either.

131. *TELLINA (TELLINIDES) TRUNCATULA* (pl. lvii. f. 69.)  
*Sowerby*, *Tankerv. Cat. App.* p. 3; *Hanl. Des. Cat.* p. 73.

T. testâ T. Vestali simillimâ, sed fulvo-aurantiâ, valdè compressâ, et magis inæquilaterali; extremitate posticâ emarginatâ; margine dorsali postico subretuso, hand autem excavato.

Very like Vestalis, but more compressed and inequilateral, of a tawny orange, with the hinder extremity clearly notched, and the posterior dorsal edge not at all excavated, but simply and slightly retuse.

Ilo Ilo, Isle of Panay, Philippines (Cuming.)

132. TELLINA VESTALIS (pl. lvii. f. 67.) *Hanley*, Zool. Proc. 1844.

T. testâ oblongo-angustâ, tenuissimâ, convexiusculâ, nitidâ, lævi, intus extusque niveâ, inæquilaterali; margine ventrali subrecto, paululum convexiusculo, anticè sursum acclinato; dorsali anticè minimè declivi et paululum convexiusculo, posticè prope ligamentum excavato, deinde convexo et subdeclivi; extremitate lateris antici longioris rotundatâ; extremitate posticâ subemarginatâ, subatenuatâ, obtusè biangulatâ; flexurâ obsoletâ; dente laterali approximate.

Elongated-oblong, extremely thin, slightly convex, inequilateral, smooth, glossy, snowy white both within and without; ventral margin but very slightly convex, and somewhat rising in front; anterior side rather the longer, its extremity rounded, its dorsal edge scarcely convex and hardly at all sloping; hinder dorsal edge excavated at the ligament, afterwards convex and slightly sloping; posterior extremity rather attenuated, slightly emarginated, and obtusely biangulated; flexure obsolete; the lateral tooth approximate.

Isles of Negros and Luzon, Philippines.

133. TELLINA LANCEOLATA (pl. lviii. f. 86, 87.) *Chemnitz*, vol. vi. p. 111. f. 103; Wood, Gen. Conch. p. 167. t. 45. f. 2; Hanl. Des. Cat. p. 67. t. 4. f. 40; Psammotæa pellucida, *Lam.* no. 4; Deles. t. 5. f. 10.

T. testâ oblongo-elongatâ, tenui, compressâ, nitidiusculâ, æquilaterali, roseâ aut albido-roseâ, concentricè et confertissimè striatâ, anticè rotundatâ, posticè lanceolatâ; margine ventrali subrecto; dorsali utrinque convexiusculo, anticè subhorizontali, posticè subdeclivi; costâ umbonali conspicuâ; ligamento satis prominente; costellâ internâ ex umbonibus posticè radiante; dente laterali distincto, subapproximato.

Elongated oblong, equilateral, thin, compressed, rather glossy,

rose-coloured or white with a rosy tinge; concentrically striated, the striæ extremely close; ventral edge nearly straight; dorsal but moderately convex on either side, nearly horizontal in front, and not greatly sloping behind; anterior extremity rounded; posterior lanceolate; ligament tolerably prominent; fold conspicuous; an internal posterior radiating riblet; the lateral tooth distinct, subapproximate.

Isle of Negros, Philippines (Cuming).

134. *TELLINA HYALINA* (pl. lxi. f. 167.) *Gmelin*, 3235; Dillw. Des. Cat. p. 85; Hanl. Des. Cat. p. 67. t. 4. f. 39; *T. complanata pellucida*, *Chem.* vol. vi. f. 99.

*T. testâ oblongâ, inæquivalvi, tenuissimâ, compressâ, valdè inæquilaterali, nitidâ, albidâ, subpellucidâ, lævigatâ; margine ventrali convexo et posticè acclivi; dorsali posticè subrecto vixque declivi, anticè arcuato et declivi; latere antico perbrevis, rotundato; postico producto, rotundato, attenuato; natibus inconspicuis; ligamento infosso; flexurâ obsoletâ; costellis duabus internis posticè radiantibus; dente laterali quamplurimum approximato.*

Oblong, inequivalve, very thin, much compressed, very inequilateral, of a brilliant smooth semi-transparent white; the ventral margin convex and rising behind; the hinder area of the dorsal edge, which is but slightly convex and not greatly sloping posteriorly, and sloping and arcuated anteriorly, flattened; anterior side short and rounded; posterior side produced, rounded, and attenuated; ligament sunken; beaks not prominent; fold obsolete; two internal rib-like lines radiating from the hinge posteriorwards; the lateral tooth almost touching the primary ones.

The only perfect specimen I have ever beheld of this extremely rare shell, belongs to Dr. Bates, who most courteously forwarded it from the United States, for my inspection.

135. *TELLINA (TELLINIDES) TIMORENSIS* (pl. lxi. f. 158, 172) *Lamarck*, *Anim. sans vert.* vol. v. = *Blainvilliei* *Sest.*

*T. testâ oblongâ, tenuiusculâ, subæquilaterali, compressiusculâ, nitidâ, subpellucidâ, extus intusque candidâ (nonnunquam marginibus roseo tinctis) concentricè substriatâ, anticè rotundatâ, posticè attenuatâ et supernè obtusè angulatâ; margine ventrali subrecto; dorsali utrinque subrecto et subdeclivi, anticè paululum convexiusculo, posticè subretuso; flexurâ costâque umbonali obsoletis; ligamento magno, prominente; dente laterali, approximato.*

Oblong, rather thin and semi-transparent, nearly equilateral, rather compressed, both within and without of a glossy white



which is often tinged with yellow and rarely with rose-colour at the margins, concentrically substriated; the ventral edge nearly straight, and usually very indistinctly inflated behind; anterior side slightly the shorter, rounded at the extremity; posterior side rather attenuated, and obtusely angulated above; dorsal edges scarcely sloping and nearly straight, the front being very slightly convex, the hinder somewhat retuse; ligament large and prominent; umbonal ridge and fold quite obsolete; teeth distinct and generally tinged with rose-colour, the lateral one approximate.

Isle of Negros, Philippines; &c.

136. *TELLINA CONSPICUA* (pl. lviii. f. 100.) *Hanley*.

T. testâ *T. Timorensis* simillimâ, sed abbreviatâ, et ovali-subtrigônâ; margine dorsali utrinque magis declivi, posticè convexiusculo.

Very like *Timorensis*, but not so elongated; the dorsal edges much more sloping, and the front one not retuse.

This may eventually prove but a young variety of *Sinuata*, but the only specimen I have ever seen (purchased with this name attached, and now in the collection of Mr. Cuming) has been pronounced distinct by every Conchologist who has inspected it.

137. *TELLINA SINUATA* (pl. lxiii. f. 203.) *Spengler*, loc. cit.; *Tellinides Timorensis*, *Sow. Gen.*; *Reeve, Conch. Syst.* t. 56. f. 2; *Young, T. nivea, Wood, Gen. Conch.* p. 177. t. 46. f. 1; *Dillwyn, D. Cat.* p. 89; *Index Test.* t. 4. f. 92.

T. testâ subovali, solidiusculâ, inæquilaterali, compressiusculâ, nitidâ, extus intusque albidâ, concentricè striolatâ, striisque obliquis anticis infernè decussatâ; margine ventrali subrecto, posticè sinuoso; dorsali anticè brevi subrecto declivi, posticè incurvato vixque declivi; extremitate posticâ latâ, rotundatâ; anticâ obtusâ; costâ umbonali obsoletâ; ligamento magno, prominente; dente laterali approximato.

Oval, rather oblique, tolerably strong, and somewhat compressed, glossy, whitish, with concentric striulæ which are decussated in front by fine raised oblique and rather distant ones; ventral edge sinuated posteriorly, and somewhat swollen in the middle; anterior side clearly the shorter, its extremity obtuse, its dorsal edge nearly straight and moderately sloping; posterior extremity obtuse, rounded below, but forming a rounded-off angle with the short concave and scarcely sloping dorsal margin; ligament large and prominent; umbonal ridge obsolete; inside white, teeth as in the last, but *never* tinged with rose-colour.

Closely allied to *Timorensis*, but differs in shape, sinuosity, and the decided inequality of its sides. The anterior oblique striæ, which in some specimens extend along the ventral edge posteriorward, are rarely present in that species.

Tranquebar; Sumatra.

138. *TELLINA TRIANGULARIS* (pl. lx. f. 150.) *Chemnitz*, vol. vi. p. 96. f. 85; *Dillwyn*, D. Cat. p. 79; *Desh.* ed. *Lam.* vol. vi. p. 209; *Tellinides triangularis*, *Hanl.* Des. Cat. p. 74. t. 4. f. 33; *Tellinides politus* *Sow.*, *Tanker.* Cat. App. p. 4; *T. trilatera*, *Gmel.*; *T. obliqua*, *Lam.* no. 47.

*T. testâ* ovato-trigonâ, inæquilaterali, solidiusculâ, compressâ, nitidissimâ, sub epidermide flavescente-candidâ, anticè et prope marginem ventralem convexum concentricè striatâ, alterâque in valvulâ striis flexuosis obliquis posticè ornatâ; margine dorsali anticè brevi et arcuato, posticè subrecto et producto; extremitate posticâ obtusè acuminatâ; anticâ rotundatâ; ligamento prominente; flexurâ nullâ; dente laterali approximatisimo.

Ovate-triangular, tolerably solid, compressed, inequilateral, both internally and externally white, the epidermis yellowish and very glossy; with concentric striæ in front, and towards the convex ventral margin which rises a little behind; anterior side rounded, its dorsal edge convex; posterior side produced and obtusely acuminated, the dorsal edge nearly straight and very oblique; ligament prominent; flexure obsolete; the lateral tooth quite close to the beaks.

Moluccas? Mus. Cuming, &c.

139. *TELLINA (TELLINIDES) ACUMINATA* (pl. lx. f. 141.) *Hanley*, Des. Cat. p. 73. sup. t. 13. f. 53.

*T. testâ* elongatâ, subovatâ, solidiusculâ, valdè inæquilaterali et inæquivalvi, compressâ, politâ, lævigatâ, extus intusque albidâ aut albido-aurantiâ; margine ventrali subarcuato; dorsali anticè subarcuato satisque declivi, posticè elongato subrecto et declivi; latere antico breviorè, rotundato; postico valdè producto, acuminato, ad extremitatem rotundato; ligamento infozzo; flexurâ costâque umbonali obsoletis; dente laterali approximato.

Elongated, subovate, moderately strong, very inequilateral and inequivalve, both valves compressed, but one much shallower than the other; smooth, polished, both within and without of a pale orange becoming white towards the subarcuated ventral edge; anterior side the shorter, rounded, its dorsal edge arched and moderately sloping; posterior side much produced, acuminated,

with the tip rounded, the dorsal edge nearly straight, moderately sloping, and very long; ligament sunken; umbonal ridge and flexure obsolete; lateral tooth approximate.

Hab.? Mus. Britannicum.

140. *TELLINA PLANISSIMA* (pl. lix. f. 124. and pl. lxii. f. 197.)  
*Anton*, Verzn. Conch. p. 4; *Ab. und Bes. Conch.* part 5. t. 5. f. 2;  
*Tellinides rosea*, *Sow.* Gen. Shells, f. 1; *Reeve*, Conch. Syst. t. 56.  
f. 1.

T. testâ ovatâ, tenuiusculâ, inæqualvi, inæquilaterali, compressâ, extus intusque roseâ, concentricè substriatâ; margine ventrali convexiusculo; dorsali utrinque declivi, anticè convexo, posticè subrecto; latere antico rotundato; postico longiore, rotundato-cuneiformi; ligamento prominente; flexurâ subobsoletâ; costellâ internâ in valvulâ complanatâ posticè radiante; dente laterali parvo, approximato.

Var. Candida.

Ovate, rather thin, inequivalve, inequilateral, compressed, rose-coloured both within and without, concentrically substriated; ventral margin moderately convex, the flexure not quite obsolete; dorsal edges sloping, the anterior convex, the posterior nearly straight; anterior extremity rounded; posterior side the longer and of a rounded wedge-shape; ligament prominent; an internal rib radiating posteriorward in the flattened valve; lateral tooth small, approximate.

Var. Quite white.

Moluccas.

The *Opalina* of Chemnitz is closely allied to, if not identical with, the white variety of this shell.

141. *TELLINA* (*TELLINIDES*) *PURPURASCENS* (pl. lxii. f. 194.)  
*Broderip* and *Sowerby*, Zool. Jour. vol. iv. p. 363; *Habl. Des. Cat.*  
p. 74. sup. t. 9. f. 18; *Zool. Beechey's Voy.* t. 42. f. 2.

T. testâ oblongo-ovali, tenui, compressâ, nitidiusculâ, subæquilaterali, roseo-purpurascente, pellucidâ, striis radiantibus et concentricis confertissimè decussatâ; margine ventrali convexiusculo; dorsali utrinque paululum declivi, anticè convexiusculo, posticè subrecto; latere antico minimè longiore, rotundato; extremitate posticâ subangulatâ, subemarginatâ; ligamento prominente; flexurâ omnino obsoletâ; dente laterali approximato.

Oblong-oval, thin, compressed, rather glossy, nearly equilateral, purplish-rose coloured, pellucid, decussated by most crowded radiating and concentric striæ; ventral margin moderately convex;

dorsal but little sloping on either side, slightly convex in front, nearly straight behind; anterior side very slightly the longer, rounded; posterior extremity subangulated, subemarginated; ligament prominent; flexure quite obsolete; lateral tooth approximate.

Real Leijos, Central America (Cuming).

142. *TELLINA* (*TELLINIDES*) *OVALIS* (pl. lx. f. 147.) *Sowerby*, Tank. Cat. App. p. 3; *Hanl. Des. Cat.* p. 73. sup. t. 13. f. 56; *T. tridentata?* *Anton*, in *Wieg. Arch.* 1837.

*T. testâ ovali, tenuissimâ, pellucidâ, subæquilaterali, compressâ, nitidâ, roseâ, radiis angustis albidis frequentibus ornatâ, concentricè substriatâ; margine ventrali convexiusculo; dorsali utrinque paululum declivi; latere antico longiore, rotundato; extremitate posticâ obtusâ, aliquantulum latiore; flexurâ costâque umbonali obsoletis; natibus inconspicuis; ligamento prominente; dente laterali approximato, distincto.*

Oval, pellucid, subequilateral, compressed, extremely thin, concentrically striated, and variegated with crowded narrow whitish rays on a rose-coloured ground; the ventral margin slightly convex; the dorsal but little sloping on either side, the ligament edge slightly retuse; posterior extremity obtusely rounded, and rather broader than the rounded termination of the more elongated anterior side; fold and flexure obsolete; beaks not prominent; ligament projecting; a distinct approximate lateral tooth.

Moluccas; San Nicolas, Zebu; (Cuming).

The *Solen striatus*, of Gmelin (from Chemnitz, vol. vi. f. 57, 58.), seems closely allied if not identical.

143. *TELLINA COCCINEA* (pl. lviii. f. 90.) *Chemnitz*, vol. vi. p. 118. f. 109; *Dillw. Des. Cat.* p. 87; *Tellinides coccinea. Hanl. Des. Cat.* p. 74. t. 4. f. 42.

*T. testâ ovali, tenuissimâ, compressâ, æquilaterali, pellucidâ, roseo-purpurascete, utrinque rotundatâ; concentricè striatâ, striis anticis frequentibus, posticis remotis et subimbricatis; margine ventrali convexiusculo; dorsali utrinque paululum declivi; flexurâ costâque umbonali obsoletis; ligamento prominente; costellis duabus internis (in junioribus) posticè radiantibus; dentibus primariis subobsoletis, dente laterali approximato, distincto.*

Oval, compressed, very thin, equilateral, of an uniform pellucid purple-rose colour, with concentric striæ which are crowded in front, but distant and subimbricated behind; the ventral edge moderately convex; the dorsal but little sloping on either side of

the little prominent beaks; extremities rounded and nearly equally broad; the fold and flexure obsolete; the ligament prominent; two rather indistinct internal rib-like lines radiating (in the young only) posteriorward; cardinal teeth nearly obsolete, the approximate anterior lateral very distinct (when young).

San Nicolas, Zebu (Cuming).

144. *TELLINA RHODON* (pl. lviii. f. 93.) *Hanley*, Zool. Proc. 1844.

T. testâ oblongâ, tenuissimâ, compressiusculâ, nitidissimâ, lævi, roseâ, pellucidâ, valdè inæquilaterali, utrinque rotundatâ; margine ventrali convexiusculo, anticè sursum acclivi; dorsali anticè convexiusculo et vix declivi, posticè subdeclivi; latere postico brevi subattenuato; natibus inconspicuis; flexurâ costâque umbonali obsolete; dente laterali approximato.

Oblong, very inequilateral, extremely thin, rather compressed, smooth, polished, and of an uniform purplish rose-colour; the ventral margin slightly convex, and rising in front; the dorsal but slightly convex on either side, scarcely sloping in front, but moderately so behind; the extremities rounded, and the shorter posterior side somewhat attenuated; beaks not prominent; the fold and flexure obsolete; lateral tooth approximate.

Hab. ? Mus. Cuming.

145. *TELLINA OUDARDI* (pl. lxvi. f. 262.) *Payraud*. Cat. Corse, p. 40. t. 1. f. 16, 17, 18; Desh. ed. Lam. vol. vi. p. 210; Hanl. Des. Cat. p. 65. sup. t. 14. f. 8.

"T. testâ ovatâ, compressâ, nitidâ, pellucidâ, obliquè cancellatâ, lineis transversis albis parvulisque longitudinalibus rubris ornatâ; anticè et posticè radiis luteo-rubrescentibus; intus et extus rubrâ.

Ovate, compressed, glossy, transparent, obliquely cancellated, adorned with transverse white lines and small longitudinal red ones; rays of a yellowish red both in front and behind; red both within and without."

I have been obliged to content myself with reprinting the original description, and giving a representation of the typical shell deposited in the Museum at Paris, as I am unable to procure a specimen in England for the purpose of comparison. Had it not been for the mention of sculpture on the surface of one of its valves, I should have doubted its distinctiveness as a species. "The valves are furnished with transverse striæ in lozenges, accompanied by a multitude of silvery-white little rays both which are crossed by very delicate longitudinal lines of a deeper red than the

ground-colour of the shell." There seems from the drawing to be but a single anterior moderately distant lateral tooth, and no *external* appearance of rays in the front of the shell. The sculpture must indeed be minute, to have escaped the attention (as it has) of our very accurate artist.

146. *TELLINA SUBROSEA* (pl. lvii. f. 64.) *Hanley*.

T. testâ oblongâ, tenuissimâ, compressâ, valdè inæquilaterali, rosâ, pellucidâ, lævi, nitidâ, posticè breviorè et obtusè cuneiformi, anticè obtusâ aut rotundatâ; margine ventrali subrecto; dorsali anticè paululum declivi, posticè declivi, utrinque subrecto aut convexiusculo; flexurâ inconspicuâ; dente laterali subobsoleto, approximato.

Oblong, extremely thin, compressed, smooth, pellucid, rosy, very inequilateral and highly glossed; ventral margin straightish; dorsal edges straightish or but slightly convex, very slightly sloping in front, moderately so behind; anterior extremity rounded or obtuse; posterior side the shorter and obtusely wedge-shaped; flexure very inconspicuous; lateral tooth approximate and almost obsolete.

Catbalonga, Isle of Samar, soft mud; (Cuming).

The specimens closely resemble the young of the *Punicea* group, but both locality and dentition are adverse to this conclusion. They appear, however to be scarcely adult.

147. *TELLINA MARGARITINA* (pl. lxvi. f. 257.) *Lam.* no. 18; *Deless. Rec.* t. 6. f. 4.

T. testâ ovali, tenui, pellucidâ, nitidâ, margaritaceâ; latere antico (postico) attenuato.

Oval, thin, transparent, glossy, nacreous white, front (hinder) side attenuated.

New Holland.

The representation of the species in *Delessert's* magnificent folio, enables me to refer this shell to the present division. I am unable, however, to add a more lengthened description than the very meagre outline furnished in the 'Animaux sans vertèbres'. The only specimens I have ever seen were the original types of the French Museum; but they sufficed to convince me of the clear distinctness of the species from any possessing a similar dentition.

\* \* \* Devoid of lateral teeth.

148. *TELLINA Cumana* (pl. lviii. f. 73.) *Psammobia Cumana*,

*Costa Cat. Test. Sicil.* 20. t. 2. f. 7.; *Tellina Costæ Philip. En. Mol. Sic.* 1. p. 28. t. 3. f. 11.

T. testâ ovatâ aut ovali-oblongâ, tenuiusculâ, inæquilaterali, valdè convexâ, subnitidâ, lævigatâ, extus intusque albidâ, umbonibus hyalinis et plerumque pallidè aurantiis; margine ventrali satis convexo; dorsali utrinque subrecto, posticè satis, anticè paululum declivi; latere antico longiore ad extremitatem rotundato; postico angustato, obtusè subtruncato; natibus acutis, et rectè incurvatis; costâ umbonali satis distinctâ, haud autem conspicuâ; ligamento infosso; dentibus parvis.

Varying from ovate to oval-oblong, rather thin, inequilateral, not very glossy, smooth, both within and without whitish, with the umbones usually hyaline and often tinged with orange; ventral edge moderately convex, but not arcuated; dorsal edges nearly straight, sloping behind, and but moderately so in front; anterior side the longer and rounded at its termination; posterior side narrowed, and obtusely subtruncated; beaks acute and not inclining to either side; ligament sunken, umbonal ridge not prominent but tolerably evident; teeth small.

Very closely resembling *Plebeia*, but proportionally more ventricose, with the ventral edge less arcuated, &c.

Mediterranean.

149. *TELLINA PLEBEIA* (pl. lix. f. 129, and f. 151.) *Hanley*, *Zool. Proc.* 1844.; *Psammobia lævigata* (?) *Lamarck*.

T. testâ subovatâ, solidiusculâ, convexâ, subnitidâ, lævigatâ, intus extusque, albidâ, umbonibus hyalinis et rubro-aurantiis; margine ventrali convexo aut subarcuato; dorsali anticè prope nates recto et paululum declivi, posticè recto satisque declivi; latere antico longiore, rotundato; extremitate posticâ obtusè angulatâ; ligamento infosso; costâ umbonali conspicuâ; flexurâ obsoletâ. Var. *Ovatâ*; castâ umbonali minus conspicuâ; umbonibus sæpe albidis.

Subovate, tolerably strong, convex, not very glossy, smooth, both within and without whitish, with the umbones hyaline and orange-red; ventral edge convex or subarcuated; anterior side the longer, rounded, its dorsal edge straight and scarcely sloping near the beaks; posterior extremity obtusely angulated, the ligamental edge straight and sloping; ligament sunken; flexure obsolete.

Var. (from Africa) ovate, the ventral edge more regularly arcuated, the fold less prominent, and the umbones often colourless.

Real Leijos, Central America; Senegal.

150. *TELLINA UMBONELLA* (pl. lvi. f. 13.) *Lamarck*, no. 48. *Hanl. Des. cat.* p. 70.

T. testâ obovali, solidiusculâ, subinæquilaterali, convexâ aut convexusculâ, subimpolitâ, extus intusque albidâ, umbonibus hyalinis et roseo tinctis, epidermide tenui fugaci cinerascete indutâ, lævigatâ; margine ventrali convexo aut subarcuato; dorsali utrinque subrecto et paulo declivi; latere antico paulo longiore, rotundato; postico subangulato; natibus haud prominentibus; ligamento infosso; costâ umbonali obsoletâ; dentibus minimis.

Oboval, tolerably strong, slightly inequilateral, merely convex, not glossy, both within and without whitish under a thin ashy epidermis, with the beaks somewhat hyaline and tinged with rose-colour, smooth or with a few antiquated lines; ventral edge decidedly convex; the dorsal edges but little sloping, and nearly straight; anterior side rather the longer, rounded at its extremity; posterior side subangulated; ligament sunken; beaks not prominent; umbonal ridge obsolete; teeth minute.

Very like *Plebeia* in all but the shape, which is much more elongated in that species.

Port Lincoln, Australia.

151. *TELLINA NOBILIS* (pl. lx. f. 143.) *Hanley, Zool. Proc.* 1844.

T. testâ ovali, solidiusculâ, convexâ, inæquilaterali, nitidissimâ, intus extusque roseâ, lævigatâ; margine ventrali convexusculo medio plerumque subrecto; dorsali anticè convexusculo et vix declivi, posticè subdeclivi et subrecto aut convexusculo; latere antico longiore, ad extremitatem obtusè rotundato; postico obtusè angulato; natibus obtusis; flexurâ costaque umbonali subobsoletis; ligamento prominulo; dentibus parvis.

Oval, tolerably strong, convex, inequilateral, very glossy, smooth, rosy within and without; ventral edge moderately convex; dorsal tolerably convex and scarcely sloping in front, moderately sloping and varying from straightness to slight convexity behind; anterior side the longer, obtusely rounded at its extremity; posterior side obtusely angulated; fold and flexure nearly obsolete; ligament rather prominent; beaks obtuse; teeth small.

Luzon, Philippines.

152. *TELLINA LUCERNA* (pl. lviii. f. 98.) *Hanley, Zool. Proc.* 1844.

T. testâ oblongâ, solidiusculâ, subventricosâ, nitidiusculâ, lævigatâ, albidâ, umbonibus aurantiis, anticè longiore et rotundatâ, posticè obtusè cuneiformi, margine ventrali subrecto; dorsali pos-



ticè declivi, anticè paululum declivi, utrinque subrecto; flexurâ obsoletâ; ligamento subinfosso; disco interno aurantio.

Oblong, tolerably solid, rather ventricose, slightly glossy, smooth, whitish, with the umbones orange, anteriorly longer and rounded, posteriorly obtusely wedge-shaped; ventral and dorsal edges straightish, the latter posteriorly sloping, anteriorly scarcely sloping; flexure obsolete; ligament rather sunken; inner disc orange.

Isles of Negros, Mindanao, and Panay.

153. *TELLINA AURORA* (pl. lviii. f. 76.) *Hanley*, Zool. Proc. 1844. p. 147.

T. testâ T. Psammotellæ simillimâ, convexiore autem, et umbonibus rubro-aurantiis; ligamento infosso; dentibus lateralibus nullis.

Very like *T. Psammotella* but more convex, and with the umbones of an orange-red. The ligament too is sunken and the hinge devoid of lateral teeth.

Panama.

154. *TELLINA GALATÆA* (pl. lxiv. f. 223.) *Psammobia Galatæa* *Lam.* no. 18; *Hanl. Des. Cat.* p. 58. sup. t. 13. f. 44. *T. Sericina* *Jonas*, Ab. und. Bes. Conch. part 3. t. 5. f. 1.

T. testâ oblongâ, solidiusculâ, convexâ, valdè inæquilaterali, subnitidâ, extus intusque candidâ, punctis minutissimis elevatis confertis radiatim asperatâ; margine ventrali subrecto; dorsali anticè subhorizontali, posticè brevi recto et declivi; latere antico producto, rotundato; extremitate posticâ truncato-cuneiformi; ligamento infosso; flexurâ obsoletâ; dentibus parvis.

Var. (*T. Diana*, Zool. Proc. 1844.) subovatâ, punctis obsoletis; margine dorsali utrinque magis declivi; extremitate posticâ magis obtusâ.

Oblong, tolerably strong, convex, very inequilateral, rather glossy, white within and without, roughened by crowded radiating rows of minute raised dots; ventral edge nearly straight; anterior side produced, rounded at the extremity; dorsal edge convex and nearly horizontal; posterior side of a truncated wedge-shape; the ligamental edge straight, short, and much sloping; ligament sunken; flexure obsolete; teeth very small.

Var. Subovate, the dots obsolete; dorsal edges more sloping; posterior end more obtuse.

China.

155. *TELLINA CANDIDA* (pl. lxii. f. 195.) *Psammotæa candida* *Lam.* no. 6. *Hanl. Des. Cat.* p. 60.

T. testâ elongato-oblongâ, subæquivalvi, maximè inæquilaterali, extus intusque candidâ, lævigatâ, convexâ aut subventricosâ; margine ventrali posticè convexiusculo, anticè arcuato et sursum acclinato; dorsali anticè convexiusculo paululumque declivi, posticè retuso et valdè declivi, extremitate anticâ rotundato-attenuatâ; latere postico brevissimo, obtusè cuneiformi; ligamento subinfosso; costâ umbonali subobsoletâ; dentibus minimis.

Elongated-oblong, subequivalve, extremely inequilateral, white both within and without, convex or subventricose, smooth, with however obsolete radiating lines, and very indistinct striulæ towards the ventral margin, which latter is straightish behind, but curves upward in front, so as somewhat to attenuate the rounded anterior extremity; front dorsal edge scarcely at all sloping, rather convex; posterior side very short and obtusely wedge-shaped, the dorsal edge retuse and decidedly sloping; ligament rather sunken; fold nearly obsolete; teeth minute.

China (?) Mus. Metcalfe.

Very closely allied to *Galatæa*, but narrower, more convex, and utterly devoid of those radiating lines of raised dots, which are so characteristic of that species. The posterior extremity is more regularly cuneiform, and the anterior termination more attenuated.

156. *TELLINA ELONGATA* (pl. lxii. f. 199.) *Hanley, Zool. Proc.* 1844.

T. testâ elongatâ, angustâ, tenuiusculâ, satis convexâ, extus intusque candidâ, lævigatâ, valdè inæquilaterali; margine ventrali medio subretuso, anticè obliquè arcuato; dorsali anticè subhorizontali et subrecto, posticè subdeclivi et subretuso; extremitate lateris antici longioris obtusâ posticâ acuminato-truncatâ; ligamento infosso; flexurâ obsoletâ; dentibus parvis.

Elongated, narrow, thinnish, moderately convex, smooth, white, very inequilateral; ventral edge rather retuse in the middle, and obliquely arcuated anteriorly; anterior side the longer, its extremity obtuse, its dorsal edge almost horizontal and nearly straight; hinder extremity with a truncated acumination, the ligamental edge slightly incurved and moderately sloping; flexure obsolete; ligament sunken; inside white; teeth small.

Chiriqui, West Columbia.

157. *TELLINA ASSIMILIS* (pl. lviii. f. 95.) *Hanley, Zool. Proc.* 1844.

T. testâ *T. elongatæ* simillimâ, convexiore autem, et latere postico obtusè cuneiformi paululumque subrostrato.

Closely resembling a young *Elongata*, but more convex, and the hinder side obtusely wedge-shaped and slightly beaked; the hinder extremity not in the slightest degree truncated, but the ligamental edge sweeping to the lower margin in a concave and then convex curve.

Luzon, Philippines.

158. *TELLINA LILIUM* (pl. lviii. f. 85.) *Hanley*, *Zool. Proc.* 1844.

T. testâ ovato-oblongâ, tenuiusculâ, convexâ aut subventricosâ, extus intusque albâ; concentricè substriatâ, striis supra costam umbonalem elevatis; margine ventrali subrecto anticè sursum acclinato; dorsali anticè subrecto paululumque declivi, posticè valdè declivi et prope ligamentum prominulum recto deindeque convexo; extremitate lateris antici producti infernè et obliquè rotundatâ; extremitate posticâ brevi, cuneiformi.

Ovate-oblong, rather thin, convex or subventricose, white both within and without, with concentric striulæ, and obsolete radiating lines, which latter are most perceptible in the younger specimens, the striulæ elevated upon and beyond the tolerably distinct fold; anterior side produced, its dorsal edge nearly straight, and scarcely sloping, its extremity obliquely rounded near the straight ventral margin; posterior side short and wedge-shaped; the ligamental edge straight, but convex at the extremity; ligament small, and moderately prominent.

Isles of Burias and Negros, Philippines.

159. *TELLINA TENTA* (pl. lxvi. f. 266.) *Say*, *Am. Conch.* t. 65. f. 3; *Gould*, *Invert. Massach.* p. 68. f. 43.

T. testâ oblongâ, tenuissimâ, subflexuosâ, convexiusculâ, extus intusque albidâ, lævigatâ, valdè inæquilaterali; margine ventrali subrecto; dorsali anticè convexo (aut subarcuato) et paululum declivi, posticè retuso et subdeclivi; latere postico brevi, ad extremitatem angustato et paululum subrostrato; extremitate anticâ rotundatâ; costâ umbonali subinconspicuâ; ligamento parvo, prominente; superficie internâ lineis radiantibus ornatâ; dentibus minimis.

Oblong, very thin, rather flexuous, moderately convex, whitish, very inequilateral, very nearly if not quite smooth, ventral margin straightish; front dorsal edge convex or subarcuated, and very little sloping; hinder one retuse and moderately sloping; posterior side short, narrowed and very slightly beaked at its termination, the tip truncated or subemarginated; anterior end rounded; fold

not very conspicuous; ligament small, projecting; inside whitish, and distinctly marked with radiating lineoles; teeth very small.

Boston.

160. *TELLINA ANCILLA* (pl. lviii. f. 94.) *Hanley*, Zool. Proc. 1844.

T. testâ oblongo-elongatâ, tenuiusculâ, satis convexâ, nitidissimâ, extus intusque candidâ, concentricè substriatâ, lineisque obsoletis radiantibus ornatâ; striis supra costam umbonalem subobsoletam remotioribus, distinctis, subimbricatis; margine ventrali subrecto; dorsali anticè subdeclivi et convexiusculo, posticè subrecto et declivi; extremitate lateris antici producti rotundatâ; latere postico brevior, obtusè cuneiformi.

Elongated-oblong, thinnish, moderately convex, very glossy, pure white both within and without, with concentric striulæ, which are most perceptible at the nearly straight ventral margin; and become subimbricated distinct and rather distant upon and beyond the almost obsolete fold; anterior side produced, its extremity rounded, its dorsal edge slightly convex and not much sloping; posterior side obtusely wedge-shaped, ligamental edge straightish, tip rounded.

Lord Hood's Island.

161. *TELLINA BODEGENSIS* (pl. lix. f. 102.) *Hinds*, Zool. Sulphur, vol. ii. t. 21. f. 2. p. 67.

T. testâ oblongo-elongatâ, inæquilaterali, solidiusculâ, compressâ, nitidâ, extus intusque candidâ, concentricè et subimbricatum sulcatâ; margine ventrali subrecto, anticè sursum acclinato; dorsali utrinque subrecto, anticè vix minimè declivi; latere postico brevior, cuneiformi; extremitate anticâ rotundatâ; flexurâ obsoletâ; ligamento satis prominente; dentibus lateralibus nullis.

Elongated-oblong, inequilateral, rather solid, compressed, of a glossy ivory white, with regular moderately distant concentric subimbricated sulci, (which in the only specimen I have ever seen disappear in one valve at the posterior end); the ventral margin nearly straight, rising in front, the anterior dorsal scarcely at all sloping; posterior side the shorter, wedge-shaped; front extremity rather obtusely rounded; fold and flexure obsolete; ligament moderately projecting; inside white; no lateral teeth.

Russian Bodejas.

162. *TELLINA RHOMBOIDES* (pl. lviii. f. 92, 96, 97.) *Quoy* and *Gaim.*, Voy. Ast. Zool. vol. iii. p. 502. t. 81. f. 4-7; T. clathrata,

*Desh.* Lam. vol. vi. p. 208; *Hanl. Des. Cat.* p. 65. sup. t. 14. f. 12.

T. testâ ovali-oblongâ, solidiusculâ, valdè inæquilaterali, magis minusve convexâ, subnitidâ, albidâ, pallidè luteâ, aurantiâ aut albido-roseâ, striis anticè concentricis et medio obliquis confertim ornatâ; margine ventrali subrecto aut paululum convexo; dorsali anticè subrecto et paululum declivi, posticè satis declivi; extremitate lateris antici producti obtusè rotundatâ; extremitate posticâ magis minusve obtusè cuneiformi; ligamento prominente; flexurâ vix perspicuâ.

Var. Pallidè luteâ, radiis angustis roseis; striis obliquis majoribus et fortioribus.

Oval-oblong, very inequilateral, tolerably strong, more or less convex, rather glossy, whitish pink, pale yellow, or orange, with very delicate striæ, which are concentric in front but become oblique in the middle; ventral edge straightish, or not much convex; anterior side produced, its extremity rounded, its dorsal edge straightish and but little sloping; posterior side rather variable in shape, but short, and more or less obtusely wedge-shaped; ligament prominent; flexure not quite obsolete.

Var. Pale yellow with narrow rosy rays, the oblique striæ larger and stronger.

Isles of Zebu and Ticao, Philippines, (Cuming).

163. *TELLINA FORMOSA* (pl. lviii. f. 83.) *Hanley, Zool. Proc.* 1844.

T. testâ obovatâ, solidiusculâ, nitidâ, convexiusculâ, subpellucidâ, albidâ, radiis interruptis roseis striisque minutis obliquis confertis undique ornatâ; margine ventrali convexo; dorsali utrinque convexiusculo, anticè subdeclivi, posticè valdè declivi; extremitate lateris antici producti rotundatâ, lateris postici brevissimi obtusissimè angulatâ; natibus roseis; ligamento parvo et satis prominente; flexurâ subobsoletâ.

Obovate, moderately strong, glossy, scarcely convex, somewhat pellucid, white, with numerous narrow rosy interrupted rays, and crowded minute oblique striæ; ventral margin convex; dorsal moderately convex on both sides, little sloping in front but greatly so behind; anterior side produced, its extremity rounded; posterior side short, its extremity very obtusely angulated; beaks rosy; ligament small, moderately prominent; flexure nearly obsolete.

Daleguete, Zebu.

164. *TELLINA DISPAR* (pl. lix. f. 108, 113, 114.) *Conrad*, Jour. Ac. N. S. Philad. vol. vii.

T. testâ ovatâ, solidiusculâ, satis convexâ, nitidâ, candidâ aut roseâ, subæquilaterali, posticè brevior et obtusè angulatâ aut obtusissimè cuneiformi, anticè rotundatâ; alterâ valvulâ lævi, alterâ striis tenuissimis anticè concentricis deinde obliquis confertim ornatâ; margine ventrali convexiusculo; dorsali utrinque haud multum declivi aut convexo; ligamento prominente; flexurâ obsoletâ.

Ovate, tolerably strong, moderately convex, glossy, white or rose-coloured, nearly equilateral, rounded in front, rather shorter and very obtusely wedge-shaped or obtusely angulated behind; one valve smooth, the other with fine striæ, which are concentric in front and then become oblique and crowded; ventral margin but slightly convex; dorsal edges not greatly sloping or convex; ligament prominent; flexure obsolete.

Sandwich Isles; and Isle of Bohol, soft mud, eight fathoms, (Cuming).

In some specimens there is a tinge of rose-colour near the dorsal edges.

165. *TELLINA LUTEA* (pl. lix. f. 103, and lxxv. f. 249.) *Gray* (1828) in Index Test. Sup. t. 1. f. 3; Hanl. Des. Cat. p. 67. sup. t. 1. f. 3. T. Guildfordiæ, *Gray* in Grif. Cuvier, Mol. t. 19. f. 2.

Young; T. alternidentata, *Brod. and Sow.*, Zool. Jour. vol. iv. p. 363; Zool. to Beechey Voy. t. 44. f. 5.

T. testâ ovali-subtrigonâ, subæquilaterali, subinæquali, compressiusculâ, sublævigatâ, albidâ aut pallidè roseâ, epidermide subviridi fugaci (in junioribus) indutâ, anticè rotundatâ; margine ventrali subrecto aut convexiusculo; dorsali anticè subdeclivi et convexiusculo, posticè declivi et prope ligamentum magnum subretuso deindeque convexo; latere postico obtusissimè cuneiformi; natibus erosis; flexurâ obsoletâ; superficie internâ albido-roseâ, utrinque saturatius fucatâ, aut roseo subradiatâ; dentibus primariis in utrâque valvulâ duobus, altero simplici, altero bicuspidato, alternatim inter se insertis.

Oval-subtriangular, subequilateral, rather compressed, slightly inequivalve, nearly smooth, whitish or pale rose-colour under (in young specimens only) a thin fugacious greenish epidermis; ventral edge but slightly (if at all) convex; anterior side rounded, its dorsal edge but slightly convex and not much sloping; posterior extremity very obtusely wedge-shaped, the dorsal edge near the

great ligament retuse but afterwards convex ; beaks eroded ; flexure obsolete ; inside pink with a darker ray or stain on either side ; primary teeth two in each valve, a simple and a bicuspidate, which are alternately inserted.

Icy Cape.

166. *TELLINA COLUMBIENSIS* (pl. lxxv. f. 246.) *Hanley*, Zool. Proc 1844. p. 71.

T. testâ ellipticâ aut oblongo-ellipticâ, compressiusculâ, subtenui, lævigatâ, extus intusque albidâ, epidermide tenuissimâ fulvocinereâ indutâ ; margine ventrali medio convexiusculo, utrinque arcuato ; dorsali utrinque convexo, anticè paulo, posticè satis declivi ; latere antico longiore, rotundato ; extremitate posticâ acuminatâ ; flexurâ costâque umbonali subobsoletis ; ligamento pro-minulo ; dentibus minimis.

Elliptic or elongated-elliptic, fragile, subinequivalve, rather compressed, whitish, covered by a very thin opalescent pale drab epidermis, quite smooth ; ventral edge arcuated, ascending posteriorly ; anterior side much the longer, its extremity rounded and somewhat attenuated, its dorsal edge convex and scarcely sloping ; posterior end acuminated, hinder dorsal edge at first hollowed out for the slightly prominent ligament, then very convex and considerably sloping ; flexure and umbonal ridge all but obsolete ; inside white, teeth minute.

Monte Christi, West Columbia.

Its more compressed valves and minute teeth will distinguish it from the few species which are allied to it in outline. The hinge-margin is very short and rather broad.

167. *TELLINA RUFESCENS*<sup>†</sup> (pl. lxxiii. f. 213.) *Chemnitz*, vol. vi. f. 97. (malè) ; *Dillw. D. Cat.* p. 85 ; T. operculata, *Gmel.* 3235 ; *Wood, Gen. Conch.* p. 165. t. 42. f. 1 ; *Lam. no. 13* ; *Sow. Gen. Shells* ; *Hanl. Des. Cat.* p. 62. t. 4. f. 37 ; *Knorr. vol. vi. t. 12. f. 1.*

T. testâ oblongâ, inæquivalvi, inæquilaterali, solidâ, subnitidâ, rubro-roseâ (intus saturatius fucatâ), sublævigatâ, lineis obsoletis radiantibus plerumque notatâ ; alterâ valvulâ convexâ, alterâ complanatâ ; margine ventrali medio subrecto, utrinque (præsertim posticè) sursum acclinato ; dorsali antico convexo, et vix declivi ; dorsali postico subdeclivi, prope ligamentum magnum prominensque retuso, deinde magis minusve convexo ; latere antico paululum brevior, rotundato ; postico attenuato obtusissimè acuminato ; costâ umbonali latâ, obtusâ ; flexurâ subobsoletâ.

Oblong, inequivalve, rather strong, scarcely glossy, within and

*T. rufescens Chemnitz vol. vi. f. 97.*

without of a lighter or darker reddish rose-colour, nearly smooth, often with obsolete radiating lines, covered when fresh with a very thin epidermis; one valve convex, the other flat; ventral edge nearly straight or subretuse in the middle, rising at the extremities and particularly posteriorly; anterior side rather the shorter, its extremity rounded, its dorsal edge convex but scarcely sloping; posterior side very obtusely acuminate, the dorsal edge at first retuse and little sloping, then moderately sloping and convex; fold broad and obtuse; flexure nearly obsolete; ligament large.

Tumbez, Peru; soft sandy mud, five fathoms (Cuming).

168. *TELLINA NITIDA* (pl. lix. f. 101.) *Poli*, Test. Sic. vol. i. t. 15. f. 2, 3, 4; Lam. no. 28; Philip. En. Moll. Sic. vol. i. p. 27; Hanl. Des. Cat. p. 64. sup. t. 14. f. 4.

T. testâ oblongâ, valdè compressâ, æquilaterali, satis solidâ, nitidâ, pallidè carneâ aut fulvâ; concentricè striatâ, striis tenuissimis, confertis, sed alterâ in valvulâ posticè magis remotis alterâque in valvulâ posticè obsolete; margine ventrali medio subrecto; dorsali anticè convexo et subdeclivi, posticè declivi et subrecto; latere antico rotundato, postico cuneiformi; flexurâ nullâ.

Oblong, much compressed, equilateral or nearly so, tolerably strong, very glossy, pale flesh-coloured or fulvous, within of a deeper hue, with fine regular and close concentric striæ, which at the posterior end of one valve are quite obsolete, and in the other are *there* replaced by stronger, more distant, and subimbricated ones; ventral edge nearly straight in the middle; anterior side rounded, its dorsal edge convex, and but moderately sloping; hinder side wedge-shaped, ligamental edge nearly straight; no flexure.

Mediterranean.

169. *TELLINA SOULEYETI* (pl. lix. f. 111.) *Hanley*, Zool. Proc. 1844.

T. testâ oblongâ, tenuiusculâ, convexo-depressâ, extus intusque albidâ, lævigatâ; margine ventrali convexiusculo; dorsali anticè convexiusculo et subdeclivi, posticè recto aut subretuso et valdè declivi; flexurâ costâque umbonali distinctis; latere antico longiore, rotundato; extremitate posticâ subrostratâ; ligamento infosso; natibus acutis.

Oblong, tolerably convex, nearly equilateral, but rather longer in front, of a rather dull white, not very thick, and quite smooth; the ventral edge more or less convex; the flexure as well as the umbonal ridge quite distinct; dorsal edge moderately convex and scarcely sloping in front, nearly straight or slightly retuse behind; anterior



extremity rounded, posterior somewhat beaked ; beaks acute ; ligament sunken ; inside white.

San. Nicolas, Zebu.

In the young, there is a slight appearance of rudimentary lateral teeth.

170. *TELLINA MICANS* (pl. lix. f. 106.) *Hanley*, Zool. Proc. 1844.

T. testâ ovato-oblongâ, tenui, compressiusculâ, subinæquilaterali, nitidissimâ, pellucidâ, nivêâ ; striis paucis remotis subobsoletis concentricis, etiamque in adultis, striis subobliquis remotis, anticè ornatâ ; margine ventrali anticè obliquè arcuato, posticè subrecto ; dorsali anticè vix declivi et convexusculo, posticè declivi subretuso et angulum obtusum cum margine postico subincurvato formante ; latere postico breviorè, cuneiformi ; extremitate anticâ obtusè rotundatâ ; flexurâ obsoletâ.

Ovate-oblong, thin, rather compressed, slightly inequilateral, extremely glossy, pellucid, snow-white, with a few indistinct and rather distant concentric striæ in front, which are crossed in adult specimens by very slightly oblique and distant ones ; ventral edge obliquely rounded anteriorly, and straightish behind ; anterior side slightly the longer, obtusely rounded, its dorsal edge scarcely sloping ; hinder extremity wedge-shaped, the ligamental edge subretuse near the moderately projecting ligament, and forming a very obtuse angle with the slightly incurved posterior edge ; flexure obsolete.

Catbalonga, Isle of Samar.

Bias, Isle of Negros.

171. *TELLINA MILES* (pl. lxii. f. 188.) *Hanley*, Zool. Proc. 1844.

T. testâ T. cuspidi affini sed oblongâ, rostratâ, compressiore ; margine antico dorsali vix paululum declivi ; ventrali convexo ; valvulâ alterâ lineis elevatis concentricis undique striatâ.

Like *T. cuspidi* but oblong, beaked, and more compressed ; front dorsal edge scarcely at all sloping ; ventral edge convex ; one valve nearly smooth, the other with regular fine elevated concentric striæ.

Hab. ? (Mus. Metcalfe.)

172. *TELLINA CUSPIS* (pl. lix. f. 125.) *Hanley*, Zool. Proc. 1844.

T. testâ ovatâ, solidiusculâ, convexâ, nitidiusculâ, rosçâ, anticè et infernè substriatâ ; margine ventrali arcuato ; dorsali utrinque subdeclivi, anticè convexo, posticè recto aut subretuso ; flexurâ costâque umbonali distinctis ; latere antico paululum longiore, rotundato ; postico subacuminato, subrostrato ; ligamento infosso.

Ovate, moderately strong, convex, rather glossy, deep rose-colour, substriated in front and towards the lower margin; ventral edge arcuated; dorsal but slightly sloping on either side, convex in front, straight or subretuse behind; fold and flexure perceptible; anterior side slightly the longer, rounded; posterior subacuminated, and somewhat beaked; ligament sunken.

Hab. ? (Mus. Cuming, Metcalfe.)

173. *TELLINA SCALPELLUM* (pl. lix. f. 116.) *Hanley, Zool. Proc.* 1844.

T. testâ oblongâ, tenuissimâ, compressâ, nitidissimâ, valdè in-æquilaterali, saturatius rosçâ, pellucidâ, sublævigatâ; margine ventrali recto aut subrecto; dorsali antico paululum declivi, subrecto; latere postico brevi, obtusè subcuneiformi; extremitate lateris antici longioris, rotundatâ; ligamento parvo, prominulo; flexurâ obsoletâ.

Oblong, very thin and pellucid, very inequilateral compressed, nearly smooth or concentrically substriated, of a very glossy deep rose-colour; ventral edge more or less straight; front side produced, rounded at its extremity, its dorsal edge nearly straight and very little sloping; hinder side obtusely wedge-shaped; ligament small and prominent; flexure almost entirely obsolete.

Isle of Zebu.

174. *TELLINA UNDULATA* (pl. lix. f. 107.) *Hanley, Zool. Proc.* 1844.

T. testâ oblongâ, tenuissimâ, compressâ, impolitâ et valdè in-æquilaterali, intus extusque albidâ, subobliquè et concentricè undulatâ; margine ventrali convexo; dorsali anticè subrecto et vix declivi, posticè incurvato satsique declivi; latere postico brevi, attenuato, rostrato, ad extremitatem biangulato; flexurâ costâque umbonali distinctis; natibus acutis; ligamento infosso.

Oblong, very thin, compressed, very inequilateral, whitish both within and without, not glossy, with oblique undulations which become concentric near the short attenuated and beaked posterior extremity, of which the tip is acute and biangulated; ventral edge convex; dorsal anteriorly straightish and scarcely sloping, posteriorly incurved and moderately sloping; anterior side produced, its extremity obtusely rounded; fold and flexure perceptible; ligament sunken; beaks acute.

St. Elena, West Columbia.

175. *TELLINA CYGNUS* (pl. lix. f. 122.) *Hanley, Zool. Proc.* 1844.

T. testâ ovatâ aut ovato-oblongâ, solidiusculâ, subæquilaterali, convexâ, nitidâ, candidâ, concentricè substriatâ; margine ventrali convexusculo; dorsali anticè subrecto et paululum declivi, posticè recto et valdè declivi; extremitate anticâ rotundato-obtusâ; latere postico cuneiformi, subrostrato; flexurâ ventrali distinctâ; ligamento infosso; superficie internâ candidâ, submargaritaceâ.

Ovate or ovate-oblong, tolerably firm, nearly equilateral, convex, glossy, white, concentrically substriated; ventral margin slightly convex; dorsal nearly straight and very slightly sloping in front; quite straight and much sloping behind; anterior end obtusely rounded; posterior side wedge-shaped, and subrostrated; flexure distinct; ligament sunken; outside white and somewhat pearly.

Bias, Isle of Negros, Philippines.

176. *TELLINA ALA* (pl. lx. f. 137, 146.) *Hanley, Zool. Proc.* 1844.

T. testâ ovatâ, solidiusculâ, subinæquivalvi, subæquilaterali, nitidâ, convexusculâ, extus intusque albidâ, radio brevi pallidè aurantio in adultis utrinque ornatâ, concentricè substriatâ; margine ventrali magis minusque convexo; dorsali anticè convexo et subdeclivi, posticè declivi et prope nates subretuso; latere antico rotundato, longiore; postico angulato, subrostrato; flexurâ costâque umbonali conspicuis; ligamento infosso; cardine dentibus primariis parvis, et nonnunquam dente laterali antico rudimentali, instructo.

Var. ovato-trigonâ, solidâ, lævi aut sublævigatâ, convexâ, nequam rostratâ; flexurâ costâque umbonali subinconspicuis.

Ovate, tolerably solid, very slightly inequivalve, nearly equilateral, rather convex, of a silky gloss, concentrically substriated, whitish within and without, with (in adult specimens) a short pale orange streak radiating on either side from the beaks; ventral edge more or less convex; anterior side rather the longer, rounded at its extremity, its dorsal edge convex, and not greatly sloping; posterior side angulated and subrostrated, its dorsal edge retuse near the beaks, and decidedly sloping; fold and flexure conspicuous; ligament rather sunken; primary teeth small, sometimes a rudimentary anterior lateral one.

Var. Ovate-triangular, solid, convex, nearly or quite smooth, not at all beaked, the orange rays clearly visible; fold and flexure by no means conspicuous.

Ceylon.

177. *TELLINA NYMPHALIS* (pl. lxii. f. 196.) *Lam. no.* 50; *Hanl. Des. Cat.* p. 70.

*T. testâ ovatâ aut obovatâ, magis minusve ventricosâ, solidiusculâ, subnitidâ, sublævigatâ, subæquilaterali, extus intusque albidâ, lineis obsoletis pallidioribus subradiatâ, epidermide pertenui lutescente indutâ; margine ventrali anticè arcuato, posticè sursum acclinato; dorsali anticè prope nates acutas subrecto deindeque arcuato et subdeclivi, posticè subretuso et valdè declivi; latere antico longiore, rotundato; postico attenuato, angulato, et paululum subrostrato; flexurâ costâque umbonali satis conspicuis; ligamento prominulo; dentibus minimis.*

Ovate or obovate, more or less ventricose, not very strong, rather glossy, nearly equilateral, whitish both within and without, covered by a very thin yellowish ash-coloured epidermis, nearly smooth being merely wrinkled by the lines of growth, peculiarly marked with obsolete radiating lines; ventral edge arcuated in front and rising behind; anterior side rather the longer, its extremity rounded, its dorsal edge at first nearly straight and then arched and moderately sloping; posterior side attenuated, angulated, and very slightly beaked; fold and flexure tolerably distinct; ligament short and slightly prominent; teeth very small.

Gambia.

In typical examples there is a slight groove-like depression behind the fold, which does not exist in the next species.

178 *TELLINA CAYENNENSIS* (pl. lxii. f. 190.) *Psammobia C. Lam.* no. 11; *Hanl. Des. Cat.* p. 58. sup. t. 11. f. 48; *Deles. Rec.* t. 5. f. 7; *T. constricta, Phil., Ab. und Bes. Conch.* part 3. t. 5. f. 5; *T. intastriata, Say, Jour. Ac. Nat. Sc. Philad.*

*T. testâ ovali-trigonâ, solidâ aut solidiusculâ, magis minusve ventricosâ, subinæquivalvi, subinæquilaterali, nitidiusculâ, extus intusque albidâ, epidermide cinerascete indutâ, sublævigatâ; margine ventrali convexiusculo; dorsali anticè convexo et subdeclivi, posticè subrecto et declivi; latere antico paulo longiore, rotundato; postico cuneiformi; natibus rectè incurvatis; ligamento prominulo, satis magno; flexurâ costâque umbonali satis conspicuis; dentibus haud parvis.*

Triangularly-oval, more or less solid and ventricose, slightly inequivalve and inequilateral, scarcely glossy, nearly smooth, whitish within and without, covered by a very thin ashy epidermis; ventral edge slightly convex; dorsal convex and moderately sloping in front, straightish (never incurved) and strongly sloping behind; front side rounded, rather the longer; hinder side wedge-shaped; beaks incurved; ligament little prominent, and as well as the teeth, of a moderate size; fold and flexure tolerably distinct.

Closely resembling the last, of which it is possibly a variety, but is more triangular, and not so distinctly marked with radiating lines. Guiana, Mexico, Florida.

179. *TELLINA ALBA* (pl. lxii. f. 193.) *Quoy and Gaim.*, Voy. Astrol. Zool. vol. iii. p. 500. t. 81. f. 1, 2, 3.

T. testâ subovatâ, solidiusculâ, subcompressâ, subpellucidâ, nitidiusculâ, extus intusque albâ, sublævigatâ; margine ventrali convexo et posticè sursum acclinato; dorsali anticè arcuato et vix subdeclivi, posticè elongato declivi et subrecto; latere antico lato, multum brevior, rotundato; postico attenuato, obtusè cuneiformi; ligamento infosso; costâ umbonali et flexurâ ventrali inconspicuis; natibus subacutis; dentibus bifidis, subvalidis, simplicibus subobsoletis.

Subovate, tolerably strong, rather flat, semi-transparent, nearly smooth, somewhat glossy, white, one of the valves with obsolete concentric striæ; the ventral edge convex and ascending posteriorly; anterior side broad, and distinctly the shorter, rounded at its extremity, arcuated and but little sloping at its dorsal margin; the posterior side obtusely cuneiform, its tip attenuated and obtusely angulated, its dorsal edge elongated scarcely convex and obliquely sloping; beaks rather prominent; ligament sunken; umbonal ridge *not* conspicuous; the bifid teeth rather strong, the simple nearly rudimentary.

In all the specimens I have seen, there are two or three obsolete radiating folds at the anterior extremity.

Australia.

180. *TELLINA PROXIMA* (pl. lxvi. f. 264. and lix. f. 115.) *Brown* in *Zoology to Beechey*, p. 154. t. 44. f. 4; *Phil. Ab. und Bes. Conch.* part 9. pl. 5. f. 4; T. sordida, *Couthouy*, *Boston Journ. Nat. Hist.* vol. ii. p. 59. t. 3. f. 11; *Sanguinolaria sordida*, *Gould*, *Invert. Massachusetts*, p. 67; *Macroma tenera*, *Leach*, *Appendix Ross. Voyage*.

T. testâ obovali, tenuiusculâ, valdè inæquilaterali, convexâ aut convexiusculâ, impolitâ, extus albidâ, intus candidâ, epidermide cinerascens fugaci indutâ, concentricè obsoletè et subimbricatim striolatâ; margine ventrali magis minusve convexo; dorsali anticè arcuato et paulo declivi, posticè recto et valdè declivi; latere postico brevi, et infernè subangulato; extremitate posticâ rotundatâ; natibus plerumque erosis; ligamento magno, prominente; costâ umbonali subobsoletâ; dentium altero in utrâque valvulâ bifurcato.

Oboval, thinnish, fragile, very inequilateral, more or less con-

vex, dull whitish (internally pure white), covered by a thin ash-coloured fugacious epidermis; ventral margin more or less convex; front dorsal edge arched and slightly sloping, hinder one straight and greatly sloping; posterior side short, subangulated below; anterior extremity rounded; beaks usually eroded; ligament large and prominent; fold almost obsolete; one of the teeth bifurcate (when the hinge is perfect, which is rarely the case) in each valve. Boston, (Gould); Arctic Ocean.

181. *TELLINA CALCAREA* (pl. lxii. f. 183.) *Chemnitz*, vol. vi. p. 140. f. 136; *Dillwyn D. Cat.* p. 93; *Hanl. Des. Cat.* p. 46. t. 4. f. 43; *T. sordida*, *Phil. Ab. und Bes. Conch.* part 9. t. 5. f. 6.

*T. testâ* ovatâ aut subovatâ, solidiusculâ, subinæquivalvi, convexiusculâ, inæquilaterali, extus intusque albidâ, epidermide cinerascente indutâ, sublævigatâ; margine ventrali satis convexo, posticè acclinato; dorsali anticè convexo et subdeclivi, posticè declivi et subrecto; latere antico longiore, ad extremitatem rotundato; postico obtusè cuneiformi; natibus acutis, prominentibus; costâ umbonali satis distinctâ; flexurâ ventrali subinconspicuâ; ligamento magno, prominulo; dentibus parvis.

More or less ovate, moderately firm, very slightly inequivalve, tolerably convex, inequilateral, whitish both within and without, never polished but often slightly glossed, covered with an ashy epidermis, under which it is nearly smooth or merely marked with the lines of growth; ventral edge ascending behind, moderately convex; anterior side the shorter, rounded at its extremity, its dorsal edge convex and somewhat sloping, but near the acute and prominent beaks slightly incurved; hinder side obtusely wedge-shaped, the ligamental edge nearly straight; umbonal ridge tolerably evident; the ventral sinus almost obsolete; ligament large and moderately projecting.

Iceland and Greenland.

182. *TELLINA NASUTA* (pl. lxiv. f. 224.) *Conrad*, *Jour. Ac. Nat. Sc. Philad.* vol. vii. p. 258; *Hanl. Des. Cat.* p. 71. sup. t. 14. f. 7.

*T. testâ* ovato-oblongâ, solidâ, inæquilaterali, convexâ, impolitâ, sublævigatâ, albidâ, epidermide cinerascente plerumque indutâ; margine ventrali medio subretuso; dorsali anticè convexo paululumque declivi, posticè elongato recto et declivi; latere antico brevior, rotundato; postico cuneiformi; extremitate posticâ subbiangulatâ; natibus obtusis; lunulâ nullâ; flexurâ costâque um-

bonali conspicuis; ligamento magno et satis prominente; superficie internâ candidâ; dentibus satis magnis.

Ovate-oblong, solid, inequilateral, convex, dull, whitish, usually covered by an ashy epidermis, nearly smooth, flexuous; ventral margin convex at the sides, but retuse in the middle; anterior side rounded, its dorsal edge convex and but little sloping; posterior side wedge-shaped, its extremity nearly biangulated, its dorsal margin elongated sloping and nearly straight; fold and flexure conspicuous; ligament tolerably large and prominent; beaks obtuse; no lunule; inside white, the teeth of a moderate size.

California.

— 183. *TELLINA EDENTULA* (pl. lxx. f. 243.) *Brod. and Sow.*, *Zool. Jour.* vol. iv. p. 363; *Zool. to Beech. Voy.* t. 41. f. 5. and t. 44. f. 6; *Hanl. Des. Cat.* p. 71.

T. testâ obovali-subtrigonâ, solidiusculâ, subventricosâ, subimpolitâ, subinæquilaterali, sordidè albidâ, sublævigatâ; margine ventrali convexiusculo; dorsali utrinque subdeclivi, anticè convexo, posticè subrecto; latere antico paulo longiore, rotundato; postico obtusè subangulato; flexurâ costâque umbonali subobsoletis; ligamento magno, prominente; dentibus minimis, sublamellosis.

Var. Abbreviatâ et magis inæquilaterali.

Oboval, subtriangular, tolerably strong, more or less ventricose, of a rather dull dirty white, nearly smooth, more or less inequilateral; ventral edge moderately convex; dorsal not greatly sloping on either side, convex in front, straightish and rather short behind; anterior side the longer, rounded at its extremity; posterior side obtusely subangulated; fold and flexure rather obsolete; ligament large and prominent; inside white; teeth very small, and usually sublamellar.

Var. Shorter and more inequilateral.

Behring's Straits.

— 184. *TELLINA INORNATA* (pl. lix. f. 123.) *Hanley*, *Zool. Proc.* 1844. p. 144.

T. testâ ovato-oblongâ, subtenui, subventricosâ, impolitâ, subæquilaterali, sordidè albidâ, epidermide tenui et cinereâ indutâ, lævigatâ; marginis ventralis parte mediâ rectâ aut subretusâ; dorsalis parte anticâ convexiusculâ et paululum declivi, parte posticâ subdeclivi; extremitate anticâ rotundatâ, posticâ attenuato-rotundatâ; flexurâ costâque umbonali obsoletis.

Ovate-oblong, rather thin, subventricose, and nearly equilateral,

smooth, dull dirty white under a thin ash-coloured epidermis; ventral edge straight or subretuse in the middle; dorsal slightly convex and very little sloping in front, moderately sloping behind; front extremity rounded, the hinder rounded but attenuated; fold and flexure obsolete.

Conception, Chili.

A fossil-like shell, which much resembles an elongated and swollen *Edentula*.

185. *TELLINA BALTHICA* (pl. lix. f. 121.) *Lin. Sys.* 1120; *Chem. vol. 6. f. 128. Hanl. Des. Cat. p. 72. t. 4. f. 85.*

*T. testâ suborbiculari, subtrigonâ, solidâ, æquilaterali, subventricosâ, albidâ aut pallidè carneâ, intus plerumque roseâ, lævigatâ; margine ventrali arcuato et posticè sursum acclinato; dorsali antico brevi, subdeclivi, et prope nates prominentes subrecto, deindeque convexo; margine ligamentali subrecto aut convexiusculo, et valdè declivi; latere antico rotundato; extremitate lateris postici angulati rotundatâ; ligamento magno, prominente; flexurâ costâque umbonali obsoletis; dentibus parvis.*

Suborbicular, slightly triangular, thick, equilateral, more or less ventricose, white or pale flesh coloured, with the interior usually rosy, smooth, not glossy; ventral edge arcuated, rising behind; front dorsal edge, short and not much sloping, nearly straight adjacent to the prominent beaks, becoming convex subsequently; hinder dorsal edge nearly straight or but slightly convex, much sloping; anterior side rounded; posterior side angulated, but with the tip rounded; ligament large, and prominent; fold and flexure obsolete; teeth small.

Baltic.

Easily distinguished from the allied species, by its more rounded shape, its ventricoseness, and arcuation of its lower edge. *T. Fabricii* which approaches it in these respects, is less angulated behind.

186. *TELLINA FUSCA* (pl. lix. f. 117.) *Psammobia fusca, Say, Jour. Ac. Nat. Sc. Philad. 5. p. 220. Sanguinolaria fusca Conrad. Am. Mar. Conc. p. 34. t. 7. f. 1; Gould, Inv. Massa. p. 66. f. 42.*

*T. testâ obovatâ, solidiusculâ aut subtenui, subæquilaterali, compressâ aut convexiusculâ, subimpolitâ, albidâ, epidermide tenuissimâ cinerascete indutâ, lævigatâ; margine ventrali maximè et regulariter arcuato; dorsali utrinque subdeclivi, anticè arcuato, posticè subrecto; latere antico rotundato; postico subangulato;*



natibus parvis acutis, prominentibus; flexurâ costâque umbonali obsoletis, ligamento prominulo; dentibus angustissimis.

Obovate, rather thin, nearly if not quite equilateral, more or less compressed or but slightly convex, scarcely at all glossy, whitish under a thin ashy epidermis, smooth; ventral edge regularly and greatly arcuated; the dorsal edges not much sloping, arched in front, and nearly straight behind; anterior side rounded, posterior subangulated; beaks small, acute, and prominent; ligament rather projecting; fold and flexure obsolete; teeth very narrow.

"Its varieties in size, strength, and colour, are very great. In muddy bays, they grow to a larger size, and are thick and of a blueish and, sometimes, a rusty colour, and covered with a firm dark epidermis. In clean sandy localities they are very thin, pure white, or sometimes of a delicate rose or lemon colour, and the epidermis very slight."—*Gould*.

North America.

187. *TELLINA INCONSPICUA*\* (pl. lix. f. 120.) *Broderip and Sow.*, Zool. Jour. vol. 4. p. 363; Zool. to Beech. Voy. t. 41. f. 6. *Hanl. Des. Cat.* p. 72.

T. testâ Solidulæ affini, sed impolitâ et epidermide fusco-cine-

\* A species which bears some resemblance to a young *Inconspicua*, or *Fusca*, but which nevertheless is probably distinct, at least judging from the figure, for I have never been able to procure a specimen of the shell, is the

*TELLINA PETALUM* (pl. lxvi. f. 263.) *VALENCIENNES* in Zool. to Humb. and Bonpl. Voy. vol. ii. p. 222. t. 48. f. 2.

T. testâ subovatâ, tenuiusculâ, fragili, subæquilaterali, compressiusculâ, sublævigatâ, supernè subrosçâ, infernè albido-luteâ, zonis subrosçis plerumque ornatâ; margine ventrali satis convexo; dorsali anticè convexo et subdeclivi, posticè convexiusculo et declivi; latere antico rotundato; postico subangulato; ligamento parvo, brevi; dentibus lateralibus nullis.

Subovate, thinnish, fragile, nearly inequilateral, rather compressed, nearly smooth, merely marked with the striæ of growth, somewhat rosy near the beaks, becoming pale yellowish with rosaceous zones towards the ventral edge, which is but moderately convex; front dorsal edge convex and not much sloping, hinder one but slightly convex and decidedly sloping; anterior extremity rounded; posterior side subangulated; ligament short and small; teeth small, no lateral ones.

Acapulco.

rascente indutâ; latere antico brevior; postico haud acuminato; ligamento permagno; natibus et umbonibus plerumque erosis.

Very like *Solidula*, but the front side is the shorter one, and the posterior extremity is not at all acuminated. The surface is not glossy, is usually eroded near the beaks, and covered with an ashy-brown epidermis near the ventral margin, which is regularly sub-arcuated. The ligament is very large and prominent.

Arctic Ocean.

188. *TELLINA FABRICII* (pl. lix. f. 112.) *Hanley*; *Venus fragilis*, *Otho Fabr.*, Faun. Groen. p. 413.

T. testâ suborbiculari, solidâ, æquilaterali, convexâ aut subventricosâ, impolitâ, extus intusque albidâ, epidermide cinerascens indutâ, concentricè subrugulosâ; margine ventrali magis minusve arcuato; dorsali utrinque convexiusculo, antici subdeclivi, posticè declivi; latere antico rotundato; postico vix etiam obtusissimè subangulato; natibus erosis; ligamento permagno, prominente; flexurâ obsoletâ; dentibus minimis.

Suborbicular, solid, equilateral, convex or subventricose, dull white, covered by an ashy epidermis, concentrically wrinkled by the lines of growth; ventral margin more or less arcuated; dorsal edges moderately convex, rather sloping in front, decidedly so behind; anterior side rounded, posterior scarcely at all angulated; beaks eroded; ligament large, and prominent; flexure obsolete; teeth very small.

There being already a *T. fragilis*, I am compelled to alter the original specific appellation of the shell.

Northern Seas.

189. *TELLINA SOLIDULA* (pl. lix. f. 109, 110.) *Pultney*, Cat. Dorset. p. 29; *Mont. Test. Brit.* p. 63; *Maton, Lin. Tr.* 8; *Wood, Gen. Con.* 193. t. 46. f. 2; *Lam. no.* 51; *Hanl. Des. Cat.* p. 70. t. 4. f. 84; *T. zonata, Dillw. D. Cat.* p. 100; *Psammobia solidula, Turton, Biv. Brit.* p. 95. t. 8. f. 2; *T. Balthica, Phil. En. Moll. Sic.* vol. i. p. 28; *T. rubra, Du Cost. Brit. Conch.* t. 12. f. 14.

T. testâ rotundato-ovatâ, solidâ, subæquilaterali, convexâ aut subventricosâ, magis minusve nitidâ, lævigatâ, albidâ, luteâ, aurantiâ aut rosâ; margine ventrali anticè arcuato, posticè sursum acclinato; dorsali anticè arcuato et subdeclivi, posticè declivi et vix convexiusculo; latere antico paululum longiore, rotundato; postico acuminato; natibus acutis, prominentibus; costâ umbonali obsoletâ; ligamento magno prominente; dentibus angustissimis et sæpe obsoletis.

Rounded-ovate, solid, subequilateral, convex or subventricose, more or less glossy, smooth or with a few ridges of growth, very variable in colouring, being white yellow orange or rosy; ventral edge arcuated in front, and rising behind so as to form a sharp angle with the slightly convex and moderately sloping ligamental margin; anterior side very slightly the longer and rounded; its dorsal edge arched and moderately sloping; ligament prominent; fold obsolete; beaks rather prominent and acute; teeth extremely narrow, and usually obsolete.

Britain; Europe; Senegal.

190. *TELLINA IRUS* (pl. lx. f. 145.) *Hanley*, Zool Proc. 1844.

T. testâ ovatâ aut obovatâ, crassâ (in adultis), subventricosâ, subæquilaterali, impolitâ, extus intusque sordidè albidâ; concentricè rugosâ, rugis interruptis, minutis, confertissimis, elevatis; margine ventrali magis minusve arcuato; dorsali anticè convexo et subdeclivi, posticè convexiusculo elongato declivi; latere antico paululum breviorè, rotundato; postico infernè angulato; lunulâ (in adultis) parvâ, profundâ; ligamento infosso; costâ umbonali rubobsoletâ; dentibus satis magnis.

Ovate or obovate, solid, coarse, subventricose, subequilateral, dull dirty white both within and without, the surface roughened by minute concentrically arranged but unconnected elevated wrinkles (remining one of those upon *Rugosa* but on a much smaller scale), which however are usually (except in portions) abraded; front dorsal edge convex after passing the small but distinct lunule; ligamental edge not greatly convex, and decidedly sloping so as to form an angle with the more or less arched ventral margin; beaks inclining to the rounded extremity of the shorter anterior side; ligament sunken but distinct; a large bifid and a simple scarcely rudimentary tooth in one valve, two rather strong diverging obtuse ones in the other.

Guinea? Mus. Cuming and Walton.

191. *TELLINA FRAGILIS* (pl. lvi. f. 14. and lx. f. 149.) *Lin.* Syst. 1117; *Chem.* vol. vi. f. 84; *Philip. Mol. Sic.* p. 28; *Poli. Test. Sic.* t. 15. f. 22, 24; *Petricola ochroleuca*, *Jam.*, no. 2; *Payr. Cat. Cors.* t. 1. f. 9, 10; *Hanl. Des. Cat.* p. 52. sup. t. 1. f. 6; *Psammobia fragilis*, *Turt. Biv. Brit.* p. 88. t. 7. f. 11, 12.

T. testâ ovato-trigonâ, solidiusculâ, subpellucidâ, valdè inæquilaterali, satis tumidâ, subimpolitâ, albidâ aut pallidè crocèâ, ad umbones flavescente aut crocèâ, striis remotiusculis elevatis concentricè asperâ; margine ventrali convexo, posticè sursum acclinato;

dorsali utrinque satis declivi, anticè convexo, posticè prope nates subretuso deindeque convexiusculo; latere antico rotundato, breviorè; postico obtusè acuminato et paululum subrostrato; natibus obtusis; flexurâ costâque umbonali obsoletis; superficie internâ plerumque crocèâ; dentibus satis magnis.

Var. Magis elongatâ, tumidâ; striis lamellosis; margine ventrali minus convexo; extremitate posticâ obtusè biangulatâ.

Ovate-trigonal, moderately solid, slightly pellucid, very inequilateral, rather tumid; almost devoid of glossiness, whitish with a tinge of yellow or saffron colour (which sometimes but rarely pervades the entire surface) upon the umbones; rough with concentric elevated rather distant striæ; ventral edge convex; dorsal slightly retuse on either side of the obtuse beaks, and then convex and rather sloping in front, scarcely convex but decidedly sloping behind; anterior side short and rounded; posterior obtusely acuminated and very slightly subrostrated; fold and flexure obsolete; inside usually of a saffron colour; teeth rather large, two diverging ones in one valve, a recurved one in the other.

Var. Elongated and not at all triangular, tumid; the striæ lamellar; ventral edge scarcely at all convex; hinder end obtusely biangulated.

Scarcely a *Tellina*, but with the teeth and general characters of so many usually included in that genus, that I know not how to separate it. From the figure in Delessert, the *Psammotœa Tarentina* of Lamarck is probably a variety of this shell.

Britain, Europe.

192. *TELLINA GUINAICA* (pl. lix. f. 130.) *Chemnitz*, vol. xi. p. 348. f. 1651, 1652, 1653; Dillw. Des. Cat. p. 69; *T. polygona*, *Gmel.*; Lam. no. 43.

*T. testâ* ovatâ, solidâ, ventricosâ, impolitâ, inæquilaterali, extus intusque albidâ, lineis rudioribus subremotis elevatis concentricè striatâ; margine ventrali magis minusve convexo; dorsali posticè subdeclivi et subrecto, anticè magis declivi et subincurvato; latere antico brevi, rotundato; latere postico rotundato-angulato aut obtusissimè cuneiformi, extremitate ejus semper rotundatâ; costâ umbonali subinconspicuâ; lunulâ distinctâ; ligamento infosso; dentibus magnis.

Ovate, solid, ventricose, dull, inequilateral, within and without whitish, roughened by coarse raised concentric and rather distant striæ; ventral edge more or less convex; dorsal edge straightish and not much sloping behind, more sloping and somewhat incurved in front; anterior side short and rounded; posterior somewhat

angulated below, or very obtusely wedge-shaped, the tip always rounded; fold not very distinct; ligament sunken; a distinct short lunule; teeth very large.

Guinea.

This is probably the *Abildgaardiana* of Spengler.

193. *TELLINA POLYGONA* (pl. lxii. f. 189.) *Chemnitz*, vol. vi. p. 91. f. 77; *Dillw. Des. Cat.* p. 77; *Hanl. Des. Cat.* p. 69. t. 3. f. 11; *T. multangula*, *Gmel.*, 3230; *Lam. no.* 42.

T. testâ ovato-oblongâ, solidâ, ventricosâ, inæquilaterali, impolitâ, extus intusque albidâ, lineis elevatis confertis concentricè striatâ, anticè breviorè, posticè subcuneiformi; margine ventrali subrecto; dorsali anticè paululum declivi, posticè longo subrecto declivi; extremitate anticâ obtusâ; posticâ attenuatâ et biangulatâ; flexurâ costâque umbonali perspicuis; ligamento subinfosso; dentibus magnis.

Ovate-oblong, solid, ventricose, inequilateral, not glossy, whitish within and without, roughened by coarse crowded concentric raised striæ, shorter in front, somewhat wedge-shaped behind; ventral edge straightish, but rendered wavy by two extremely obsolete and subcentral radiating ridges; anterior extremity obtuse, dorsal edge short and scarcely at all sloping; posterior tip narrow and biangulated, the ligamental margin long straightish and sloping; fold and flexure evident; teeth large.

Guinea.

194. *TELLINA BRUGUIERI* (pl. lxii. f. 192.) *Hanley*, *Zool. Proc.* 1844; *Enc. Mèth. t.* 231. f. 2.

T. testâ rotundato-trigonâ, solidâ, subæquilaterali, convexâ, impolitâ, extus intusque albidâ, sublævigatâ, (aut striis exilibus obsolete radiantibus strias concentricas tenuissimas decussantibus); margine ventrali convexo aut subarcuato; dorsali posticè elevatiore valdè declivi et paululum convexo, anticè arcuato et declivi; extremitate anticâ obtusâ; posticâ rotundatâ; natibus prominentibus, et anticè incumbentibus; flexurâ costâque umbonali obsolete; ligamento infosso; lunulâ parvâ; dentibus maximis.

Suborbicular, subtriangular, solid, nearly equilateral, convex, white within and without, dull, nearly smooth, when closely examined some of the adult specimens present the appearance of a most minute decussation, caused by the intersection of delicate obsolete radiating lines and very fine concentric striæ, which latter are chiefly apparent near the convex or subarcuated ventral margin; posterior dorsal edge abruptly sloping, slightly convex, and

more elevated than the arcuated and moderately sloping anterior; front extremity broad and rounded, hinder one obtuse; ligament sunken; fold and flexure obsolete; beaks prominent and inclining forward; teeth extremely large.

Isle of Panay, Philippines (Cuming).

195. *TELLINA ALTA* (pl. lxii. f. 200.) *Conrad*, Jour. Ac. N. S. Philad. vol. vii. p. 258; *Hanl. Des. Cat.* p. 71.

T. testâ obovali, subventricosâ, crassiusculâ, subinæquilaterali, impolitâ, concentricè et confertim rugosâ, extus intusque albidâ; margine ventrali subrecto; dorsali utrinque paululum declivi, anticè convexo, posticè recto; latere antico longiore, rotundato; extremitate posticâ obtusè biangulatâ; costâ umbonali latâ, planulatâ; natibus rectè incurvatis; ligamento infosso; dentibus angustis.

Oboval, slightly ventricose, thickish, slightly inequilateral, dull white, with crowded concentric wrinkles; ventral edge straightish; dorsal little sloping on either side, convex in front, rectilinear behind; anterior side rather the longer, rounded at its extremity; posterior side obtusely biangulated; umbonal ridge broad but flattened; beaks directly incurved; ligament sunken; inside white; the teeth narrow; the hinge margin broad.

California.

196. *TELLINA EPHIPPIMUM* (pl. lxxv. f. 244.) *Spengler*, loc. cit.

T. testâ *Lacunosa* simillimâ, sed valdè inæquivalvi et planè inæquilaterali; costâ umbonali angustissimâ, carinatâ.

Closely resembling *Lacunosa*, but decidedly inequivalve and inequilateral; the umbonal ridge narrow and carinated in the flatter valve, replaced by a linear indentation in the other.

East Indies (Spengler).

197. *TELLINA LACUNOSA* (pl. lxxv. f. 252.) *Chem.*, vol. iv. p. 92. f. 78; *Lam. no.* 39; *Enc. Mèth. t.* 290. f. 14; *Hanl. Des. Cat.* p. 69. t. 3. f. 12; T. papyracea, *Gmel.*

T. testâ obovali, ventricosâ, subinæquivalvi, subæquilaterali, tenuiusculâ, subpellucidâ, extus intusque candidâ, sublævigatâ, medio contorto-lacunosâ; margine ventrali subrecto sed contorto; dorsali antico subarcuato; ligamentali subdeclivi, subrecto; latere antico lato, rotundato; postico lato, obtuso; natibus prominentibus, incurvatis; ligamento subinfosso; costâ umbonali latâ, obtusâ.

Var. Testâ crassiore; parte superiore lateris postici partem inferiorem superante.

Oboval, rather thin, very slightly inequivalve, nearly inequilateral, ventricose, somewhat pellucid, white, nearly smooth or with delicate concentric wrinkles, which are chiefly apparent near the rounded anterior extremity; ventral edge nearly straight (as to convexity), but much twisted, owing to the concavity in front of the obtuse but prominent umbonal slope in one of the valves; dorsal edge scarcely sloping, arcuated in front, straightish behind; extremity of the hinder side (which is slightly the shorter) very obtusely biangulated; ligament rather sunken; beaks broad and directly incurved; teeth small.

Var. Thicker and coarser; a broader space from the umbonal ridge to the posterior end, which latter projects at its upper extremity.

Guinea.

198. *TELLINA SPECTABILIS* (pl. lxxv. f. 254.) *Hanley*, Zool. Proc. 1844.

T. testâ ovato-trapeziformi, subtenui, maximè inæquilaterali, ventricosâ, impolitâ, extus intusque albidâ; concentricè striatâ, striis elevatis, tenuibus, rugosis; margine ventrali paululum convexo; dorsali anticè convexo et declivi, posticè recto aut subincurvato et maximè declivi; extremitate lateris antici longioris obtusâ; extremitate posticâ truncato-cuneiformi, obtusè biangulatâ; flexurâ costâque umbonali conspicuis; ligamento magno, haud prominente; dentibus minimis.

Ovate, subtrapeziform, not solid, extremely inequilateral, ventricose, dull, white both internally and externally, with fine raised concentric wrinkles, which are rather distant near the beaks, but become closer towards the but little convex ventral margin; anterior side produced, its dorsal edge convex and sloping (in the young it is at first almost parallel with the ventral); its extremity obtuse or more rarely rounded; posterior side extremely short, its extremity truncato-cuneiform and obtusely biangulated, the ligamental edge abruptly sloping and straight or slightly incurved; fold and flexure prominent, the former subcarinated in the young; ligament large but not projecting; teeth minute.

Bay of Manilla and Isle of Siquijor.

199. *TELLINA DOMBEI* (pl. lxxii. f. 182, and pl. lxxiv. f. 222.) *Hanley*, Zool. Proc. 1844. — *Dombeyi* corr. in index.

T. testâ obovatâ aut ovali, subinæquivalvi, magis minusve so-

lidâ, subventricosâ, subimpolitâ, sublævigatâ, extus intusque albidâ, umbonibus aurantio-roseis; margine ventrali subrecto aut paululum convexo; dorsali antico vix declivi et prope nates prominulas subretuso, deinde magis minusve convexo; margine ligamentali declivi, subrecto; extremitate anticâ rotundatâ; latere postico brevior, obtusè cuneiformi; flexurâ subobsoletâ, (rarius distinctâ); ligamento infosso.

Var. Testâ majore, impolitâ, opacâ, concentricè subrugosâ, umbonibus haud erubescens; flexurâ costâque umbonali distinctis.

Varying in form from obovate to oval, more or less solid, rather ventricose, rarely at all glossy, whitish with the umbones reddish crimson, nearly smooth, (in full grown individuals with a few concentric striæ near the scarcely convex ventral margin, whose flexure is rarely distinct); anterior side the longer, sometimes greatly, at others scarcely so, its dorsal edge for about half its length from the rather prominent beaks somewhat retuse and nearly parallel to the lower edge, thence declining in a convex slope to the rounded termination; hinder extremity more or less obtusely wedge-shaped, the ligamental edge sloping strongly, and being more or less straight; ligament sunken; the inner disk red; one umbone projecting beyond the other.

Var. Larger, dull, opaque, the umbones not coloured, the surface somewhat wrinkled concentrically; fold and flexure distinct.

Panama, the variety from Tumbez.

200. *TELLINA ANGULATA* (pl. lkv. f. 250.) *Chemnitz*, vol. vi. p. 89. f. 74, 75; *Wood*, *Gen. Conch.* p. 146. t. 43. f. 1; *Schroet. Einl.* vol. ii. t. 7. f. 8; *Lutraria Tellinoides*, *Lam.*; *Deles.* t. 3. f. 5; *Hanl. Des. Cat.* p. 27. sup. t. 11. f. 33.

T. testâ subovali, inæquivalvi, solidiusculâ, inæquilaterali, convexâ aut subventricosâ, extus intusque albâ, sublævigatâ; margine ventrali subrecto aut convexiusculo; dorsali utrinque paululum declivi aut subdeclivi et subrecto, posticè elevatiore, anticè prope nates subincurvato; latere antico obtuso aut rotundato, longiore; extremitate posticâ obtusâ, subattenuatâ, aut obtusissimè biangulatâ; natibus prominentibus incurvatis, acutis; costâ umbonali satis distinctâ; ligamento parvo, angusto, infosso; dentibus obsoletis.

Somewhat oval, inequivalve, tolerably strong, moderately inequilateral, convex or subventricose, white both within and without, nearly smooth; ventral edge straightish or but slightly convex; anterior side the longer, its extremity rounded when the straightish



dorsal edge (which is slightly incurved near the beaks) is moderately sloping, when but little sloping obtuse; posterior extremity very obtusely biangulated, or attenuated and obtuse; beaks acute prominent and incurved; fold tolerably distinct; ligament small narrow and sunken; teeth obsolete.

Ceylon and Isle of Negros, Philippines; (Cuming).

This is not the *T. angulata* of Linnæus, which I believe to be a *Psammobia*.

201. *TELLINA GUBERNACULUM* (pl. lxii. f. 186.) *Hanley*, Zool. Proc. 1844.

T. testâ subovatâ, subinæquivalvi, tenui, compressiusculâ, lævi, extus intusque albidâ, valdè inæquilaterali; margine ventrali convexiusculo, anticè sursum acclinato; dorsali anticè magis minusve convexo et declivi, posticè recto brevi et subito declivi; extremitate lateris antici longioris rotundato-attenuatâ; extremitate posticâ brevissimâ, truncato-cuneiformi; flexurâ costâque umbonali obsolete; ligamento infosso; dentibus parvis.

Var. Testâ magis elongatâ; latere postico brevissimo; margine dorsali posticè maximè declivi, anticè subrecto.

Subovate, rather inequivalve, thin, somewhat compressed, smooth, whitish, very inequilateral; ventral edge moderately convex, sloping upwards in front; dorsal more or less convex and sloping anteriorly, but short straight and abruptly declining posteriorly; anterior side produced, its extremity attenuated and rounded; posterior side extremely short and truncato-cuneiform; fold and flexure obsolete; ligament sunken; teeth small.

Var. More elongated; hinder side extremely short; dorsal edge nearly straight in front, most abruptly sloping behind.

Real Lejos, Central America (Cuming).

202. *TELLINA TRUNCATA*. (pl. lxii. f. 198.) *Jonas*, Ab. und Bes. Conch. part 3. t. 5. f. 2.

T. testâ ovali-subtrigonâ, solidiusculâ, subinæquivalvi, subventricosâ, subimpolitâ, extus intusque albâ, sublævigatâ, valdè inæquilaterali; margine ventrali subrecto; dorsali anticè convexiusculo et declivi aut subdeclivi, posticè recto subelongato et valdè declivi; latere antico producto, ad extremitatem rotundato-attenuato; latere postico subcuneiformi, ad extremitatem angusto et subangulato; flexurâ subobsoletâ; ligamento magno, infosso; dentibus primariis distinctis.

Oval-subtriangular, tolerably strong, slightly inequivalve, rather ventricose, rather dull white, nearly smooth, very inequilateral;

ventral edge nearly straight; anterior side attenuated, produced and rounded, its dorsal edge sloping and tolerably convex; posterior side nearly wedge-shaped, the extremity narrow and subangulated, the dorsal edge straight, rather long in proportion and strongly sloping; flexure almost entirely obsolete; ligament large and unken; inside white; teeth distinct.

Bay of Manilla (Cuming).

Not unlike *Gubernaculum* but the hinder side in that species is less wedge-shaped, the tip being broader, and the ligamental edge decidedly shorter.

203. *TELLINA PELLUCIDA* (pl. lix. f. 118.) *Philippi*, Ab und. Bes. Conch. part 3. t. 5. f. 4.

T. testâ ovatâ aut obovatâ, tenui, inæquilaterali, convexâ, nitidâ, extus intusque niveâ, lævigatâ; margine ventrali convexo, posticè sursum acclinato; dorsali anticè convexo et paulo declivi, posticè convexiusculo et declivi; extremitate lateris antici longioris rotundatâ; extremitate posticâ obtusâ aut rotundatâ; natibus acutis, haud prominentibus; ligamento fulvo, infosso; costâ umbonali obsoletâ; dentibus angustis, minimis.

Var. Testâ abbreviatâ; margine dorsali utrinque magis declivi et minus convexo.

More or less ovate, thin, decidedly convex, inequilateral, pure glossy white both within and without, smooth, or with a few lines of growth; anterior side the longer, its extremity rounded, its dorsal edge convex and but little sloping; hinder side clearly the shorter, its dorsal edge sloping and but slightly convex, tip obtusely rounded; ventral edge convex (or even subarcuated) and inclining upward posteriorly; ligament yellow and sunken; beaks acute but not prominent; umbonal ridge obsolete; teeth minute.

Var. Abbreviated; the dorsal edges more sloping and less convex.

Bay of Manilla (Cuming).

204. *TELLINA CHINENSIS* (pl. lvi. f. 18.) *Hanley*, Zool. Proc. 1844.

T. testâ ovali, solidiusculâ, convexâ, subinæquilaterali, impolitâ, extus intusque candidâ, lævigatâ; margine ventrali subrecto; dorsali anticè convexiusculo et paululum declivi, posticè subrecto satisque declivi; extremitate posticâ obtusâ; latere antico longiore, rotundato; ligamento — ?; costâ umbonali obsoletâ.

Oval, tolerably strong, convex, slightly inequilateral, dull white, smooth; ventral edge nearly straight; anterior side the longer,

rounded at its extremity, its dorsal edge scarcely at all sloping and moderately convex; posterior extremity obtuse, the hinder dorsal edge moderately sloping and nearly straight; ligament —?; umbonal ridge obsolete; inside white.

China (Mus. Brit.).

205. *TELLINA FRIGIDA* (pl. lix. f. 119.) *Hanley, Zool. Proc.* 1844.

T. testâ ovali, solidiusculâ, convexâ, inæquilaterali, nitidâ, albidâ (intus candidâ), lævigatâ, utrinque rotundatâ; margine ventrali convexo; dorsali antico subdeclivi, convexo; latere postico planè breviorè; natibus inconspicuis; flexurâ costâque umbonali obsoletis; dentibus minimis.

Oval, tolerably strong, convex, inequilateral, glossy, whitish, smooth, rounded at each extremity; ventral edge convex; front dorsal convex and moderately sloping; posterior side decidedly the shorter; beaks not prominent; fold and flexure obsolete; teeth minute; inside white.

Kamskatchka.

206. *TELLINA GRANDIS* (pl. lxxv. f. 247.) *Hanley, Zool. Proc.* 1844. p. 187.

T. testâ ovali, subtrigonâ, solidâ, subinæquivalvi, lævi, convexâ, subimpolitâ, extus intusque albidâ, anticè rotundatâ; margine ventrali convexiusculo; dorsali utrinque declivi, posticè recto aut subrecto, anticè vix convexiusculo; extremitate lateris antici brevioris obtusâ; flexurâ costâque umbonali subobsoletis; ligamento magno

Subtriangularly oval, solid, slightly inequivalve, convex, smooth, rather of a dull white under an almost imperceptible epidermis, which is only visible near the but little convex ventral margin; anterior side clearly the longer, its extremity rounded, its dorsal edge but little convex and moderately sloping, ligamental edge straight, and strongly sloping; posterior termination obtuse; fold and flexure almost entirely obsolete; inside white, the hinge-plate strong, the teeth of a moderate size.

Tumbez, Peru.

207. *TELLINA SECTA* (pl. lxxv. f. 245, 248.) *Conrad, Jour. Ac. N. S. Philad.* vol. vii. p. 257. *Hanl. Des. Cat.* p. 67.

T. testâ solidiusculâ, inæquivalvi, nitidâ, extus intusque candidâ, lævigatâ; alterâ valvulâ convexâ, alterâ complanatâ; margine ventrali subarcuato; dorsali utrinque haud multum declivi,

et prope ligamentum magnum inciso; extremitate anticâ rotundatâ; costâ umbonali magis minusve conspicuâ; dentibus primariis distinctis; margine cardinali sub ligamento intus incrassato, superficie internâ impolitâ.

*Testâ seniore* obovali, æquilaterali; margine dorsali subhorizontali, convexiusculo; latere antico lato; extremitate posticâ obtusâ; margine postico angulum obtusum cum margine ventrali formante; ligamento maximo.

*Testâ juniore* sicut in seniore, sed ovali, inæquilaterali; margine dorsali antico subdeclivi; latere antico longiore, attenuato.

Var. *Testâ ovali*, inæquilaterali, margine dorsali antico subdeclivi; latere antico longiore, plerumque ad extremitatem attenuato; extremitate posticâ angulatâ; margine ventrali posticè subincurvato et magis sursum acclinato.

Inequivalve, tolerably strong, smooth, glossy, rounded in front; one valve convex, the other flattened; ventral margin subarcuated; dorsal edges not greatly sloping, near the large ligament (under which the hinge margin is internally thickened) with an incision; umbonal ridge more or less conspicuous; teeth distinct.

*Adult.* Oboval, equilateral, hinge-margin subhorizontal and slightly convex; anterior side broad; hinder extremity obtuse; posterior edge forming one obtuse angle with the ventral; ligament very large.

*Young.* Resembling the adult, but oval and inequilateral; the front dorsal edge moderately sloping; anterior side the longer, attenuated at its termination.

Var. Oval, inequilateral; front dorsal edge moderately sloping; anterior side the longer, and generally attenuated at its extremity; posterior end angulated, the hinder ventral edge being somewhat incurved and sloping upward.

California.

---

To this section belong, likewise, the *T. lateralis* and *brevifrons* of Say, which, however, I am unable to identify; and as the species have not been recognised by any subsequent conchologists, and are unknown even to the naturalists of the United States, I cannot, consequently, add to descriptions which will not exactly apply to any Tellens I am acquainted with.

“*TELLINA LATERALIS*, Say, Journ. Ac. Nat. Sci. Philad.

Transversely subovate, subequilateral; posterior (anterior) mar-

gin regularly rounded, anterior (posterior) margin beaked, the rostrum turned to the left and slightly gaping, ligament slope straight; basal margin regularly arcuated, a little contracted near the beak; valves whitish, often tinged with ferruginous, with small concentric wrinkles and slight waves; within white, a little sculptured by the external waves; no lateral teeth, two cardinal in one valve, and one with another hardly elevated filiform one in the other.

South coast of the United States."

*TELLINA BREVIFRONS* Say, Am. Conch. t. 64. f. 1.

Ovate; thin, fragile, not very convex, white, tinged particularly on the umbo with pale dull fulvous; with transverse (concentric) slender striæ, and in a particular light obsolete longitudinal (radiating) striæ are visible; beaks much anterior (posterior) to the middle, forming an angle in consequence of the anterior and posterior hinge-margin being rectilinear to a considerable distance, the latter parallel to the base; anterior (posterior) side short and abrupt, rounded at the tip, and with a submarginal undulation; posterior (anterior) side more than as long again as the anterior (posterior), rounded at tip; within much more deeply coloured with fulvous! cardinal teeth, two in the left and one in the right valve; lateral teeth none."

The shell is represented in the drawing as nearly an inch long, of a pale rose-colour (!), and of a subelliptical contour, a shape neither agreeing with the description nor with the further statement that "its outline corresponds with *T. donacina*, but is more convex" (!). I regard it as a very doubtful and ill-defined species. A single specimen only has been met with by Dr. Ravenel in South Carolina.

---

The *T. linteæ* of Conrad, which, being described in company with recent shells, I had introduced into my 'Descriptive Catalogue,' I have since ascertained to be fossil.

## TELLINA.

## LIST OF PLATES LVI. TO LXVI.

- | PLATE LVI. |                                     | Fig.                                   |  |
|------------|-------------------------------------|----------------------------------------|--|
| Fig.       | 1. <i>Tel. Guildingii</i> , Sp. 16. | 37, 38. <i>Tel. carnaria</i> , Sp. 69. |  |
|            | 2, 3. — <i>robusta</i> , 56.        | 39. — <i>splendida</i> , 67.           |  |
|            | 4. — <i>pulchella</i> , 17.         | PLATE LVII.                            |  |
|            | 5. — <i>virgulata</i> , 18.         | 40. — <i>spinosa</i> , 77.             |  |
|            | 6. — <i>distorta</i> , 19.          | 41. — <i>pumila</i> , 105.             |  |
|            | 7. — <i>pudica</i> , 46.            | 42. — <i>virgo</i> , 118.              |  |
|            | 8. — <i>semen</i> , 51.             | 43. — <i>crystallina</i> , 89.         |  |
|            | 9. — <i>casta</i> , 57.             | 44. — <i>inæqualis</i> , 104.          |  |
|            | 10. — <i>balaustina</i> , 58.       | 45. — <i>ostracea</i> , 87.            |  |
|            | 11. — <i>obliquilineata</i> , 59.   | 46, 47. — <i>lineata</i> , 22.         |  |
|            | 12. — <i>donacina</i> , 20.         | 48. — <i>subtruncata</i> , 83.         |  |
|            | 13. — <i>umbonella</i> , 150.       | 49. — <i>mera</i> , var. 101.          |  |
|            | 14. — <i>fragilis</i> , 191.        | 50. — <i>Corbuloides</i> , 108.        |  |
|            | 15. — <i>carnicolor</i> , 75.       | 51. — <i>culter</i> , 107.             |  |
|            | 16. — <i>bimaculata</i> , 52.       | 52. — <i>felix</i> , 110.              |  |
|            | 17. — <i>Senegalensis</i> , 68.     | 53. — <i>Hiberna</i> , 112.            |  |
|            | 18. — <i>Chinensis</i> , 204.       | 54. — <i>hilaris</i> , 111.            |  |
| 19, 20. }  | — <i>bimaculata</i> , 52.           | 55, 56. — <i>Philippinarum</i> , 106.  |  |
| 21, 22. }  |                                     | 57. — <i>Corbuloides</i> , 108.        |  |
| 23, 24. —  | <i>ampullacea</i> , 60.             | 58. — <i>inæquistriata</i> , 31.       |  |
|            | 25. — <i>tulipa</i> , 21.           | 59. — <i>tenera</i> , 114.             |  |
|            | 26. — <i>Gouldii</i> , 92.          | 60. — <i>polita</i> , 113.             |  |
|            | 27. — <i>decora</i> , 120.          | 61. — <i>Cycladiformis</i> , 102.      |  |
| 28, 29. —  | <i>flexuosa</i> , 72.               | 62. — <i>fabula</i> , 123.             |  |
|            | 30. — <i>pisiformis</i> , 71.       | 63. — <i>juvenilis</i> , 125.          |  |
|            | 31. — <i>Cyrenoidea</i> , 66.       | 64. — <i>subrosea</i> , 146.           |  |
|            | 32. — <i>Lucinoidea</i> , 53.       | 65. — <i>similis</i> , 119.            |  |
|            | 33. — <i>nux</i> , 54.              | 96. — <i>Deshayesii</i> , 2.           |  |
|            | 34. — <i>pinguis</i> , 55.          | 97. — <i>vestalis</i> , 132.           |  |
| 35, 36. —  | <i>lineata</i> , 22.                | 68. — <i>Valtonis</i> , 116.           |  |

- Fig.  
69. *Tel. truncatula*, Sp. 131.  
70. — *emarginata*, 130.  
71. — *lux*, 126.

## PLATE LVIII.

72. — *Cumingii*, 3.  
73. — *Cumana*, 148.  
74, 75. — *Antonii*, 5.  
76. — *aurora*, 153.  
77. — *verrucosa*, 6.  
78, 79. — *crucigera*, 4.  
80. — *inaequistriata*, 31.  
81, 82. — *tenuis*, 124.  
83. — *formosa*, 163.  
84. — *vernalis*, 127.  
85. — *lilium*, 158.  
86, 87. — *lanceolata*, 133.  
88. — *iridescens*, 122.  
89. — *punicea*, 33.  
90. — *coccinea*, 143.  
91. — *eburnea*, 36.  
92. — *rhomboides*, 162.  
93. — *rhodon*, 144.  
94. — *ancilla*, 160.  
95. — *assimilis*, 157.  
96, 97. — *rhomboides*, 162.  
98. — *lucerna*, 152.  
99. — *Burneti*, 90.  
100. — *conspicua*, 136.

## PLATE LIX.

101. — *nitida*, 168.  
102. — *Bodegensis*, 161.  
103. — *lutea*, 165.  
104. — *exilis*, 117.  
105. — *rhodora*, 109.  
106. — *micans*, 170.  
107. — *undulata*, 174.  
108. — *dispar*, 164.

- Fig.  
109, 110. *Tel. solidula*, Sp. 189.  
111. — *Souleyeti*, 169.  
112. — *Fabricii*, 188.  
113, 114. — *dispar*, 164.  
115. — *proxima*, v. *macroma*, 180.  
116. — *scalpellum*, 173.  
117. — *fusca*, 186.  
118. — *pellucida*, 203.  
119. — *frigida*, 205.  
120. — *inconspicua*, 187.  
121. — *Balthica*, 185.  
122. — *cygnus*, 175.  
123. — *inornata*, 184.  
124. — *planissima*, 140.  
125. — *cuspis*, 172.  
126. — *obliquilineata*, 59.  
127. — *decora*, 120.  
128. — *deltoidalis*, 62.  
129. — *plebeia*, 149.  
130. — *Guinaica*, 192.

## PLATE LX.

131. — *Antonii*, 5.  
132. — *fimbriata*, 73.  
133. — *Owenii*, 25.  
134. — *Psammotella*, 129.  
135. — *asperrima*, 9.  
136. — *insculpta*, 128.  
137. — *ala*, 176.  
138. — *sulcata*, 10.  
139. — *perplexa*, 86.  
140. — *regia*, 35.  
141. — *acuminata*, 139.  
142. — *incarnata*, 115.  
143. — *nobilis*, 151.  
144. — *sincera*, 70.  
145. — *irus*, 190.  
146. — *ala*, 176.  
147. — *ovalis*, 142.

Fig.

148. *Tel. staurella*, Sp. 15.  
 149. — *fragilis*, 191.  
 150. — *triangularis*, 138.  
 151. — *plebeia*, 149.  
 152. — *prora*, 39.  
 153. — *rubescens*, 38.  
 154. — *punica*, 33.  
 155. — *imbellis*, 103.

## PLATE LXI.

156. — *gargadia*, 76.  
 157. — *rostrata*, 1.  
 158. — *Timorensis*, 135.  
 159. — *alternata*, 37.  
 160. — *pristis*, 84.  
 161. — *striata*, 34.  
 162, 163. — *vulsella*, 27.  
 164. — *albinella*, 29.  
 165. — *pulcherrima*, 8.  
 166. — *albinella*, 29.  
 167. — *hyalina*, 134.  
 168. — *laceridens*, 40.  
 169. — *crassa*, 78.  
 170. — *rosea*, 30.  
 171. — *staurella*, 15.  
 172. — *Timorensis*, 135.  
 173. — *crassa*, 78.  
 174. — *planata*, 99.  
 175. — *striatula*, 61.  
 176. — *laceridens*, 40.  
 177. — *pectrum*, 79.

## PLATE LXII.

178. — *crucigera*, 4.  
 179. — *Brasiliana*, 45.  
 180. — *Mars*, 94.  
 181. — *gargadia*, 76.  
 182. — *Dombeï*, 199.  
 183. — *calcareo*, 181.

Fig.

184. *Tel. decussata*, Sp. 74.  
 185. — *Capsoides*, 85.  
 186. — *gubernaculum*, 201.  
 187. — *lyra*, 91.  
 188. — *miles*, 171.  
 189. — *polygona*, 193.  
 190. — *Cayennensis*, 178.  
 191. — *plicata*, 88.  
 192. — *Bruguieri*, 194.  
 193. — *alba*, 179.  
 194. — *purpurascens*, 141.  
 195. — *candida*, 155.  
 196. — *nymphalis*, 177.  
 197. — *planissima*, 140.  
 198. — *truncata*, 202.  
 199. — *elongata*, 156.  
 200. — *alta*, 195.

## PLATE LXIII.

201. — *magna*, 96.  
 202. — *perna*, 28.  
 203. — *sinuosa*, 137.  
 204. — *virgata*, 12.  
 205. — *Sowerbii*, 47.  
 206. — *princeps*, 32.  
 207. — *interrupta*, 11.  
 208. — *chloroleuca*, 49.  
 209. — *Tongana*, 42.  
 210. — *chloroleuca*, 49.  
 211. — *strigosa*, 98.  
 212. — *virgata*, 12.  
 213. — *rufescens*, 167.  
 214. — *jubar*, 13.  
 215. — *Pharaonis*, 26.  
 216. — *marginalis*, 14.  
 217. — *perna*, 28.  
 218. — *Madagascariensis*, 41.  
 219. — *perna*, 28.



Fig.  
221, 220. *Tel. radiata*, Sp. 44.

## PLATE LXIV.

222. — *Dombei*, 199.  
 223. — *Galathæa*, 154.  
 224. — *nasuta*, 182.  
 225. — *lævigata*, 50.  
 226. — *remies*, 65.  
 227. — *lævigata*, 50.  
 228. — *discus*, 64.  
 229. — *deltoidalis*, 62.  
 230. — *fausta*, 63.  
 231. — *rastellum*, var. 7.  
 232. — *discus*, 64.  
 233. — *rugosa*, 82.  
 234. — *fausta*, 63.  
 - 235. — *scobinata*, 80.  
 - 236. — *lingua-felis*, 81.  
 237. — *lata*, 48.  
 238. — *rugosa*, 42.

## PLATE LXV.

239. — *magna*, 96.  
 240. — *Sol*, 97.  
 241. — *elegans*, 43.  
 242. — *rastellum*, 7.  
 243. — *edentula*, 183.

Fig.

244. *Tel. ephippium*, Sp. 196.  
 245. — *secta*, 207.  
 246. — *Columbiensis*, 166.  
 247. — *grandis*, 206.  
 248. — *secta*, 207.  
 249. — *lutca*, 165.  
 250. — *angulata*, 200.  
 251. — *Listeri*, 103.  
 252. — *lacunosa*, 197.  
 253. — *foliacea*, 95.  
 254. — *spectabilis*, 198.

## PLATE LXVI.

- 255. — *rubicunda*, 93.  
 256. — *serrata*, 44. 2 /  
 257. — *margaritosa*, 147.  
 258. — *laccridens*, var. 40.  
 259. — *donacina*, v. *Lantivyi*, 20.  
 260. — *decora*, 120.  
 261. — *staurella*, v. *scalaris*, 15.  
 262. — *Oudardi*, 145.  
 263. — *petalum*, p. 317, note.  
 264. — *proxima*, 180.  
 - 265. — *incarnata*, 115.  
 266. — *tenta*, 159.  
 267. — *iris*, 121.  
 268. — *mera*, 101.

## ALPHABETICAL INDEX.


*acuminata*, Sp, 139.  
*ala*, 176.  
*alba*, 179.  
*albinella*, 29.  
*alta*, 195.  
 - *alternata*, 37.  
*ampullacea*, 60.

*ancilla*, Sp. 160.  
*angulata*, 200.  
*Antonii*, 5.  
*assimilis*, 157.  
*asperrima*, 9.  
*aurora*, 153.  
*balaustina*, 58.


- Balthica, Sp. 185.  
 -bimaculata, 52.  
 Bodegensis, 161.  
 Brasiliana, 45.  
 Brugueri, 194.  
 Burneti, 90.  
 calcarea, 181.  
 -candida, 155.  
 Capsoides, 85,  
 -carnaria, 69.  
 carnicolor, 75.  
 casta, 57.  
 Cayennensis, 178.  
 Chinensis, 204.  
 chloroleuca, 49.  
 coccinea, 143.  
 Columbiensis, 166.  
 conspicua, 136.  
 Corbuloides, 108.  
 crassa, 78.  
 crucigera, 4.  
 crystallina, 89. W. 3.  
 culter, 107.  
 Cumana, 148.  
 Cumingii, 3.  
 cuspis, 172.  
 Cycladiformis, 102.  
 cygnus, 175.  
 Cyrenoidea, 66.  
 decora, 120.  
 decussata, 74.  
 deltoidalis, 62.  
 Deshayesii, 2.  
 discus, 64.  
 -dispar, 164.  
 distorta, 19.  
 Dombi, 199.  
 donacina, 20.  
 eburnea, 36.  
 -edentula, Sp. 183.  
 elegans, 43.  
 elongata, 156.  
 emarginata, 130.  
 ephippium, 196.  
 exilis, 117.  
 Fabricii, 188.  
 -fabula, 123.  
 -fausta, 63. W. 5.  
 felix, 110.  
 fimbriata, 73.  
 flexuosa, 72.  
 -foliacea, 95.  
 formosa, 163.  
 fragilis, 191.  
 frigida, 205.  
 fusca, 186.  
 Galathæa, 154.  
 gargadia, 76.  
 gelida, 23.  
 Gouldii, 72.  
 grandis, 206.  
 gubernaculum, 201.  
 Guildingii, 16.  
 Guinaica, 192.  
 Hiberna, 112.  
 hilaria, 111.  
 hyalina, 134.  
 imbellis, 100.  
 inæqualis, 104.  
 inæquistriata, 31.  
 incarnata, 115.  
 inconspicua, 187.  
 -inornata, 184.  
 insculpta, 128.  
 -interrupta, 11. W. 5.  
 iridescens, 122.  
 iris, 121.  
 irus, 190.

- jubar, Sp. 13.
- juvenilis, 125.
- laceridens, 40.
- lacunosa, 197.
- lævigata, 50. *W. S.*
- lanceolata, 133.
- lata, 48.
- lilium, 158.
- lingua-felis, 81.
- lineata, 22. *W. S.*
- Listeri, 103.
- lucerna, 152.
- Lucinoides, 53.
- lutea, 165.
- lux, 126.
- lyra, 91.
- Madagascariensis, 41.
- magna, 96. *W. S.*
- margaritina, 147.
- marginalis, 14.
- Mars, 94.
- mera, 101.
- micans, 170.
- miles, 171.
- ✓ nasuta, 182.
- nitida, 168.
- nobilis, 151.
- nux, 54.
- nymphalis, 177.
- obliquilineata, 59.
- ostracea, 87.
- Oudardi, 145.
- ovalis, 142.
- Owenii, 25.
- pellucida, 203.
- perna, 28.
- perplexa, 86.
- petalum, p. 317, note.
- Pharaonius, 26.
- Philippinarum, Sp. 106.
- pinguis, 55.
- pisiformis, 71.
- planata, 99.
- planissima, 140.
- plebeia, 149.
- plectrum, 79.
- plicata, 88.
- polita, 113.
- polygona, 193.
- princeps, 32.
- pristis, 84.
- prora, 39.
- proxima, 180.
- Psammotella, 129.
- pudica, 46.
- pulchella, 17. *W. S.*
- pulcherrima, 8.
- pumila, 105.
- punicea, 33. *W. S.*
- purpurascens, 141.
- radiata, 44.
- rastellum, 7.
- regia, 35.
- remies, 65.
- rhodon, 144.
- rhodora, 109.
- rhomboides, 162.
- robusta, 56.
- rosea, 30.
- rostrata, 1.
- rubescens, 38.
- rubicunda, 93.
- rufescens, 167. *W. S.*
- rugosa, 82.
- scalpellum, 173.
- scobinata, 80.
- secta, 207.
- semen, 51.


- Senegalensis, Sp. 68.  
 serrata, 24.  
 similis, 119.  
 sincera, 70.  
 sinuata, 137.  
 sol, 97.  
 solidula, 189.  
 Souleyeti, 169.  
 Sowerbii, 47.  
 spectabilis, 198.  
 spinosa, 77.  
 splendida, 67.  
 staurella, 15.  
 striata, 34.  
 striatula, 61.  
 strigosa, 98.  
 subrosea, 146.  
 subtruncata, 83.  
 sulcata, 10.  
 tenera, Sp. 114. *WCS*  
 tenta, 159.  
 tenuis, 124.  
 Timorensis, 135.  
 Tongana, 42.  
 triangularis, 138.  
 truncata, 202.  
 truncatula, 131.  
 tulipa, 21.  
 umbonella, 150.  
 undulata, 174.  
 Valtonis, 116.  
 vernalis, 127.  
 verrucosa, 6.  
 vestalis, 132.  
 -virgata, 12.  
 virgo, 118.  
 virgulata, 18.  
 -vulsella, 27.




1




2



3



4



5



6



7



8




9




10




11




12




13




14



15



16



17



18



19



20




21



22



23



24



25



26



27




28



29



30



31




32




33



34




35




36



37




38



39


*C. G. S. S. S. S. S.*






































































## MONOGRAPH OF THE GENUS

LINGULA, *Brug.*

BY G. B. SOWERBY, F.L.S., ETC.

*Char. Gen.* Testa tenuis, subæquivalvis, æquilateralis, ovato-elongata, depressa, apice subtruncato, basi subacutâ, pediculo affixâ. Impressiones musculares in utrâque valvâ quatuor, ad utrumque latus duæ, altera basalis, altera subcentralis.

Shell thin, consisting for the most part of a horny epidermis, lined with a small quantity of calcareous matter within; its general form when the valves are closed is an elongated oval, compressed and much resembling a duck's bill, (whence the specific appellation of the first known species). The two valves are very nearly equal, the dorsal being somewhat thicker and more acuminate at the base. When in a state of rest, the two valves are closed at the edges in all parts. There is neither ligament nor hinge, the two valves being held together by the adductor muscles. Muscular impressions several in each valve, arranged in pairs, one pair nearly central, another toward the posterior part, and a third, small and double, close to the posterior apex. The shell is attached by a peduncle which is fixed within the apex of the dorsal valve.

Of recent species seven are here described and figured, more than double that number are known in a fossil state, as characterizing the various secondary and tertiary deposits.

## DESCRIPTIONS.

1. LINGULA ANATINA (pl. lxvii. f. 1, 2, 3, 9, 10.) *Lam.*

L. testâ oblongâ, compressâ, longitudinaliter substriatâ, apicis basisque valvarum latitudine æquali, apice subtrilobo, valvâ ventrali medio intus longitudinaliter incrassatâ.

Shell oblong, compressed, slightly striated longitudinally, equal

in width at both ends; apex somewhat trilobate; ventral valve longitudinally thickened near the centre within.

Fig. 1 represents the usual variety with its peduncle. Fig. 2, inside of the dorsal valve. Fig. 3, inside of the ventral valve. Fig. 9, a variety of a brown colour, found in sand at low water, at the Island of Siquijor, by Mr. Cuming. Fig. 10, a dark coloured variety.

From the Indian Ocean and the Moluccas, H. Cuming.

2. *LINGULA HIANS* (pl. lxvii. f. 4.) *Swains.*

L. testâ elongato-oblongâ, tenuissimâ, angustiori, prope basin paulatim attenuatâ, convexiusculâ.

Shell of a lengthened oblong form, very thin and rather narrow, slightly attenuated and rather convex toward the base.

Distinguished from *L. anatina* by being much thinner and of a more delicate green colour; it is also slightly and gradually attenuated toward the base; it only gapes at the extremities when the valves are separated from the animal and dried.

3. *LINGULA AUDEBARDI* (pl. lxvii. f. 5.) *Brod. Zool. Proc.* 1833. p. 100.

L. testâ elongato-oblongâ, glabrâ, pallidè flavâ, viridi transversim pictâ, margine antico rotundato, viridi.

"The rounded anterior edge of this shell is given and the transverse lines of that colour are produced by the progressive increase of the shell which is small and parchment like. In all the dried specimens the thin anterior edge is contracted into a square form, but in its natural state this edge is rounded. A general contraction moreover gives the dried shells a narrower and more ventricose character than they really possess, and the remains of the *cilia* of the mantle give to their edges a bearded appearance."—*Brod.*

Found by Mr. Cuming in hard coarse sand at about half tide, from four to six inches below the surface, at the Island of Punam in the Bay of Guayaquil.

4. *LINGULA SEMEN* (pl. lxvii. f. 11.) *Brod. l. c.*

L. testâ ovato-oblongâ, crassiusculâ, planâ, albidâ, lævissimâ, politâ, margine antico rotundato.

Shell ovate-oblong, thickish, flattened, whitish, very smooth and polished; anterior margin rounded.

"This shell is much firmer than is usual to *Lingula*, so firm indeed as not to have contracted at all in drying. In size and appearance it bears a near resemblance to a melon seed."—*Brod.*

Dredged off the Isle of Plata, West Columbia; H. Cuming.

5. *LINGULA TUMIDULA* (pl. lxvii. f. 7.) *Reeve, Zool. Proc.* 1841. p. 100.

L. testâ tenui, corneâ, subquadratâ, latâ, rufescente-olivaceâ, radiatim irregulariter striatâ, apice subtrilobo.

Broader in proportion than any other known species; it is somewhat narrower toward the base than at the apex, and it is irregularly and somewhat radiately striated from the base to the broad and subtrilobate apex.

New Holland.

*Lingula compressa* of Reeve is only a badly preserved specimen of this species. Masbate, H. Cuming.

6. *LINGULA OVALIS* (pl. lxvii. f. 8.) *Reeve, Zool. Proc.* 1841. p. 100.

L. testâ oblongo-ovali, subcompressâ, tenui, anticè rotundatâ.

Distinguished from other species by its regularly oblong-oval shape, both ends being nearly similar.

Sandwich Islands, H. Cuming.

7. *LINGULA ALBIDA* (pl. lxvii. f. 6.) *Hinds, Moll. Voy. of the Sulphur*, p. 71. t. 12. f. 4.

L. testâ oblongâ, lævi, complanatâ, anticè truncatâ, ubique albidâ; pediculo brevi, cylindræo.

"This species is distinguished by its uniform dead white colour, and very short pedicle, which does not exceed half the length of the shelly portion."—*Hinds*,


Bay of Magdalena, California.

---

#### EXPLANATION OF PLATE LXVII.

1. 2. 3. *Lingula anatina*, Sp. 1.
4. ——— *hians*, 2.
5. ——— *Audebardi*, 3.
6. ——— *albida*, 7.
7. ——— *tumidula*, 5.
8. ——— *ovalis*, 6.
9. ——— *anatina*, var. 1.
10. ——— *anatina*, var. 1.
11. ——— *semen*, 4.





THE HISTORY OF THE  
NATIVE PLANTS OF THE  
WEST INDIES  
BY  
JAMES SMITH  
M.D.  
OF THE COLLEGE OF PHYSICIANS IN GREAT BRITAIN  
AND OF THE SOCIETY OF PHYSICIANS IN FRANCE  
AND OF THE SOCIETY OF PHYSICIANS IN ITALY  
AND OF THE SOCIETY OF PHYSICIANS IN SPAIN  
AND OF THE SOCIETY OF PHYSICIANS IN PORTUGAL  
AND OF THE SOCIETY OF PHYSICIANS IN GREECE  
AND OF THE SOCIETY OF PHYSICIANS IN TURKEY  
AND OF THE SOCIETY OF PHYSICIANS IN PERSIA  
AND OF THE SOCIETY OF PHYSICIANS IN INDIA  
AND OF THE SOCIETY OF PHYSICIANS IN CHINA  
AND OF THE SOCIETY OF PHYSICIANS IN JAPAN  
AND OF THE SOCIETY OF PHYSICIANS IN SIBERIA  
AND OF THE SOCIETY OF PHYSICIANS IN RUSSIA  
AND OF THE SOCIETY OF PHYSICIANS IN POLAND  
AND OF THE SOCIETY OF PHYSICIANS IN PRUSSIA  
AND OF THE SOCIETY OF PHYSICIANS IN AUSTRIA  
AND OF THE SOCIETY OF PHYSICIANS IN HUNGARY  
AND OF THE SOCIETY OF PHYSICIANS IN ITALY  
AND OF THE SOCIETY OF PHYSICIANS IN FRANCE  
AND OF THE SOCIETY OF PHYSICIANS IN GREAT BRITAIN



THE HISTORY OF THE  
NATIVE PLANTS OF THE  
WEST INDIES  
BY  
JAMES SMITH  
M.D.  
OF THE COLLEGE OF PHYSICIANS IN GREAT BRITAIN  
AND OF THE SOCIETY OF PHYSICIANS IN FRANCE  
AND OF THE SOCIETY OF PHYSICIANS IN ITALY  
AND OF THE SOCIETY OF PHYSICIANS IN SPAIN  
AND OF THE SOCIETY OF PHYSICIANS IN PORTUGAL  
AND OF THE SOCIETY OF PHYSICIANS IN GREECE  
AND OF THE SOCIETY OF PHYSICIANS IN TURKEY  
AND OF THE SOCIETY OF PHYSICIANS IN PERSIA  
AND OF THE SOCIETY OF PHYSICIANS IN INDIA  
AND OF THE SOCIETY OF PHYSICIANS IN CHINA  
AND OF THE SOCIETY OF PHYSICIANS IN JAPAN  
AND OF THE SOCIETY OF PHYSICIANS IN SIBERIA  
AND OF THE SOCIETY OF PHYSICIANS IN RUSSIA  
AND OF THE SOCIETY OF PHYSICIANS IN POLAND  
AND OF THE SOCIETY OF PHYSICIANS IN PRUSSIA  
AND OF THE SOCIETY OF PHYSICIANS IN AUSTRIA  
AND OF THE SOCIETY OF PHYSICIANS IN HUNGARY  
AND OF THE SOCIETY OF PHYSICIANS IN ITALY  
AND OF THE SOCIETY OF PHYSICIANS IN FRANCE  
AND OF THE SOCIETY OF PHYSICIANS IN GREAT BRITAIN

PREPARING FOR PUBLICATION,

*In Numbers, profusely illustrated, price 2s. 6d. each, plain; 3s. 6d. coloured, size royal 8vo.*

---

# ELEMENTS OF CONCHOLOGY,

COMPRISING

**The Physiological History of Shells**

AND

**THEIR MOLLUSCOUS INHABITANTS,**


**THEIR GEOGRAPHICAL DISTRIBUTION, HABITS, CHARACTERS, AFFINITIES  
ARRANGEMENT, AND ENUMERATION OF SPECIES.**

BY

**LOVELL REEVE, A.L.S. ETC.**

AUTHOR OF THE 'CONCHOLOGIA ICONICA'.

---



*Voluta angulata.*

THE object of the present work is to invest the Science of Conchology with a high and interesting character, in an elementary form; to introduce in an explanatory and popular style, what has almost been confined (except in the admirable example begun by Mr. Gray in 'Turton's British Shells') to the regions of physical anatomy; and to ennoble the spirit of collecting shells, by inducing a love of enquiry into the nature and habits of their molluscos inhabitants. Shells are too frequently collected in the present day as mere objects of fancy or ornament, fascinating the amateur by their varieties of form and colour, without exciting the least enquiry as to their nature or zoological character. The Conchologist should look upon his shells, in the

absence of the living parts, as the medals of a creation whose history may to a great extent be ascertained by their structure and animal-impressions; and to exhibit their connection it is only necessary to present him with a few of the typical kinds with the animal *in situ*. Many erroneous notions have prevailed in regard to this most interesting science, for want of properly estimating the very important relation between shells and their molluscous inhabitants. Collectors are still prejudiced in favour of the antiquated notions of Linnæus. They arrange their shells according to the varieties of form and external aspect, forgetting, that they are merely a collection of skeletons, the calcified portions of animals possessing an organization far superior to that of the Bee, and a thousand creatures whose structure and sagacity of habits have so often excited their admiration.

It is proposed in the "ELEMENTS OF CONCHOLOGY" to give an account, not of the anatomical detail of the Mollusca, but of their habits, affinities, and mode of producing shells, their structure, geographical distribution, and systematic arrangement, with the names of all the species known up to the time of publication; accompanied by a characteristic shell-figure of every genus, and one of a shell with its living animal in illustration of each particular group.

The drawings of the shells with animals will be mainly derived from those magnificent, but almost inaccessible, Zoologies of Surveying Expeditions published by the French Government, including that by M. Lesson in the 'Voyage de la Coquille', by M. M. Quoy and Gaimard in the 'Voyage de l'Astrolabe', by M. D'Orbigny in the 'Voyage dans l'Amerique Méridionale' and by M. Deshayes in the 'Mollusques de l'Algérie'; and the author trusts by a popular generalization of the facts at present veiled in physiological obscurity, to give the elementary study of Conchology a more legitimate tone, and make it as easy as it is full of scientific interest.

To expect a work of this nature to become popular it must be well illustrated, and it must be cheap; and the author has assuredly the means of accomplishing these objects with peculiar facility in the possession of a well-organized printing establishment.

It is fully expected that the Work will not exceed about twelve numbers, and as a guarantee for its completion it may be added that Mr. G. B. Sowerby Jun. has been for some time occupied in engraving the figures upon copper-plate, of which between two and three hundred are already finished.

The first Number will be published in the course of January.


\*\* Subscribers are earnestly solicited to favour the Publishers with their names as early as possible.

LONDON :

REEVE, BROTHERS, KING WILLIAM STREET, STRAND ;  
 FORTIN, MASSON ET CIE., PLACE DE L'ECOLE DE MEDECINE, PARIS.










Carded







SMITHSONIAN INSTITUTION LIBRARIES



3 9088 00555 9539